THE

NAUTILUS

A MONTHLY JOURNAL
DEVOTED TO THE INTERESTS OF
CONCHOLOGISTS

VOL. XXVI.

MAY, 1912, to APRIL, 1913.

EDITORS AND PUBLISHERS:

H. A. PILSBRY, Curator of the Department of Mollusca, Academy of Natural Sciences
PHILADELPHIA.

C. W. JOHNSON, Curator of the Boston Society of Natural History,
BOSTON.

INDEX

TO

THE NAUTILUS, XXVI.

INDEX TO SUBJECTS, GENERA AND SPECIES.

Abnormal shells (Pl. V) 38
Agriolimax, Feeding habits of 108
Alabama, Additions to the list of Mollusca from Monte Sano 14
Amnicola winkleyi Pils., n. sp. (Pl. I, figs. 9, 10) . . 1
Ancellaria castanea Sowb. 77
Anculosa minor Hinkley, n. sp. (Pl. I, figs. 7, 8) . . 47
Annularia ramsdeni Pils. & Hend., n. sp. . . . 42
Annularia pseudalatum Torre, n. sp. 43
Aplexa hypnorum Linné 64
Ashmunella ashmuni Dall 81
Ashmunella thomsoniana, Pecos River, N. M. . . . 69
Baldwin, D. D. (obituary and portrait) . . . 48, 82
Bermuda shells 12
Bifidaria contracta 17
Bifidaria pellucida hordeacella (Pl. II) . . 17, 21, 31
Bifidaria pellucida parvidens 81
Boston Malacological Club 144
Bryant, F. W. (obituary) 119
Buccinum solutum Herm.=Chrysodomus heros Gray . 46
Buccinum stromboides Herm. 47
California, Two new land shells from 37
Campeloma decisum Say, reversed 41
Cannibal snails 132
Carocolus sagemon var. semiaperta 106
Carychium minimum at Quincy, Mass. 24
Cepolis alauda Fer. 2
Cerion pinerium Dall 112
Chondropoma carenasense Pils. & Hend., n. sp. . . 44
Chondropoma wilcoxi Pils. & Hend., n. sp. . . . 45
Chondropoma dissolutum. 112
Chondropoma picta Pfr. 104
Cincinnatia in the Delaware drainage 96
Circinaria concava from Sheboygan County, Michigan . 3
Colombia, Land shells of Carthagena 84
Columbella misera Sowb. var. californica Rve. (Pl. IV, fig. 15). 76
Columbella picta Reeve (Pl. IV, fig. 13) 76
Columbella propinqua E. A. Smith (Pl. IV, fig. 14). . 76

Columbella terpsichore Leathes (Pl. IV, fig. 12) . . 76
Columella=Sphyradium 60
Conchological chat from London 85
Conchological Museum 34
Cryptogirasia 70
Cryptosoma, Theob., 1857, not Bruelle = Megaustenia, n. n. 70
Cuba, A day's collecting at Antilla 2
Cuba and the Isle of Pines, A collector in western . 99, 111
Cuban land shells, New 42
Cuban, Pleistocene fossils from Eastern 69
Cuban Urocoptis, New 55, 109, 125
Cumberlandia, A new genus of Naiades 13
Cumberlandia monodonta Say 14
Cyclas albula Prime 139
Cyclostoma semicanum 111
Cypraea erosa carmen Smith, n. var. (Pl. IV, fig. 10) . 78
Cypraea erosa nebrites Melv. (Pl. IV, fig. 9) . . . 78
Cypraea erosa subalba Smith, n. var. (Pl. IV, fig. 11) . 78
Cypraea helvola Linné (Pl. IV, fig. 8) 78
Cypraea helvola Linné, var. mascarena Milv. (Pl. IV, fig. 6) 78
Cypraea hirasei Roberts, n. sp. (Pl. VI, figs. 1, 2, 3) . 97
Cypraea hungerfordi küensis Robt., n. subsp. (Pl. VI, figs. 5, 6, 7, 8) 98
Cypraea pantherina and vars. 78
Cypraea turdus Lam. 78
Epiphragmophora dupetithouarsi cuestana Edson, n. subsp. 37
Epiphragmophora dupetithouarsi 49
Epiphragmophora oregonensis Lea, Notes on the validity of 49
Eulima tenostoma 55
Floridian Viviparus, A new 48
Florida, Land shells of southern 16, 31
Fresh-water Mollusca living out of the water . . . 64
Galba montanensis Baker, n. sp. 115
Glandinella poeyanas Pfr. 112
Gripp, Charles W. (obituary) 132
Guatemala, Notes on shells from Quirigua 11
Harmandia somboriensis Rochebrune 141
Helix alonensis Fer. (Pl. IV, fig. 5) 26
Helix hortensis Müll. 74
Helix (Iberus) qualtierana Linn. (Pl. IV, figs. 1, 2, 3) . 26
Helix (Iberus) qualtierana, var. grandae Smith, n. var. (Pl. IV, fig. 4) 27
Hermann (Dr. Johann), Work in "Der Naturforscher," 1781 45
Holospira bilamellata Dall (Pl. V, figs. 8, 9) . . . 41
Holospira mesolia Pils., n. sp. 89
Hyria 141

Lamarck's Collection of Shells 119
Lampsilis ventricosa (Barnes) in the upper Potomac drainage 51
Lymnæa auricularia near Philadelphia 27
Lymnæa from Montana, A new 115
Lymnæa parva. 65
Lymnæa (Radix) auricularia at Cambridge, Mass. . . 116
Lymnæa (Radix) auricularia in Colorado 84
Macroceramus canimarensis rotundibasis Pils., n. subsp. . 130
Maine mollusca, Notes on 73
Marine gasteropoda from Aden with description of new forms 74
Marginella obscura Reeve, (Pl. IV, fig. 16) . . . 77
Megaustenia Ckll. n. n. for Cryptosoma preoccupied . . 70
Mexico, Shell collected at Manzanillo, west . . . 143
Microceramus perconicus trinidadensis Pils., n. var. (Pl. VIII, figs. 15, 17, 18) 129
Microceramus sanctispiritensis Pils., n. sp. (Pl. VIII, figs. 11, 12, 13) 127
Missouri, Shells of southeast 90
Mitra costellaris Lam. 77
Mitra deshayesii Reeve 77
Montana, A list of the Mollusca from the Mussel Shell valley 130
Murex despectus Linné 47
Murex fulvescens Sowerby 120
Musculium orbiculare Sterki, n. sp. 140
Nassa pulla var. minor Smith n. var. (Pl. IV, fig. 17) . 77
New England notes 54
Nerita cancellata Herm. 46
New Mexico, Mollusca from Northern 80
New Orleans mollusca 36
Nodularia jourdyi Morelet 141
Notes . . 12, 24, 35, 48, 59, 69, 84, 96, 108, 119, 132, 144
Nyctilochidæ 59
Nyctilochus Gistel = Septa Perry = Triton Auct. . . 59
Odostomia trifida, winkleyi, etc. 54, 74
Oliva andamanensis Bdg. (Pl. IV, fig. 7) 77
Opeas clavalinum kyotoense at Cambridge, Mass. . . 24
Oreohelix alta Pils., n. var. 144
Oreohelix carinifera Pils., n. sp. 89
Oreohelix colonies in Colorado 9
Oreohelix cooperi W. G. B. 9
Oreohelix cooperi minor Ckll. 10
Oreohelix haydeni betheli n. subsp. Pils. and Ckll. . . 144
Oreohelix haydeni Gabb. 11, 89
Oreohelix haydeni gabbiana Hemp. . . . 11, 30
Oreohelix hendersoni Pils., n. sp. 11, 29
Oreohelix strigosa depressa Ckll. (Pl. V, fig. 16) . 41, 81
Ovula (Neosimnia) hirasei Pils., n. sp. (Pl. VII, fig. 4) . 114

Ovula nipponensis Pils., n. sp. (Pl. VII, fig. 9) . . 114
Panama, Collecting in 121
Paludestrina salsa at Rowley, Mass. 54
Parreysia dalliana Frierson, n. sp. 142
Physa sayi 65
Physunio micropterus Morelet 141
Pisidium abditum Hald. 6
Pisidium abditum subrotundum Sterki . . . 7, 65
Pisidium alleni Sterki, n. sp. 9
Pisidium columbianum Sterki, n. sp. 117
Pisidium fraudulentum Sterki, n. sp. 95
Pisidium furcatum Sterki, n. sp. 118
Pisidium (furcatum var.?) rhombicum Sterki . . . 119
Pisidium glabellum Sterki, n. sp. 137
Pisidium lermondi Sterki, n. sp. 138
Pisidium minus Ad. 7
Pisidium (nevadense var. or sp.?) modicum Sterki, n. . 137
Pisidium overi Sterki, n. sp. 136
Pisidium peraltum Sterki. 95
Pisidium regulare Prime 7
Pisidium sphaericum Sterki, n. sp. 8
Planorbis bicarinatus Say, abnormal (Pl. V, fig. 4) . . 41
Pleistocene fossils from Eastern Cuba 69
Pleurodonte (Labyrinthus) goldmani=sipunculata . . 60
Pleurodonte marginella Gmel. 3
Polygyra albolabris Say (Pl. V, fig. 13). 40
Polygyra appressa Say. 1, 93
Polygyra cereolus (Pl. V, figs. 1, 2, 3). 39
Polygyra cereolus and var. carpenteriana . . 16, 31, 33
Polygyra elevata Say (Pl. V, fig. 15) . . . 40, 91
Polygyra exoleta Binn. (Pl. V, fig. 14) 40
Polygyra fallax Say. 80
Polygyra fallax goniosoma Pils., n. subsp. . . . 80
Polygyra fraterna Say (Pl. V, fig. 5). . . . 40, 91
Polygyra hirsuta Say 40, 91
Polygyra hopetonensis var. obsoleta. 79
Polygyra inflecta Say (Pl. V, fig. 6). . . . 39, 91
Polygyra mobiliana, banded 120
Polygyra palliata Say, in New England 59
Polygyra profunda Say (Pl. V, fig. 12) 40
Polygyra texasiana. 90
Polygyra tridentata Say (Pl. V, figs. 10, 11) . . . 39
Polygyra vannostrandi Bld. 79
Polymita muscarum 2
Pseudogalba new name for Simpsonia 120
Publications received 60, 71
Pyramidella bartschi=Odostomia bartschi. . . . 54
Pyramidella fusca and producta at Chatham, Mass. . . 55
Radix auricularia at Cambridge, Mass. 116
Rostellaria curvirostris Lam. (young) 79

Rumina decollata in Mobile and New Orleans . . . 4
Septa Perry, Note on the genus 58
Snail trap 60
Solenaia (Mycetopus) rugatus Sowerby 142
Sonorella argus Edson, n. sp. 37
Sonorella coloradoensis Stearn 39
Spain, Notes on the Molluscan fauna of the Sierra Elvira . 25
Sphaeriidae old and new 6, 95, 117
Sphaerium acuminatum Prime. 139
Sphaerium occidentale Prime 64
Sphaerium ohioense Sterki, n. sp. 139
Sphyradium, American species of 60
Spondylus americanus Hermann (1781)=echinatus Mart. (1784). 46
Strange names for old acquaintances. . . . 45
Strombus columba Lam. 79
Taylor, D.D., George W. (obituary) 82
Terebra clarkei Smith, n. sp. (Pl. IV, fig. 18). . . . 75
Thysanophora cockerellæ Pils., n. sp. (Pl. I, figs. 5, 6) . 12
Trapezoideus misellus Morelet 141
Trivia eos Roberts, n. sp. (Pl. VII, figs. 10, 11) . . 99
Trochetella stellata Pory 111
Turbo canaliculatus Herm.=T. spenglerianus Gmel. . 46
Unio (Lampsilis?) purpuriatus Say (Pl. III, figs. 4, 5) . 22
Unio (Nodularia) putzeysi Preston, n. n. for N. submigra Preston 70
Unio (Obovaria) jacksonianus Frier., n. sp. (Pl. III, figs. 1, 2, 3. 23
Unio silongweensis, n. n. for N. vicinus Preston not Lea . 35
Unionidae, A criticism of Dr. Haas' monograph . . 141
Urocoptis acicularis Torre, n. sp. (Pl. VI, fig. 6) . . 68
Urocoptis bacillaris Torre. n. sp. (Pl. VI, figs. 7, 8). . 57
Urocoptis bacillaris exilis Torre, n. var. (Pl. VI, fig. 10) . 58
Urocoptis bacillaris albicostulata Torre, n. subsp. (Pl. VI, fig. 9). 66
Urocoptis bacillaris sordidula Torre, n. subsp. (Pl. VI, fig. 11) 66
Urocoptis boqueronis Torre, n. sp. (Pl. VI, figs. 1, 2, 3) . 67
Urocoptis camagueyana Torre, n. sp. (Pl. VI, figs. 12, 13) 68
Urocoptis canteroiana Arango (Pl. VIII, figs. 4, 5) . . 126
Urocoptis cara Pils. & Hend., n. sp. (Pl. VI, figs. 27–30) 110, 128
Urocoptis cienfuegosensis Pils., n. sp. (Pl. VIII, figs. 6, 7, 8) 127
Urocoptis cinerea Pfr. (Pl. VI, figs. 14, 15) . . . 55
Urocoptis (Cochlodinella) mediana Pils., n. sp. (Pl. VIII, figs. 1, 2, 3). 126
Urocoptis fortiuscula Torre, n. sp. (Pl. VI, figs. 24, 25, 26) 57
Urocoptis hidalgoi cabrasensis Pils. & Hend., n. var. (Pl. VIII, fig. 16) 110

Urocoptis ischna Pils. 105
Urocoptis livida Torre, n. sp. (Pl. VI, figs. 21, 22) . . 56
Urocoptis livida occulta Torre, n. subsp. (Pl. VI, fig. 23). 56
Urocoptis longa Pils. & Hend., n. sp. (Pl. VIII, figs. 9, 10) 128
Urocoptis pallida Torre, n. sp. (Pl. VI, figs. 18, 19, 20) . 57
Urocoptis parallela Torre, n. sp. (Pl. VI, figs. 4, 5) . . 67
Urocoptis philippiana Pfr. 104
Urocoptis pruinosa Mor. 112
Urocoptis scaeva Gund. 104
Urocoptis stricta Torre, n. sp. (Pl. VI, figs. 16, 17) . . 66
Viviparus contectoides in Philadelphia 84
Viviparus walkeri Pils. and Johns, n. sp. (Pl. III, figs. 6, 7) 48
White, James Johnson (obituary) 35

INDEX TO AUTHORS.

Baker, H. B. 3
Baker, F. C. 115, 120
Barret, Wm. T. 61
Berry, S. S. 130
Clapp, Geo. H. 120
Clapp, W. F. 24, 116
Cockerell, T. D. A. 69, 70, 144
Dall, W. H. 58, 108, 132, 143
Daniels, L. E. 38
Edson (Harry), Henry M. 37, 49
Frierson, L. S. 22, 23, 141
Gaines, Marshall R. 34
Hanham, A. W. 83, 133
Hedley, C. 45, 85
Henderson, Jr., John B. 2, 42, 109
Henderson, Junius 9, 80, 84
Hinkley, A. A. 36, 47
Johnson, C. W. 48, 59, 120
Long, Bayard 27
Ortmann, A. E. 13, 51
Pilsbry, H. A., 1, 11, 29, 42, 45, 48, 60, 72, 79, 84, 88, 109, 114, 125, 144
Presbrey, Eugene W. 121
Preston, H. B. 35, 70
Roberts, S. Raymond 97
Sampson, F. A. 90
Smith, H. H. 4
Smith, Maxwell 25, 74
Sterki, V. 6, 64, 95, 117, 136
Torre, Carlos de la 55, 66
Vanatta, E. G. 12, 16, 31, 69, 84, 96
Wheeler, H. E. 14, 99, 111
Winkley, H. W. 54, 73, 144



1, 2. POLYGYRA CLARKII BRADLEYI VAN. (VOL. XXV, p. 120).
3, 4. ALDRICH & PILSBRY: A NEW PLIOCENE NERITINA.
5, 6. THYSANOPHORA COCKERELLÆ PILSBRY.
7, 8. HINKLEY: ON ANCULOSA.
9, 10. AMNICOLA WINKLEYI PILSBRY.
11, 12. ASHMUNELLA FERRISSI PILS. ARIZONA.

The Nautilus.

Vol. XXVI. MAY, 1912. No. 1

A NEW SPECIES OF AMNICOLA.

BY H. A. PILSBRY.

Amnicola winkleyi, n. sp, Plate I, figs. 9, 10.

The shell is umbilicate, ovate-conic, composed of 4½ to nearly 5 strongly convex whorls, separated by a very deep suture; whitish corneous when cleaned of the thin brown stain, glossy, faintly striate; apex rather acute, aperture ovate, nearly vertical; peristome continuous, a little flattened where it is very shortly in contact with the proceeding whorl.

Length 4.8, diam. 3.1 mm.; length of aperture 2.2 mm.

Saco, Maine.

By its dentition this species belongs to the group of *Amnicola lustrica* Pils., the shape of the central tooth and its denticulation being quite unlike *A. limosa*. It differs from *limosa* by its projecting first whorl, that of *limosa* being flat on top. *A lustrica* is smaller and more slender. *A schrökingeri* Ffld. is closely related but is a very much smaller species. Frauenfeld gives the length as 3.15 mm., and the largest in a series of over 100 before me is 3.2 mm. long, with 4½ whorls.

A. winkleyi has some resemblance to *A. (Cincinnatia) cincinnatiensis* Anth. by its acute apex and very convex whorls, but it is less solid, and differs very widely in dentition.

This species is one of the fruits of the careful examination of the New England fauna by the Rev. Henry W. Winkley, during the past decade or more. Doubtless the species exists in many collections under one name or another. My attention was called to it in the course of a study of the northern species for a monograph of the New York fauna.

A DAY'S COLLECTING AT ANTILLA, CUBA.

JOHN B. HENDERSON, JR.

The very new little town of Antilla, at the head of Nipe Bay, Cuba, is a very unlikely looking place for land-shells. Off in the distance are fine ranges of high mountains that beckon to the collector, but the few hills near Antilla are low and seem always to be surrounded by mangrove swamps and therefore quite inaccessible. In the hopes of finding something we turned to the only approachable hill, one lying just at the edge of the town. It is a wretched little apology for a hill, without rocks or any good feature for the collector other than a somewhat vigorous crown of vegetation. Arrived at the spot we found the place most uninviting. Goats, pigs, chickens and the whole assortment of Cuba's domesticated fauna ranged the woods, and a very dead horse sent clear to Heaven a protest against our invasion.

But no collector of land-shells happening along Antilla way should neglect devoting an hour or so to these goat-infested woods, for right there live in reasonable abundance the finest *Polymita muscarum* that ever came out of Cuba. Practically all the color varieties known of that really exquisite arboreal snail appear to live there. They range from black (with one orange band) through the yellows and reds (generally brilliantly painted in bands) to the pure china-ware glazed white. One would naturally suppose that so bright a color spot on a tree as this shell presents would be most easily detected. When once seen it certainly looks easy enough, yet I feel sure that we collected but a small proportion of the inhabitants. The reason is that the snail usually hides in the crotch of a limb, or fastens itself to the trunk of shrub or tree at just the spot where a convenient leaf covers him. The best manner of stalking them is to look along the ground for "bones" and when a "cemetery" is found then turn the eyes upward to the vegetation, for these creatures seem to have a gregarious habit and live in colonies. This may not be true in the rainy season when all tropical snails are more active.

From this little wooded hill we took besides the *muscarum* some very fine *Cepolis alauda* Fer. of the more typical form than is usually seen in collections. This is also a tree snail with much the same habit as the *muscarum* but they do not affect the colonial mode of life as definitely, and they are less shy in exposing themselves on

the tree to full view of the eager collector. The great range of variation in this species, not only in color but in size and form causes one to wonder if there are not possibly several distinct species included in the one comprehensive name of "alauda." There is also there, in these woods, but rather sparingly, *Pleurodonte marginella* Gmel. of a very dark color form. These also live on the trees, but only on the large trees and then invariably on the trunks. They look just like great big buttons and may be detected a hundred feet away. On the ground, quite common, were *Pleurodonte auricoma provisoria* Pfr. and a Helicina that we may call, for the present, *submarginata* Gray, also, *Oleacina subulata* Pfr. On the tree trunks, low down near the ground, very strictly of social habit, were *Macroceramus pictus* (Gundl.) Pfr. and, more commonly distributed, the *Chondropoma revinctum* Poey.

We answered the call of the more distant mountains later and I would like to tell of that trip when I shall have had time to identify the material collected. That is up in the "picta" country. I know of nothing more wonderful for jaded nerves and "that tired feeling" than to range the woods and banana fields for *Polymita picta.*

NOTES ON CIRCINARIA CONCAVA (SAY) FROM SHEBOYGAN COUNTY. MICHIGAN.

BY H. BURRINGTON BAKER.

Circinaria concava was considered to be a typical shell of the Upper Austral, and in the southern peninsula of Michigan was not known to extend much farther north than the old Saginaw-Grand River outlet. In consequence, it was a surprise to find it last summer near the north shore of Burt Lake, Cheboygan County, about one-hundred and fifty miles north of this line. Even more peculiar, however, was its habitat, as it was living in and along the edge of a large, boreal, cedar-spruce-fir (*Thuja-Picea-Abies*) bog, along with such a typically boreal species as *Pyramidula asteriscus* (Morse).

A quite thorough search of the hardwoods and other transition habitats of the region failed to reveal this shell, so it appears we have an *austral* shell living in a *boreal* bog in a *transition* region. One is tempted to conclude that this species, weakened by living too far north, is unable to cope with its competitors in the more favor-

able habitats, and is, for that reason, restricted to the unfavorable, sparsely populated, boreal bog, just as the boreal plants and animals are confined to these relict islands throughout the transition and the northern parts of the Upper Austral zones. These rivals would probably not be other shells in the case under consideration, as *Circinaria* is the only snail in the region that is mainly carnivorous in habits, but would probably be some of the carnivorous arthropods.

The shells themselves show the influence of poor conditions as they are, thin, fragile, and dwarfed. The largest specimen collected, which was a shell of four and one-half whorls measures: greater diameter 11.9 mm., lesser diameter 10.4 mm., altitude 5.3 mm., while a shell with five whorls from southern Michigan measures: 18.9 mm., 16 mm., 8.5 mm.; and a young shell of four and one-half whorls from the same region as the latter: 13.9 mm., 11.8 mm., 6.4 mm. This difference in size is even more marked in the embryonic first whorl, and the northern shells are more transparent and vitreous than the southern, have little of the latter's clouded, milky appearance, and truly resemble in texture an over-grown *Vitrea*. In addition, the revolving, microscopic wrinkles, which mark the typical shells over the entire upper surface but most strongly along the inner, sutural slope of each whorl, are more delicately and strongly marked in the northern form.

The question as to how this form got into this region is, at present, unanswerable, but the fact that the bog, in which it was found, lies between the Nipissing fossil beach and the present shore of Burt Lake is at least interesting in this connection.

RUMINA DECOLLATA IN MOBILE AND NEW ORLEANS.

BY. HERBERT H. SMITH.

This species is common in Southern Europe and has been carried by commerce to Bermuda, Havana and other places. As far as I know, the only recorded North American locality is Charleston, S. C., where it has been fully naturalized for many years. For its discovery in Mobile and New Orleans we are indebted to Mr. L. H. McNeill, of the former city. This gentleman sent to me a drawing of the shell, stating that he had first noticed it in New Orleans as

early as 1907,[1] and subsequently found it common in Mobile. Mr. McNeill is interested in shells and had collected them with the late Dr. Charles Mohr : he did not remember seeing this species in the Mohr cabinet or elsewhere, and asked me for information. The drawing, though he apologized for it, was so good as to leave little doubt in my mind ; moreover, Mr. McNeill stated that the shells were always decollate ; and, from their occurrence in the older or French parts of the two cities, he believed them to be introduced.

At my request he took the trouble to gather and forward more than 100 living specimens ; and a few days after, he sent some dead but good shells which he had brought from New Orleans. These are like Charleston or Bordeaux specimens. Mr. McNeill's notes are so good that I gladly transcribe them :

" The Mobile specimens were gathered at the S. E. corner of Monroe and Royal Sts. Another location is the S. E. corner of St. Emanuel and Canal Sts. Both are in the oldest parts of the town, and the first on the site of old Fort Charlotte (Fort Conde of the French ;) there were dozens of them crawling across the sidewalk, it being a damp and cloudy day, and many had been crushed by pedestrians. The shell is found over quite a large area of the city."

With regard to the New Orleans specimens he writes : " I noticed them first while putting up a building on the uptown side of Jackson Ave., two doors towards the river from Prytania St., and afterwards in other locations, including Girod and Claiborne Cemeteries. I also saw a gardener destroying several of them, remarking that they were a pest among the violets, which suggests that they are well known all over the city."

These exact locality notes are interesting. In Mobile, at least, the snail appears to be common only in the older or French streets, and probably that is why so large a species has been overlooked by Mohr, Showalter and other good collectors. During a recent brief visit, I collected in vacant lots where rubbish piles, stones, etc., gave plenty of shelter : snails were much in evidence, but no Rumina was seen. I was not looking for it, not knowing, at the time, that it lived in the city ; but I could hardly have missed so large a species had it been at all common. These lots were in newer parts of the

[1] After this was written and sent to press I learned that Mr. Ferriss collected *Rumina* at New Orleans, some years ago. As far as I know the fact was not published.

city, a mile or more from the French streets. Probably the species, though fully naturalized, has not extended its range so far.

All the indications are that Rumina was introduced in Mobile a century or more ago, during the French rule, and in all probability it was brought from some French port. The New Orleans race may be as old or older; both cities date from the 17th century. I shall not be at all surprised if Rumina turns up in Pensacola, another old colony; and it is more than probable that other European species may be found in these Southern cities.

SPHAERIIDAE, OLD AND NEW.

BY V. STERKI.

A revision, or monograph, of the North American *Sphæriidæ* has been demanded for years. But the way is not clear: very much is yet to be learned with respect to many species, their interrelations, the extent of their variation, and their distribution. A great deal of material for study from all over the continent is still desirable. A catalogue of the known species, with some notes will be published soon, and on the following pages will be found some descriptions and rectifications.—The numbers cited are those of the special collection of *Sphæriidæ*.

Piridium abditum Haldeman, Proc. Acad. Phila., I. p. 53, 1841; Prime, Mon. Corb. 1865, p. 68.—The *P. abditum* muddle has been the greatest stumbling block in the otherwise difficult study of *Pisidium*. Beginning with T. Prime, this "species" has been the general dumping ground into which almost anything was thrown that was not plainly "a black bass or a snapping turtle," or to remain serious, any *Pisidium*, even such as *noveboracense*, *variabile*, *æquilaterale*, *compressum*, etc.. and occasionally *Sph. occidentale*.

The authentic specimens, from Lancaster Co., Pa., in the T. Prime collection[1] are of a rather small form but serve to define the

[1] They had been there for about seventy years, and remained covered with a ferruginous crust so that their real shape, surface sculpture and color could not be seen. Now they are clean. Of the seven, five are immature, of a rather small form, as noted, 2.8–3.3 mm. long, the others, two pair of valves, are 3.8-4 mm. long. The description, in Mon. Corb., is quite inadequate, even with respect to the original form: "very inequilateral" [should be inequi-

species, which is really widely distributed, common and very variable, even if restricted after eliminating the three following species and some others which are clearly distinct.

The mussels of the typical and near-typical forms average larger and more inflated. It appears that there are several geographical subspecies; the following is one of them:

P. abditum subrotundum St. (*P. subrotundum*, The Nautilus XX, p. 19, 1906) The mussel is less inflated than in *abditum*, averaging smaller, the beaks are broader and less elevated; rather uniform. Appears to be a northern form; New England to Ontario, Michigan and northern Ohio.

P. minus (Ad.), *Cyclas minor* Ad., Proc. Boston Soc. N. H. I, p. 48, 1841; ranged under the syn. of *P. abditum* by Prime, Mon. Corb., p. 68. Is distinct so far as can be judged from the authentic specimens (3 valves, from Waybridge, Vt.) in the T. Prime collection, but appears to be rather rare. Specimens fairly well agreeing with them have been seen from: Duxbury, Mass. (W. F. Clapp, No. 6969); Denisville, Me. (Dr. Jas. Lewis' collection, now Mr. Bryant Walkers' No. 2582); Klinger's Lake, St. Joseph Co. Mich. (B. Walker, 6846); Ann Arbor, Mich. (H. E. Sargent, No. 3099, none full-grown); Cowley Farm, Nepean, Ont. (Justice F. R. Latchford, No. 6929), and apparently Greenport, L. I, N. Y. (J. B. Henderson from Sanderson Smith, No. 675).

P. regulare Pme., Boston Journ. VI, p. 363, pl. XII, figs. 11, 12, 1852. Also arranged under the synonymy of *P. abditum*, in Mon. Corb., l. c.; certainly not a form of *abditum*, to judge from the single specimen from "Ohio" in the Prime collection, ranging nearer *P. noveboracense* Pme., but appears to be distinct. Specimens identical were collected near Douglas Lake, Cheboygan County, Michigan, 1911, by Mr. H. B. Baker, Nos. 7084 and 7085, and probably the same near Geauga Lake, southeast of Cleveland, Ohio, by Mr. John A. Allen, 1910, No. 6542. Others were seen before but not recognized.

partite]: generally not so, and the figure shows rather the contrary and, by the way, has the beaks too high and otherwise faulty; "generally light straw color" is exceptional: generally light to dark corneous, often yellowish "cardinal teeth separate" is meaningless; "long. 0.15, lat. [= alt.] 0.14: diam. 0.09 inches" does not agree with "shell elongated," and with the figure "The surface is at times also quite rough and the striæ coarse"—conf. with; "the exterior surface is much smoother" (than in *P. noveboracense*).

Pisidium sphæricum, n. sp.—Mussel very inequipartite, strongly inflated, nearly globular; superior margin nearly straight, short, posterior subtruncate, the others rounded, supero-anterior slope barely marked; beaks near the posterior end, high, prominent over the hinge margin, rounded; scutum and scutellum distinctly marked; surface dull to slightly glossy, with fine, irregular striæ and a few lines of growth, color yellowish to grayish corneous, lighter along the margins, often with brown to blackish mottlings, whitish to straw in the young; shell rather strong, posterior muscle scars and pallial lines distinct; hinge stout, curved, cardinal teeth rather small, the right angular, with the posterior part thicker and grooved, the left anterior short, abrupt, angular, the posterior steeply oblique, little curved; ligament strong. Soft parts not examined.

Long. 5.6, alt. 5.3, diam. 4.6 mm. (100: 95: 82).

Long. 4.8, alt. 4. diam. 3.4 mm. (100: 84: 71).

Hab.; New England to Virginia, Ontario and Wisconsin (?): Westbrook, Me., collected by Prof. A. H. Norton; Saco, Me., Danvers and Lynnfield, Mass., by the Rev. H. W. Winkley; George R. North, Warren, Knox, Co., Me., by N. W. Fermond: Cambridge and Woburn, Mass., by Wm. F. Clapp.; Va. oppos. Washington, D. C., Sterki; Unalitoulin Id., Ont., by Justice F. R. Latchford; Fox River, Wis., by Geo. T. Marston, sent by E. W. Roper (if no label was misplaced; one of the two specimens in the collection is true, the other is a *P. abditum*).

Types: No. 797, Westbrook Me., for full-grown specimens, No. 727 Saco Me., and 6641, Lynnfield, Mass., for juv. and adolescent of a smaller form; yet anyone of the lots in the collection might serve as types.

This very distinct and characteristic *Pisidium* has been known for many years, but unfortunately much misunderstood and ill-treated. Generally it has been taken for a form of *abditum;* the late E. W. Roper named it *adamsi*, and under that name, specimens are in various collections; I have named it *P. minus*, having seen two specimens under that name, in a collection, but from that species it is also distinct, as evidenced by authentic specimens of *minus*.—From larger forms of *abditum* it is easily distinguished by the large and prominent beaks situated close to the posterior end of the mussel, and the strong hinge. In the young, the superior and supero-anterior margins form a peculiar somewhat angular curve, much as

in the young of *P. compressum* Pme.—It cannot be mistaken for any other species.

Pisidium alleni, n. sp.—Mussel rather large, well inflated, oval to ovoid in outline, without any projecting angles, beaks slightly behind the middle, large, prominent, rounded; scutum and scutellum slightly marked, small; surface shiny, with irregular striæ and several (4–6) strongly-marked lines of growth, color straw to light yellowish, plumbeous around the beaks in old specimens; shell and hinge medium strong, the latter rather long, cardinal teeth placed on the upper part of the hinge plates, the right angular with the posterior part thicker and grooved, left anterior short, pointed, the posterior oblique, rather long, nearly straight, ligament strong.

Long. 6.2, alt. 5, diam. 3.8 mm. (100 : 81 : 61). Half-grown specimens are rather different in shape, somewhat oblique, moderately inflated, posterior margin subtruncate, beaks markedly low and broad, apparently more posterior; young mussels are very flat.

Hab.—New England, New York; Hebron and Buckfield, Oxford county, Me., collected by Mr. John A. Allen; Franklin county, Me., by Mr. N. W. Lermond; "N. H." in the collection of Dr. James Lewis (B. Walker's), and that of the Cincinnati Soc. N. H.; Clifton, Staten Island, N. Y., Mr. J. B. Henderson, from Sanderson Smith. Types No. 5871a from Hebron, Me.

The only species to which this has resemblance is *roperi* St., which is much smaller, with shell and hinge slighter, and half-grown specimens not so oblique, and markedly constant over a wide territory. I take pleasure in naming the species in honor of Mr. J. A. Allen, who has done much good collecting in New England and Ohio.

(*To be continued.*)

OREOHELIX COLONIES IN COLORADO.

BY JUNIUS HENDERSON.

(*Concluded from Vol. XXV, p. 139*)

Oreohelix cooperi (W. G. B.)

Where the formations were badly cut up by igneous intrusions and displaced by folds and faults, in the neighborhood of Rabbit Ears, and elsewhere, attention was sometimes called to the existence of the calcareous formations in unexpected places by the presence of

numerous dead snail shells lying on the soil-covered surface. The species is very abundant about nine miles east of Meeker, on a Lower Mancos shale slope, with *gabbiana*. We have but one reversed specimen of the species.

These records show that *cooperi* is much more generally distributed than any other *Oreohelix* in Colorado, with its variety *minor* a close second, and *depressa* third.

Oreohelix cooperi minor Ckll.

I recently found dead shells very abundant, including one reversed specimen, on the exceedingly dry slopes of a gulch where the red Carboniferous sandstones were almost barren of soil and vegetation, except dwarf cedar and pinyon pine, altitude 7,200 feet. There are thin limestone bands in the formation. By 15 minutes' search 37 live ones were obtained, the majority being, to my surprise, under a small pinyon log and the cones which had lodged against it, the balance two or three at a time under small cedar logs. This form also occurs on a similar slope about two miles distant, and I found no other species of *Oreohelix* there. A few were found on a steep Niobrara slope southeast of Sulphur Springs. A very fine colony occurs on a steep, rather dry terrace south of White River, about two miles southeast of Meeker, under *Amelanchier*, etc., and another four miles southwest of Meeker. These were by no means as unfavorable localities as the McCoy and Sulphur Springs colonies of *minor* occupy, and seemed much better suited to snail life than the flourishing Muddy Creek colony of *cooperi*, and as well suited as the big *depressa-cooperi* colony at Steamboat Springs. My first impression was that perhaps *minor* was merely an ecological form of *cooperi*, dwarfed by unfavorable environment, but more careful consideration of the facts of distribution and an examination of our large series of both, seem to render that supposition untenable. Four dead specimens are from Placerville, San Miguel county, and twelve from Montrose, were collected by Mr. E. R. Warren, the well-known mammalogist. We have a lot from near Ohio City, Colo. (Frank Rohwer), and ten dead ones from Johnson county, Wyoming (N. DeW. Betts). This form has been likened to *O. s. concentrata* Dall (Nautilus, XVI, 106–107), but I have compared it with authentic *concentrata* received from Dr. Dall and find it quite distinct. I should say that *minor* is a dwarf variety of *cooperi* and *concentrata* perhaps a dwarf form of *depressa* type.

The Ohio City specimens were abundant at 11,000 feet, just above timber line, near a snow bank. That it was not a case of dwarfing of specimens of *cooperi* on account of lack of moisture or low temperature is indicated by the fact that *O. s. depressa* at that and higher altitudes seems at its best.

OREOHELIX HAYDENI (Gabb).

The only place in Colorado from which this species has been received is Glenwood Springs, where it was found alive in such abundance by Bethel and Dakan, and some fossil specimens collected near there. I see no reason for Binney's conclusion that it is only a variety of *strigosa*.

OREOHELIX HAYDENI GABBIANA (Hemphill).

In the Glenwood Springs colony hereinbefore described there is some tendency to intergrade with *haydeni*, and the sculpture of the embryonic whorls further indicates a close relationship. Dakan found another astonishing colony near Newcastle, from which he brought over 1,000 very fine adult specimens collected alive in a few moments. I found them more sparingly at another point near there, on Lower Mancos shales. Nine miles east of Meeker I found a colony on Lower Mancos shales, where they were mingled with *cooperi* in about equal proportions. It was raining, and the snails were crawling about under the bushes by the hundreds.

OREOHELIX HENDERSONI Pilsbry.

Known only from the type locality, where dead shells were collected three years ago by Mr. A. Dakan, on Little Thompson Creek, ten miles northwest of Longmont, on a talus slope of Niobrara limestone. Dr. Pilsbry considered it new, so I asked Mr. Dakan to search for live ones. Last July, during better snail weather, he returned to the place and had no difficulty in getting 160 live ones and many more dead ones. He says "it appears to be a small, healthy colony."

NOTES ON SHELLS FROM QUIRIGUA, GUATEMALA.

BY HENRY A. PILSBRY.

In the course of a recent visit to Guatemala, chiefly for studying insects, Mrs. T. D. A. Cockerell collected the following snails, all from Quirigua.

Aperostoma dysoni (Pfr.)
Ampullaria sp. (young).
Helicina amœna Pfr.
Thysanophora plagioptycha (Shuttl.)
Thysanophora cockerellæ n. sp.
Opeas micra (Orb.)
Euglandina cumingi (Beck.)
Cæcilioides consobrina veracruzensis (C. and F.)
Succinea sp. (immature).
Aplexa princeps Phillips.

THYSANOPHORA COCKERELLÆ n. sp. Pl. I, figs. 5, 6.

Related to *T. conspurcatella*, *hornii* etc. The spire is conic, composed of nearly $4\frac{1}{2}$ very strongly convex whorls united by a very deep suture, the last whorl regularly rounded, rather coarsely striate. Umbilicus contained $3\frac{1}{4}$ times in the diameter. Aperture subcircular, about one-fifth of the circumference occupied by the preceding whorl.

Alt. 2.4, diam. 3.4 mm.

Quirigua, Guatemala, W. P. Cockerell, 1912.

The spire is decidedly higher than in *T. conspurcatella* Morel. It is nearer *T. proxima* Pils., from the state of Michoacan which agrees in size of umbilicus, but the spire is higher in *T. cockerellæ*. *T. fischeri* Pils. is a more delicate shell with narrow umbilicus. Seven specimens were collected, the largest being measured above and figured. Most of the lot are decidely smaller with 4 whorls. The surface is not in good condition, but will probably prove to have cuticular processes like those of the related species mentioned.

It is named for Mrs. Cockerell.

NOTES.

BERMUDA SHELLS.—Mr. Arthur Haycock presented to the Academy of Natural Sciences of Philadelphia a set of shells from Whitby Cave, Bermuda, including *Strobilops hubbardi* Br., *Zonitoides bristoli* Gul., *Thysanophora hypolepta* Shutt., *Bifidaria rupicola* Say, and *Striatura milium meridionalis* Pils. The *Striatura* is new to the Bermuda fauna.—E. G. VANATTA.

ERRATA.—February number, page 110, end of sixteenth line from bottom, for "the rows" read *two rows*.

THE NAUTILUS.

VOL. XXVI. JUNE, 1912. NO. 2

CUMBERLANDIA, A NEW GENUS OF NAIADES.

BY A. E. ORTMANN.

The type-species of this new genus is *Unio monodonta* of Say, which stands, in Simpson's system (Pr. U. S. Mus. 22, 1900 p 679) as *Margaritana monodonta* (Say).

As Simpson has correctly recognized, this species has affinity with the genus *Margaritana* (Type: *margaritifera* Linnaeus), and, consequently belongs to my family *Margaritanidæ* (NAUTILUS, 25, 1911 p. 129). Through the courtesy of Mr. Bryant Walker I had the opportunity to examine the soft parts of a number of individuals from the Cumberland River, and was able to ascertain that all essential characters given by me for this family are present, chiefly so the conformation of the diaphragm, of the posterior end of the gills, and of the anal and branchial openings. But there is one interesting difference. While in other species of the genus *Margaritana* the interlaminar connections of the gills are scattered, and irregularly distributed, or fall into more or less distinct oblique rows, in the present species these irregular, interrupted rows are *developed as continuous septa, which run obliquely forwards.* In the middle of the gills, these septa are well developed and quite continuous, but rather distant from each other. Toward the ends of the gills they are more irregular, shorter, and chiefly at the posterior end they are somewhat interrupted, and thus it is clearly seen that they represent a *further development* of the oblique, interrupted rows present in other species of *Margaritana*.

The septa of this species (*monodonta*) have nothing to do with the septa of the family *Unionidæ*, for in the latter they run parallel to

the gill-filaments while here they cross them obliquely. Moreover, the septa of the *Unionidæ* are much more closely set (chiefly in the females), and much more regular.

Margaritana monodonta differs by this structure markedly from all species of *Margaritana* I know. I have seen the typical *Margaritana*-structure not only in *Margaritana margaritifera* (L.) (both from Europe and northeastern America), but also in two specimens of *Margaritana sinuata* (Lam.) (= *crassa* Simpson, l. c. p. 678) from near Perpignan, France (sent to me by Mr. W. Israël), and in a number of specimens of *Margaritana hembeli* (Conr.) from southern Alabama (I am obliged to Mr. H. H. Smith for these). While I thus consider these three species as congeneric (the creation of *Pseudunio* for *sinuata* by Haas notwithstanding), *M. monodonta* differs from them in an important structure, which undoubtedly represents a higher specialization, a forward step in evolution, and it is, as I believe, entitled on this account to the rank of a separate genus, which I propose to call Cumberlandia (Type: *monodonta* Say).

As to the distribution of *Cumberlandia monodonta*, Walker's papers should be consulted (Pr. Mal. Soc. 9. 1910 pp. 137–139, and NAUTILUS, 25. 1911 p. 57–58). According to this, the metropolis of it seems to be in the Cumberland and Tennessee Rivers, but it has spread into the lower Ohio drainage, and into Indiana and Illinois. In the Ohio, it was not known, up to the present, above Cincinnati: but I may mention here, that I found last summer (July 13, 1911) a dead specimen of *Cumberlandia monodonta* in the Ohio River at Buffington Island, near Portland, Meigs Co., Ohio. I also may add, that according to the experience gained by my last year's collecting, it is entirely excluded that this species occurs anywhere in the Big Sandy or Great Kanawha Rivers, and that I do not think that it reached its present range by coming "down" the Ohio. But this will be discussed elsewhere.

ADDITION TO THE LIST OF MOLLUSCA FROM MONTE SANO, ALABAMA.

BY. H. E. WHEELER.

Through the kindness of Prof. Herbert H. Smith, conchologically the "discoverer" of Monte Sano, and Mr. George H. Clapp, who

determined the collections from this region, I am able to make the following additions to the list published in the March number of the NAUTILUS for the current year. It should be noted that some of the species listed then and now were not found on the mountain proper, but, living on the foothills or in the adjacent valleys, are entitled to enumeration in the fauna of the region.

Polygyra auriformis, Bld. This and the species *clausa* are low land shells, collected near the base of the mountain.

Polygyra pustuloides, Bld. Near Huntsville, and also on the mountain.

Polygyra tridentata Say. "Common in the whole region, and probably on the table land of the Mountain." Smith.

Polygyra clausa Say.

Polygyra hirsuta Say. Typical *hirsuta* is found on Monte Sano according to Clapp.

Strobilops virgo Pils.

Bifidaria contracta Say.

Bifidaria corticaria Say.

Bifidaria pentodon Say.

Bifidaria pentodon gracilis Sterki.

Bifidaria clappi Sterki. See NAUTILUS, Vol, XXII, pp. 108, 109. Sterki gives Huntsville, Alabama as one of the type localities. Clapp has it also from Monte Sano.

Vertigo milium Gld.

Vertigo rugosula ovalis Sterki. "In the Huntsville region, and almost surely from the mountain." (Smith).

Vertigo gouldii Binn.

Cochlicopa lubrica morseana P. From the western part of Vincent Mountain. This is one of the numerous connected spurs of the same mountainous chain, and the species is likely to be found on Monte Sano in suitable situations.

Omphalina pilsbryi Clapp. See NAUTILUS, Vol. XVIII, p. 30. Collected by the writer at Normal, several miles north of Huntsville, Alabama, and by Smith on Monte Sano.

Euconulus chersinus dentatus, Sterki. "More correctly a 'form'; certainly not a variety." (Smith).

Gastrodonta intertexta Binn. Collected by the writer at Huntsville, and by Prof. Smith on the mountain.

Gastrodonta acerra Binn. Huntsville region. Should be found on the mountain.

Punctum blandianum Pils.

Pyramidula cumberlandiana, Lea. Rarely at the southern end of Monte Sano. This record, furnished by Smith, cancels the negative note in previous article.

There is a variety of *Pyramidula alternata*, Say, which is quite constant, distinguished principally by its sub-carinate contour, which was collected on an isolated peak (Ward's Mountain?) three miles south west of Huntsville.

Pyramidula sub-sp. nov. (Pilsbry). A splendid sub-species of *bryanti* which has been described by Dr. Pilsbry, but not yet published. Locality a dry ridge about 6 miles north of Monte Sano.

Succinea avara Say. Reported by Clapp.

Carychium exiguum Say. "Associated with *exile*, H. C. Lea, but rare and local." (Smith).

LAND SHELLS OF SOUTHERN FLORIDA.

BY E. G. VANATTA.

This list is based upon shells picked from leaf-mould collected in Florida by Messrs. Clarence B. Moore, H. A. Pilsbry, C. T. Simpson, Stewardson Brown, H. W. Fowler, and J. S. Raybon.

All are in the collection of the Academy of Natural Sciences.

Many of the specimens of *Truncatella bilabiata* are a rather smooth form parallel to *T. c. succinea* Ad.

The typical *Polygyra cereolus* Mühl. were at first divided by size from the var. *carpenteriana* Bld., but these two forms intergrade as shown by the following count of specimens in a large series from two localities. In a lot from Little Pine Key there are 46 *Polygyra cereolus* measuring from 14.5 to 10.5 mm.; 700 *Polygyra c. carpenteriana* measuring 10.3 to 7.5 mm., and 40 measuring 8.3 to 7 mm. in diameter.

From Big Pine Key at Robert Watson's place there are 12 *P. cereolus* measuring 14.5 to 11 mm. in diameter; 1009 var. *carpenteriana* measuring 11.3 to 9 mm., and 16 from 9 to 7 mm. It will be seen that no definite division can be made. Each of these lots represents a single colony, the snails being scraped up with the dirt, without selection, from under stones in an area of a square rod or two.

The *Bifidaria contracta* from Florida form a race in which the inner portion of the parietal lamella is divided from the forward part by a depression, as if it were another lamina joined to the main one.

In picking them over, one has a tendency to divide each lot of *Bifidaria pellucida hordeacella* from Florida into four forms, figured on plate II, as follows:—form A, *long*, figs. 1. 3, 4, 5, 9, 10, 15; form B, *medium-sized*, figs. 6, 7, 8, 13, 16, 17, 19; form C, *small*, figs. 14, 20, 21, 24, 25, 26, 27, and form D, *obese*, figs. 12, 18, 22. All these forms seem to intergrade in size through a small number of examples. The variation observed is largely due to the formation of a lip when the shells have made a different number of volutions, as the long forms have about $5\frac{1}{2}$ whorls and the short forms only about 4 whorls. It is peculiar that the long shells from Florida have weak teeth and the short specimens have strong teeth; while in Texas the converse is often the case.

To give an idea of the relative abundance of the forms, in a lot of *Bifidaria p. hordeacella* from Bahia Honda Key I counted 25 long, 31 medium, 15 small, and 7 obese shells. In another series from Key West there are 111 long, 358 medium, 22 small, and 140 obese specimens. Each of these lots was from dirt gathered in the area of a square yard or less. The size in mm. of the several forms may be measured by the scale supplied on the plate.

Vertigo variolosa is very scarce, as only four perfect specimens were found. *Helicina tantilla* also continues to be excessively rare. Many other equally small species were obtained by thousands.

For lists of species from other keys visited, see the THE NAUTILUS, xix, 1905, p. 38, and xxi, 1908, p. 99. Altogether over 50,000 specimens have been picked from dirt and débris obtained on the Florida Keys.

BAHIA HONDA KEY, Lee Co., collected by Dr. H. A. Pilsbry.

Chondropoma dentatum Say.
Truncatella bilabiata Pfr.
Truncatella caribæensis succinea Ad.
Thysanophora selenina Gld.
Thysanophora plagioptycha Shutt.
Polygyra cereolus Mühl.
Polygyra cereolus carpenteriana Bld.
Pupoides modicus Gld.
Bifidaria p. hordeacella Pils.

Bifidaria rupicola Say.
Varicella gracillima floridana Pils.

Blue Hill, Horr's Island, Lee Co. (near Key Marco), collected by Clarence B. Moore.
Thysanophora plagioptycha Shutt.
Pupoides modicus Gld.
Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.
Vertigo variolosa Gld.
Guppya gundlachi Pfr.
Vitrea dalliana ' Simps.' Pils.

Boca Grande Key, Monroe Co., collected by Stewardson Brown and H. W. Fowler.
Drymæus multilineatus Say.
Cerion incanum Binn.
Succinea floridana Pils.

Boca Chica Key, Monroe Co. Collected by Dr. H. A. Pilsbry, C. T. Simpson, S. Brown and H. W. Fowler.
Chondropoma dentatum Say.
Truncatella caribæensis ' Sby ' Rve.
Truncatella bilabiata Pfr.
Helicina orbiculata Say.
Helicina orbiculata clappi Pils.
Thysanophora selenina Gld.
Thysanophora incrustata Poey.
Thysanophora plagioptycha Shutt.
Praticolella jejuna Say.
Cerion incanum Binn.
Pupoides modicus Gld.
Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.
Varicella g. floridana Pils.
Guppya gundlachi Pfr.
Vitrea dalliana ' Simps.' Pils.

Fikahatchee Key, Lee Co. Collected by Clarence B. Moore.
Helicina orbiculata Say.
Helicina o. clappi Pils.
Polygyra c. carpenteriana Bld.



VANATTA: VARIATION OF BIFIDARIA PELLUCIDA HORDEACELLA.



1, 2, 3. UNIO JACKSONIANUS FRIERSON.
4, 5. UNIO PURPURIATUS SAY.
6, 7. VIVIPARUS WALKERI PILS. & JOHNS.

Thysanophora selenina Gld.
Thysanophora plagioptycha Shutt.
Pupoides modicus Gld.
Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.
Bifidaria contracta Say.
Strobilops hubbardi Br.
Euglandina rosea Fer.
Vitrea dalliana 'Simps.' Pils.
Guppya gundlachi Pfr.
Zonitoides minuscula Binn.
Zonitoides singleyana Pils.

A SMALL KEY OPPOSITE FLAMINGO, Monroe Co. Collected by Clarence B. Moore.

Polygyra c. carpenteriana Bld.

GILBERT'S, Lee Co. (near Matanzas Pass). Collected by Clarence B. Moore.

Bifidaria rupicola Say.

GRASS KEY, Monroe Co. Collected by J. S. Raybon.

Truncatella caribæensis 'Sby' Rve.

KEY WEST, Monroe Co. Collected by Dr. H. A. Pilsbry and C. T. Simpson.

Truncatella caribæensis 'Sby.' Rve.
Helicina o. clappi Pils.
Thysanophora incrustata Poey.
Thysanophora plagioptycha Shutt.
Polygyra cereolus Mühlf.
Urocoptis poeyana variegata Pfr.
Microceramus pontificus Gld.
Pupoides modicus Gld.
Bifidaria p. hordeacella Pils.
Opeas gracile Hutt.
Opeas micra Orb.
Euglandina rosea minor Binn.
Varicella g. floridana Pils.
Succinea campestris Say.

LIGNUM VITÆ KEY, Monroe Co. Collected by Clarence B. Moore.

Helicina o. clappi Pils.

Lossman's Key, Monroe County. Collected by Clarence B. Moore.

Helicina orbiculata Say.
Helicina o. clappi Pils.
Thysanophora selenina Gld.
Thysanophora plagioptycha Shutt.
Thysanophora caeca Guppy.
Polygyra cereolus Mühl.
Polygyra c. carpenteriana Bld.
Polygyra septemvolva Say.
Microceramus pontificus Gld.
Strobilops hubbardi Br.
Bifidaria rupicola Say.
Bifidaria contracta Say.
Vitrea indentata Say.
Vitrea dalliana 'Simps.' Pils.
Euconulus chersinus Say.
Guppya gundlachi Pfr.
Zonitoides arborea Say.
Zonitoides minuscula Binn.
Zonitoides minuscula alachuana Dall.
Zonitoides singleyana Pils.

"McIlvaine's Place," on a small key two miles northeast of Marco, Lee Co. Collected by Clarence B. Moore.

Polygyra c. carpenteriana Bld.
Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.
Bifidaria contracta Say.
Vertigo rugosula oralis St.
Guppya gundlachi Pfr.
Zonitoides minuscula Binn.
Zonitoides singleyana Pils.

Little Marco Key, Lee Co. Collected by Clarence B. Moore.

Microceramus floridanus Pils.

Mormon Key, Monroe Co. Collected by Clarence B. Moore.

Truncatella caribæensis 'Sby.' Rve.
Truncatella bilabiata Pfr.
Polygyra c. carpenteriana Bld.

Nameless key (Mr. Addison's), two miles east of Marco, Lee Co. Collected by Mr. Moore.

Pupoides modicus Gld.

No Name Key, Monroe Co. Collected by Dr. Pilsbry.

Chondropoma dentatum Say.
Helicina o. clappi Pils.
Thysanophora selenina Gld.
Thysanophora incrustata Poey.
Thysanophora plagioptycha Shutt.
Praticolella jejuna Say.
Drymæus multilineatus Say.
Microceramus pontificus Bld.
Cerion incanum Binn.
Strobilops floridanus Pils.
Varicella g. floridana Pils.
Vitrea dalliana 'Simps.' Pils.
Guppya gundlachi Pfr.

Big Palo Alto Key, Dade Co., collected by Clarence B. Moore.

Helicina o. clappi Pils.

Panther Key, Lee Co., collected by Clarence B. Moore.

Polygyra c. carpenteriana Bld.
Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.
Guppya gundlachi Pfr.
Zonitoides singleyana Pils.

Explanation of Plate II.

Bifidaria pellucida hordeacella (Pils).

Fig. 1, 3, 5. Form *a*. Key West Fla. A. N. S. 104029.
Fig. 2. Form *a*. New Braunfels, Texas. A. S. N. 68888.
Fig. 4. Form *a*. Summerland Key, Fla. A. N. S. 104021.
Fig. 6, 7, 8. Form *b*. Key West. A. N. S. 104029.
Fig. 9, 10. Form *a*. Bahia Honda Key, Fla. A. N. S. 105521.
Fig. 11. Form *d*. New Braunfels, Texas. A. N. S. 68888.
Fig. 12. Form *d*. Key West. A. N. S. 104029.
Fig. 13. Form *b*. Key West. A. N. S. 104029.
Fig. 14. Form *c*. Bahia Honda Key. A. N. S. 105521.
Fig. 15. Form *a*. Key West. A. N. S. 104029.

Fig. 16. Form *b*. Key West. A. N. S. 104029.
Fig. 17. Form *b*. Bahia Honda Key. A. N. S. 105521.
Fig. 18. Form *d*. Bahia Honda Key. A. N. S. 105521.
Fig. 19. Form *b*. Bahia Honda Key. A. N. S. 105521.
Fig. 20, 21. Form *c*. Key West. A. N. S. 104029.
Fig. 22. Form *d*. Key West. A. N. S. 104029.
Fig. 23. Form *c*. New Braunfels Texas. A. N. S. 68888.
Fig. 24, 26, 27. Form *c*. Summerland Key. A. N. S. 102742.
Fig. 25. Form *c*. Bahia Honda Key. A. N. S. 105521.

(Key West, figs. 1, 3, 5, 6, 7, 8, 12, 13, 15, 16, 20, 21, 22. Bahia Honda Key, figs. 9, 10, 14, 17, 18, 19, 25. Summerland Key, figs. 4, 24, 26, 27. New Braunfels, Texas, figs. 2, 11, 23).

(*To be concluded*).

UNIO (LAMPSILIS?) PURPURIATUS SAY.

BY L. S. FRIERSON.

Though Mr. Say describes very well his *Unio purpuriatus*, its identity seems to have been nearly lost. That it may again be known to students, it is now figured, plate III, figs. 4, 5, and his description repeated. "Transversely oval, slightly oblong, in some specimens with a little tendency towards ovate; dirty yellowish or fuscous, obscurely radiate with blackish green: beaks behind [*before*] the middle, hardly raised; anterior [*posterior*] edge rounded, or rather composed of two nearly rectilinear lines; posterior [*anterior*] margin rounded; hinge margin regularly rounded; cavity of the hinge membranes (behind [*before*] the beaks) narrow but very obvious; basal margin arquated, a little compressed in the middle and sometimes almost contracted in that part; within purple, margin livid, anterior [*posterior*] submargin iridescent; primary teeth nearly direct, rather thick, striated; lateral teeth hardly extending beyond the sinus of the hinge margin. Length [*height*] one inch and three-fifths; breadth [*length*] three inches and one-tenth. Convexity over nine-tenths of an inch.

Remarks.—Conrad's identification of this species with *U. medellinus* Lea no doubt has confused the situation. Crosse and Fischer doubted this identification of Conrad's, but they erred in thinking that the name was badly spelled and *meant* to be *purpuratus*, preoc-

cupied by Lamark. Mr. Say intended this name to show that his shell was the Mexican analogue of *purpureus*, and hence his name *purpuriatus* is a valid name. Mr. C. T. Simpson's remark in his *Synopsis* that *U. strebeli* Lea was a *young medellinus* was an error, and still further confused the situation.[1] Examples of *U. purpuriatus* were collected by Mr. A. A. Hinkley and distributed as *U. strebeli*.

The species occurs in both white and purple nacre, and many are *entirely* rayless. The example figured is smaller than Say's, but some of Mr. Hinkley's specimens attained the dimensions given by Say.

UNIO (OBOVARIA) JACKSONIANUS, NEW SPECIES.

BY L. S. FRIERSON.

Shell ovate, smooth, rounded before and below, nearly straight from beak to post-point, which is about half way the height of the shell; umbonal ridge low, and the posterior area very narrow; beaks not high, sculpture not seen; nacre bluish white; iridescent behind; teeth double in left, single in right valve; cardinals stout, erect, laterals not very large; muscle scars confluent behind, separate before; pallial line obsolete behind. Length, 45 mm.; height, 33 mm.; thickness, 20 mm.

Habitat: Pearl River, Miss. (and also in the Yalabusha River, Miss.).

Numbers of this species have been in hand for several years and have been distributed largely under the name of "*U. castaneus* Lea, Variety." Collectors will rectify their names accordingly. The novelty of the present species was determined by gravid specimens collected by Mr. A. A. Hinkley, and by a comparison with *castaneus* of the Lea collection by Mr. Bryant Walker. This species has not the *female* form of *castaneus* and the dorsal scars are in the bottom of the beaks in place of being on the teeth.

In outline it is nearly exactly the same as *U. strodeanus* B. H. Wright (Proc. Acad. Nat. Sci. Phil., 1900, Plate 1, fig. 3), but dif-

[1] This error was caused by someone having marked a young *U. medellinus* "type" and misplacing it; and it finally got into the tray of *strebeli*. This explanation was given by Dr. Wm. H. Dall.

fers in being greenish, while *strodeanus* is dead black; its beak cavties are deeper, and its teeth are much longer, stouter and more erect.

From *U. curtus*, to which it is allied, it differs, lacking the pronounced swelling before.

Many of these shells show a constriction like the *U. constrictus* of Conrad, and it may be that this is a sexual mark.

Figs. 1 and 2 represent the normal shape, Fig. 3 the constricted form.

Types in the Academy of Natural Sciences, cotypes in my own collection, in the National Museum, and specimens of the same lot are in many other cabinets as "*castaneus*."

NOTES.

CARYCHIUM MINIMUM Mull.: It may interest students of geographical distribution, to learn that a thriving colony of *Carychium minimum* Mull., has been found in a greenhouse at Quincy, Mass. I have observed this species at intervals for nearly a year, and as it is becoming more plentiful, there seems to be a reasonable chance that it may escape to the surrounding country. Considering its natural habitat, there is a possibility of its becoming accustomed to our climate. W. F. CLAPP.

OPEAS CLAVULINUM KYOTOENSE Pils.: I have found a few specimens of *Opeas clavulinum kyotoense Pils.*, in a greenhouse at Cambridge, Mass., but only in the hottest house, although they are very hardy if kept in a warm vivarium. They will not survive a slight drop in temperature. Therefore there seems little likelihood of this species becoming a permanent addition to our northern fauna. Unlike the majority of our North American pulmonates this species does not appear to mind direct sunlight, but will leave the shade of a protecting leaf, to spend an hour or two in crawling on the glass, exposed to the rays of a bright midday sun.

I am indebted to Mr. George H. Clapp for the identification of this species. He states that specimens have also been found by him in Phipp's Conservatory, Pittsburgh, Pa., in 1901, and by Mr. Bryant Walker in a "Bigonia House" at Buffalo, N. Y.

W. F. CLAPP.



SMITH: MOLLUSCA OF THE SIERRA ELVIRA.
MOLLUSCA OF ADEN.

THE NAUTILUS.

VOL. XXVI. JULY, 1912. No. 3

NOTES ON THE MOLLUSCAN FAUNA OF THE SIERRA ELVIRA, SPAIN.

BY MAXWELL SMITH.[1]

While staying in Granada, Spain, during April 1910, the writer made several excursions for the day to the Sierra Elvira mountains. This striking range rises from the plain before Granada and may be distinctly seen from the battlements of the Alhambra.

The Sierra Elvira is a treeless Jurassic range named after the ancient town of Iliberris. It is covered with rocks so sharp in places that shoe leather offers little protection to the feet and climbing is necessarily a slow operation. On a clear day the sun reflected on the rocks makes the place not unlike a furnace. When the day is a cloudy one the biting cold wind from the snow-covered Sierra Nevada constantly keeps up. Nevertheless the splendid view amply repays the climber even if he is not in quest of snails. There are several small caves in a lofty valley several miles from the railway which skirts the base of the Sierra far below. Several species were found near the mouth of one of these caves. The more distant portions of the range no doubt contain other species not taken by the writer.

One day the ascent of the Sierra Elvira was made from Pinas Puente a quaint old town in an attractive situation on the Cubillas. In the brooks flowing into this stream a species of Melanopsis was noticed in abundance. It was at Pinas-Puente that Columbus was overtaken by the messenger of Queen Isabella in 1492, when in

[1] Read before the Brooklyn Conchological Club, January 29, 1912.

despair he had given up negotiations with the Spanish monarchs and was actually on his way to France.

I was accompanied on these trips by William M. Bunker, Honorary President of the San Francisco Chamber of Commerce. Mr. Bunker assisted in turning over many large stones, adhering to the under surface of which the majority of the specimens were found.

Leucochroa candidissima Drap. This species lives on the southern sides of the Sierra.

Helix sp. A small form of the *caperata* group.

Helix alonensis Fér. (pl. IV, fig. 5). One example obtained.

Helix (*Iberus*) *gualtierana* Linn. Plate IV, fig. 1 (typical), 2, 3, (convex form).

Eighteen specimens are before me: a selection made with a view to proportion, arranging these in lots according to the height of the spire, there are :

10 shells with a flattened spire (a few are sunken above) (fig. 1).

4 shells in which the spire is more produced (fig. 2).

4 shells with the highest spire (fig. 3).

In the first lot the examples appear to approach closely the typical form of *gualtierana*, which was described by Linn. The typical form may easily be distinguished by the flattened or sunken spire, the half or nearly covered umbilicus, the strongly carinated periphery and the aperture which is decidedly broader than high. Pilsbry's figure (Manual, Vol. IV, plate 53 fig. 6) agrees perfectly.

A variety (plate 4 fig. 2) has the aperture of the type but with a more developed spire. In the third lot (fig. 3) the shells run about the same size as the type. The peripheral keel is similar. On the other hand the aperture is of a distinctly different shape. In the majority of the examples examined *it is nearly as high as wide*, less extended laterally than in the typical form.

Helix gualtierana var. *laurenti* Bourguignat (Revue et mag. de zoologie, 1870, p. 95, pl. 14, figs. 5–7) was based on a shell from Mt. Gador, near Almeria. It is decidedly *conic* above, the apical portion is not depressed as in convex forms of *gualtierana* though a little flattened at the tip ; the base is figured as rather strongly convex. Figs. 11–13 of pl. 53, Man. Conch. IV, are similar though larger, but figs. 9, 10, copied from Kobelt, cannot be this race.

I took a solitary example, dead and bleached, which agrees with the original account of *laurenti* in the conic top. It is solid, much

more finely sculptured than *gualtierana*, umbilicus wholly closed. It differs from *laurenti* by the more deeply descending last whorl and the much less convex base. Alt. 20, diam. 31 mm. If found to be a different race, it may be called ver. *granadæ* (pl. IV, fig. 4).

Helicodonta (*Caracolina*) *lenticula* Fér. The examples before me are smaller than the Sicilian specimens.

Buliminus quadridens Müll. Several dozen specimens were taken. Dr. Pilsbry, to whom I submitted the form, writes, "The *Buliminus* is *B. quadridens* Müll., rather a long form of the species, but certainly that."

LYMNAEA AURICULARIA NEAR PHILADELPHIA.

BY BAYARD LONG.

The writer's proper field of activity belongs in the realm of botany; but like many another, he sometimes wanders from the beaten path of his own special science. A note, however, upon the occurrence of a foreign species markedly thriving in our own natural habitats, and there associating with indigenous forms in so perfectly a naturalized state that only a knowledge of the normal distribution of the species makes it possible to detect the status of its occurrence: such a note seems rather familiar ground, since there is such a host of introduced plants whose migrations must be kept under observation. So the writer may, in a measure, be justified in breaking into Conchology. Botanists make rather a point of putting upon record the early occurrences of foreign elements entering into our flora, because, besides being of an historical interest, a lack of such data, among plants at least, has often led to considerable confusion and error as to what are actually indigenous plants of any given area. Such a condition is probably not so likely to occur among shells, but still it may be of no less importance to have definite record of observations of species becoming naturalized, in case the forms should possibly become widespread and a knowledge of their original centres of dissemination lost.

In 1910 there was brought to the Philadelphia Academy by Samuel R. Jacob some shells which he had casually collected in Europe, and also some specimens of *Lymnæa auricularia* (Linn.) which, however, were said to have been collected from near

Philadelphia in a pond about one mile south of Ogontz, Montgomery County, Penna., along Old York Road. A possibility of accidental mixture of specimens was thought to exist, and no published record made of it.

Upon Mr. E. G. Vanatta's learning that I lived near the *auricularia* locality, he mentioned the case to me and I felt confident that the station referred to must be that known as Dobbin's pond at Old York Road and Ashbourne Road. He asked me if I would not sometime examine the pond with the hope of verifying the Jacob record. I accordingly did so on May 22nd of this year and found, associated with *Planorbis antrosus* Conrad, an abundance of a large *Lymnæa* which proved to be the desired *L. auricularia*, as was verified the following morning by Dr. Pilsbry and Mr. Vanatta when I brought in living material.

Dobbin's Pond was doubtless at one time well kept, since it is situated on private property, but for many years the estate has been largely left to take care of itself, and consequently much debris of refuse, dead leaves and sticks has collected in it. A delta formation of considerable extent occurs at the inlet stream. Few, if any, large water-plants, like pond-lilies, occur, but there is an abundance of algal growth, and pond-scums cover much of the surface.

The *Lymnæa* is a very striking species, both because of the large size of full-grown individuals, and, of the conspicuous dark mottling of the mantle on a light yellow ground as seen through the shell on the younger individuals. The mottling was commonly obscured in the large specimens because of algal growth and dirt. Most of the animals, especially the half-grown ones, were found crawling around in shallow, quiet water on the muddy bottom, along the sides of the delta. The large individuals were generally found attached to floating logs or sticks, and one was seen sticking to grass stems hanging into the water. The majority of the specimens seen were doubtless half-grown individuals from the past season. The shells were found to measure on the average about 17 mm. × 12 mm. The largest shells measured about 28 mm. × 21 mm., but some of Jacob's collection are as large as 30 mm. × 24 mm. One specimen, found dead, has the aperture very beautifully and widely flaring but in most of the material this is not well-developed, or else has been broken away.

Except for a very possible solution, suggested by Mr. Vanatta,

I am rather at a loss to account for the occurrence of this European species at this station. In both the localities previously reported, that of Mr. H. E. Walter at Lincoln Park, Chicago, reported by Mr. F. C. Baker,[1] and that of Mr. R. E. Call, at Flatbush, Brooklyn, N. Y.,[2] the introduction of the species seems to be satisfactorily traced to European plants cultivated in either green-houses or ponds. But neither of these conditions certainly prevail at the present time at the new locality, or to furnish a particularly satisfactory solution. Mr. Vanatta suggests that, from the common use of these snails in aquariums, and the pond being on private property once doubtless well taken care of, it is very possible that the contents of an aquarium may have been transferred to the pond at some time or other. I have since made inquiries and am told that the pond has contained gold fish for many years and that they are at present still frequent there. So the evidence seems rather suggestive that this solution may be correct.

Suffice it, however, that the species has certainly become extensively naturalized in this pond and is actually one of its most prominent molluscan elements.

A NEW OREOHELIX FROM COLORADO.

BY H. A. PILSBRY.

We have recently received from the Hon. Junius Henderson a series of shells collected by Mr. Albert Dakan, which seem to represent a new specific form, which may appropriately be named for Mr. Henderson, whose scientific works include an excellent account of the mollusks of Colorado.

OREOHELIX HENDERSONI, n. sp.

The shell is solid, in shape much like *O. strigosa depressa* (Ckll.), the last whorl angular in front of the aperture, becoming rounded in the last half, shortly descending in front. First $2\frac{1}{2}$ whorls rusty brown; the ground color then becomes very pale brown, and a narrow darker band midway between sutures appears, usually not continuing upon the last whorl, which is lighter, more cream-colored,

[1] Nautilus xv, p. 59, 1901.

[2] Science, New Series, xvi, p. 65, 1902.

beneath. Old shells, by loss of the cuticle, have the last $1\frac{1}{2}$ whorls opaque white with some faint corneous or ruddy stains. Embryonic shell of $2\frac{1}{3}$ whorls, the first somewhat convex, first half whorl smooth, next whorl having fine, close, radial thread-like striæ, last $1\frac{1}{3}$ whorls, having similar striæ and spiral cords, becoming coarse on the last half whorl. This sculpture stops abruptly at the end of the embryonic stage. Neanic and adult stages have rather fine, irregular wrinkles of growth, indistinctly cut by very weak lightly impressed spiral lines which are often almost obsolete. Aperture quite oblique, nearly round.

Alt. 8.6, diam. 16.3 mm.; whorls $4\frac{3}{4}$.

Genitalia as figured for *O. s. depressa* and *O. elrodi*, the end of the penis closely resembling the latter species. Epiphallus tapers at the end. Length of penis 13.6, of epiphallus $5\frac{1}{2}$, of spermatheca and duct $14\frac{1}{2}$ mm.

North bank of Little Thompson Creek, about 10 miles N.-W. of Longmont, Colorado. About 160 living and over 100 dead shells found among roots of "woodbine" in loose limestone talus, at southern point of ridge of Niobrara Limestone, July 26, 1911, by Mr. Albert Dakan.

The strong spiral sculpture of the embryonic shell, abruptly giving place to oblique wrinkle-striæ at the inception of the neanic stage, distinguishes this species from *O. strigosa depressa* (the common Colorado form of *strigosa*), to which it is related by the soft anatomy. The early stages are much as in *O. yavapai*, but that species and its varieties differ strongly by the very short penis.

O. gabbiana (Hemph.), from the Oquirrh Mts., differs from *hendersoni* by the less strong embryonic spiral and more strongly developed, widely spaced spirals of the last whorl, which is carinate or angular. An *Oreohelix* from Glenwood Springs, Colo., which has been provisionally referred to *gabbiana* is probably a form of *hendersoni*. Another form which seems referable here is from Newcastle, Colorado, also collected by Mr. Albert Dakan. The shells are very large, alt. 14.5, diam. 24 mm., with $5\frac{1}{3}$ whorls. It was found on a hillside, out by thousands on a moist day.

I should not complete this account without offering an apology to Mr. Henderson for allowing the name of this species to appear in his article on Oreohelix Colonies (p. 10 of the May number), in advance of any published description. I had written the above account some

months before, when the shells were received. The press of other matter and the habit of giving other contributors precedence over the Editors in our limited space, explain the appearance of the MS. name. To tell the honest truth, I had forgotten that it was not published!

LAND SHELLS OF SOUTHERN FLORIDA.

BY E. G. VANATTA.

[Concluded for p. 22.]

Pavilion Key, Monroe Co., collected by Clarence B. Moore.

Truncatella caribæensis 'Sby.' Rve.

Truncatella c. succinea Ad.

Bifidaria p. hordeacella Pils.

Pelican Key, Lee Co, (A small key near the entrance to Chokoloskee Key), collected by Clarence B. Moore.

Polygyra c. carpenteriana Bld. Some of the specimens lack the internal lamella.

Pine Island, Lee Co., collected by Clarence B. Moore.

Truncatella caribæensis 'Sby.' Rve.

Truncatella c. succinea Ad.

Truncatella bilabiata Pfr.

Truncatella clathrus Lowe.

Practicolella jejuna Say.

Polygyra septemvolva volvoxis Pfr.

Bifidaria p. hordeacella Pls.

Euglandina rosea Fer.

Euglandina rosea parallela Binn.

Zonitoides singleyanus Pils.

Little Pine Key, Monroe Co., collected by Dr. H. A. Pilsbry and C. T. Simpson.

Truncatella caribæensis 'Sby.' Rve.

Thysanophora selenina Gld.

Thysanophora incrustata Poey.

Thysanophora plagioptycha Shutt.

Praticolella jejuna Say.

Polygyra c. carpenteriana Bld.

Microceramus pontificus Gld.

Pupoides modicus Gld.

Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.
Varicella g. floridana Pils.
Vitrea dalliana 'Simp.' Pils.
Guppya gundlachi Pfr.
Succinea floridana Pils.

Big Pine Key, Monroe Co. Collected by H. A. Pilsbry, C. T. Simpson, S. Brown and H. W. Fowler.

Truncatella caribæensis succinea Ad.
Helicina o. clappi Pils.
Helicina tantilla Pils.
Thysanophora incrustata Poey.
Polygyra c. carpenteriana Bld.
Urocoptis poeyana variegata Pfr.
Microceramus pontificus Gld.
Strobilops hubbardi Br.
Pupoides modicus Gld.
Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.
Vertigo variolosa Gld.
Euglandina r. parallela Binn.
Vitrea dalliana 'Simps.' Pils.
Guppya gundlachi Pfr.
Zonitoides minuscula Binn.
Succinea floridana Pils.

Porpoise Point, Monroe Co. Part of a key. Collected by Clarence B. Moore.

Polygyra c. carpenteriana Bld.
Bifidaria p. hordeacella Pils.

Rabbit Key, Monroe Co. Collected by Clarence B. Moore.

Truncatella bilabiata Pfr.

Polygyra cereolus carpenteriana Bld. Many of the specimens from this place lack the internal lamina.

Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.

East Cape, Cape Sable, Monroe Co. Collected by Clarence B. Moore.

Helicina o. clappi Pils.

Seminole Point, Monroe Co. Part of a key. Collected by Clarence B. Moore.

Truncatella caribæensis 'Sby.' Rve.

Truncatella bilabiata Pfr.

Polygyra c. carpenteriana Bld. Many of the specimens in this lot have the internal lamina reduced to a few indistinct nodules.

Bifidaria p. hordeacella Pils.

Bifidaria rupicola Say.

Vertigo rugosula oralis St.

Snake Key, Monroe Co. Collected by Clarence B. Moore.

Truncatella bilabiata Pfr.

Polygyra c. carpenteriana Bld. The specimens from this place are very small and many have a very high spire; they also almost lack the internal lamella.

Bifidaria p. hordeacella Pils.

Snipe Key, Monroe Co. Collected by S. Brown and H. W. Fowler.

Helicina o. clappi Pils.

Sugar-loaf Key, Monroe Co. Collected by H. A. Pilsbry and C. T. Simpson.

Truncatella bilabiata Pfr.

Truncatella clathrus Lowe.

Helicina orbiculata Say.

Helicina o. clappi Pils.

Thysanophora selenina Gld.

Thysanophora incrustata Poey.

Thysanophora plagioptycha Shutt.

Polygyra cereolus Mühlf.

Polygyra c. carpenteriana Bld.

Microceramus pontificus Gld.

Strobilops floridanus Pils.

Strobilops hubbardi Br.

Varicella g. floridana Pils.

Pupoides modicus Gld.

Bifidaria p. hordeacella Pils.

Vitrea dalliana 'Simps.' Pils.

Summerland Key, Monroe Co., collected by Dr. Pilsbry.

Truncatella bilabiata Pfr.

Helicina o. clappi Pils.
Helicina tantilla Pils.
Thysanophora selenina Gld.
Thysanophora incrustata Poey.
Thysanophora plagioptycha Shutt.
Polygyra cereolus Mühl.
Pupoides modicus Gld.
Bifidaria p. hordeacella Pils.
Bifidaria rupicola Say.
Varicella g. floridana Pils.
Vitrea dalliana 'Simps.' Pils.
Guppya gundlachi Pfr.
Zonitoides minuscula Binn.

Torch Key, Monroe Co., collected by H. A. Pilsbry.
Chondropoma dentatum Say.
Truncatella caribæensis succinea Ad.
Truncatella bilabiata Pfr.
Truncatella clathrus Lwe.
Polygyra cereolus Mühlf.
Polygyra c. carpenteriana Bld.
Cerion incanum Binn.
Bifidaria p. hordeacella Pils.

A CONCHOLOGICAL MUSEUM.

Conchologists the world over are interested in the work of Mr. Y. Hirase, of Kyoto, Japan. I doubt if many of them are aware how deeply they are indebted to him. His quarter century of exploration of Japan, by land and by sea, of Corea, and of China, at great personal sacrifice of money and time, has given to science many new species, enriching cabinets all the world round with choice specimens most delicately handled.

He has an extensive and valuable collection of material, containing co-types of the numerous new species brought to light by his efforts. He has plans for the establishment of a Conchological Museum for the preservation and exhibition of this collection. He has in hand cash for about three-fourths of the cost of this undertaking. He still needs $5,000 to finish the work as planned.

Mr. Hirase has expressed to me, in writing, his willingness, in case he is aided by friends in America to the extent of $5,000, to have that sum expended under the supervision of the Trustees of the Doshisha University, in paying the last bills on the plant; to guarantee perpetually the free use of the Museum to students of the Doshisha University; and, in case of the discontinuance of the Museum, to cover into the treasury of the Doshisha for the use of that institution, the $5,000 advanced in aid of his undertaking.

A number of gentlemen have already by word of mouth or in writing indicated a desire to aid Mr. Hirase. Is not the present the best time to carry out this purpose?

Please send your checks—large or small—to Mr. C. W. Johnson, Boston Society of Natural History, Boston, Mass., and thus discharge, in part, the debt due Mr. Hirase. He does not seek pay. He is worthy of our best sympathy and encouragement.

A good museum of the kind he plans will be of immense value to Japan and the world of science, will be a generous crown to his life of hard work, and an encouragement to other lovers of science to emulate his example. MARSHALL R. GAINES.

NOTES.

JAMES JOHNSON WHITE.—We regret to announce the death of our friend Mr. J. J. White, which occurred at Rockledge, Florida, April 26, 1912. He was born in Kittanning, Pa., Feb. 13, 1846. Moved with his parents to Illinois, and later lived in Iowa. In 1885 he removed to Florida, living at Palm Beach, and since 1892 at Rockledge. Mr. White was deeply interested in the study of mollusca, and by correspondence and exchange was known to many of the readers of THE NAUTILUS.

UNIO SILONGWEENSIS, a new name for *Unio vicinus* Preston not Lea. Mr. Bryant Walker has called my attention to the fact that the name *Unio vicinus*, used by me in the Annals & Magazine of Natural History, Ser. 8, Vol. VI, 1910, p. 60, Pl. IV, fig. 7 for a Central African species, was preoccupied by Lea in 1856 (Jl. Acad. Nat. Sci., Philadelphia, Vol. 4, p. 61 and Lea, Obs. on genus Unio, Vol. 6, p. 61). I now propose the new name *Unio silongweensis* for the African form.—H. B. PRESTON.

Mr. C. T. Simpson, the well-known author of the "Synopsis of the Naiades," has arranged for the publication of his "Descriptive Catalogue of the Naiades." This is the companion volume to the Synopsis and was prepared immediately after that work appeared, but for various reasons its publication has been delayed until now. It will contain full descriptions of all the species recognized in the "Synopsis" with comparative notes, keys, etc. Descriptions and references to all the species described since the publication of the "Synopsis" have been added, bringing the subject down to date. Unfortunately it is not to be illustrated, but its publication will for the first time place in the hands of the students of the *Unionidæ* adequate descriptions of all the species in one volume. The cost and in many cases the practical impossibility of obtaining the literature necessary for any accurate work in studying this family has been a great drawback to interesting our American collectors in this group, and, in the absence of a fully illustrated manual, this work of Mr. Simpson's will go far towards supplying the long-felt want of such a publication.

It is expected that the Catalogue will be published early in the fall and will be a volume of from 800 to 1000 pages. It is to be published by Bryant Walker of Detroit, Mich.

New Orleans Mollusca.—The writer with Mr. J. B. Upson, spent the best part of one day the last week of December, 1880, on the shore of Lake Pontchartrain, to the east of Spanish Fort and collected the following species:

Polygyra espiloca Ravenel, common.

Polygyra febigeri Fér., abundant. These two species of Polygyra were determined for the writer by Thomas Bland, in a letter dated May 20, 1881.

Zonitoides arborea Say.

Zonitoides minuscula Binney.

Glandina bullata Gould. Mr. Bland said of this form "*Glandina truncata*, var. (scarcely *bullata*)" several fine specimens were found and some quite young.

Rumina decollata L. One dead example.

Succinea campestris Say, common.

Planorbis lentus Say, scarce.

Segmentina havanensis Pfr., common.

Lymnæa techella Hald., few.

Gnathodon cuneatus Gray, great quantities of dead shells.

Also young specimens of *Viviparus*, *Campeloma* and *Sphærium*.—A. A. Hinkley.



DANIELS: ABNORMAL SHELLS.

THE NAUTILUS.

VOL. XXVI. AUGUST, 1912. No. 4

TWO NEW LAND SHELLS FROM CALIFORNIA.

BY HARRY EDSON

SONORELLA ARGUS, n. sp.

Shell small, depressed, flesh-colored, with a narrow brown band encircling the whorls a little above the periphery. This shell approaches very nearly *S. fisheri*, Bartsch but differs in color, size, and in having a wider umbilicus.

Collected by Mr. A. M. Strong at the Iron Cap copper mine, Argus range, Inyo Co., California. Cotypes in the geological museum of Stanford University, and A. N. S. Phila.

The type measures: maj. lat. 11.3 mm., min. lat. 9.5 mm., alt. 5.7 mm., aperture: maj. lat. 6 mm., alt. 5.3 mm., umbilicus, 2 mm.

EPIPHRAGMOPHORA DUPETITHOUARSI CUESTANA, n. subsp.

Shell large, depressed, polished, obliquely striated, with seven whorls, the last whorl descending slightly near the aperture; deeply and widely umbilicated, showing three volutions; lip slightly reflexed, the ends approaching; color translucent brown, circled near the top of the whorls by a dark brown band edged with yellow. The type measures; greater diam. 26½ mm., lesser diam. 22½ mm., alt. 15½ mm.; aperture, width 16 mm., alt. 13½ mm., including peristome.

This shell is quite similar to *sequoicola* Cooper, but shows no signs of being hirsute and has much coarser striations.

Habitat Cuesta Pass, Santa Lucia Mts., California, collected by Mr. Harold Hannibal. Cotype in coll. A. N. S. P.

Both species will be figured on a future plate.

ABNORMAL SHELLS

BY L. E. DANIELS

In the United States as elsewhere shells are frequently met with that are abnormal to a greater or less degree due to the shell having met with an injury during its growth.

Tryon, in the American Journal of Conchology, Vol. 2, plate 1, fig. 4, figures an abnormal specimen of *Planorbis bicarinatus* Say, from the collection of Charles M. Wheatley Esq., of Phoenixville, Pa., collected by him at Mexico, Oswego Co., N. Y. He says "this is the first American specimen which we have observed to deviate much from the normal form; reversed or scalariform shells being very much rarer in America than in Europe. Some of our species of *Lymnæa* are sometimes slightly scalariform, and in the *Viviparidæ* this is more frequent, but the deviation has never been observed to be very great. *Melantho integra* and *decisa* are both occasionally met with, reversed. I believe that no scalariform *Helix* has been observed in the United States, and the reversed specimens known do not exceed ten or twelve."

Thomas Bland in "Remarks on Certain Species of North American Helicidæ Part 3," 1862, says. "In one of my specimens of *H. tridentata*, after completion of the shell, the animal continued the last whorl about 2 mill beyond the lip, partially reflected the new lip, and added two incipient teeth in advance of the old ones. On plate 4, fig. 20 a sketch is given of the very singular reparation of one of my specimens of *H. septemvolva.* The fracture was behind the aperture, which the animal in consequence abandoned, but it formed a new one by reflecting the outer lip, on the lower part of which it added a small tooth,—it found insufficient space for another partial tooth behind the old aperture."

Binney in the Manual of American Land Shells, page 282, fig. 300, figures an abnormal specimen of *Stenotrema monodon.* The peristome having been broken after the animal's arrival at maturity, a new peristome has been formed somewhat in the rear of the first, and a new parietal tooth added. On page 291 he figures *P. tridentata* with a double peristome.

S. C. Wheat in Bulletin of the Brooklyn Conchological Club figures a specimen of *Pyramidula alternata* Say (fig. 5) found near

Ithaca, N. Y. It is terraced, the suture being placed far below the carina of the preceding whorl. The carina is unusually sharp. Another specimen (fig. 4) was flat, the suture being above the carina and deeply impressed. He also figures an abnormal *P. thyroides* and *P. tridentata.*

H. A. Pilsbry in Mollusca of the Southwestern States I, page 274, say: "Scalariform monsters are not rare, but the inception of that abnormal condition seems to be invariably traceable to an injury of the shells." He figures on plate XXIV fig. 32 a scalariform specimen of *Oreohelix strigosa concentrata* (Dall).

Through the kindness of the Editors of "The Nautilus" I am enabled to present to its readers the following descriptions and figures of abnormal specimens in my cabinet.

Sonorella coloradoensis (Stearns).

East side of Powell's Plateau near Mojave Creek, Grand Canyon, Arizona, 1909.

The shell is scalariform, due no doubt to an injury when young.

Polygyra cereolus (Mühlf). Pl. V, figs. 1, 2, 3.

Long Boat Key. Tampa Bay, Fla., 1900.

One specimen (fig. 1) is normal up to the end of the eighth whorl. A break then occurred, after which the whorl deviates downward, is abruptly recurved, traversing the base across the umbilical region. A nearly normal aperture is formed. In another shell, fig. 2, the whorl adjacent to the aperture dips down for a short distance, subsequently rising above the level of the spire. A well-formed normal example is shown in fig. 3.

Polygyra tridentata (Say.) Pl. V, figs. 10, 11.

Wyandotte, Crawford Co., Ind., 1903.

This shell had been broken back of the peristome, the animal instead of repairing the break retreated eight mm., and formed a new peristome and parietal tooth.

I succeeded in extracting the animal without breaking the outer peristome although the attachment to the shell was very slight, this (the outer) peristome being dead.

Polygyra inflecta (Say). Pl. V, fig. 6.

Jeffersonville, Clark Co., Ind., 1903.

Peristome malformed, the callus wanting above, teeth rudimentary.

Polygyra profunda (Say). Pl. V, fig. 12.

Near Hoge's Bridge, Morris, Ills., 1910. This shell had formed a peristome, then carried the shell forward twelve mm. and formed a second peristome. The shell between the two peristomes is white, has no epidermis, and has a rough appearance.

Polygyra albolabris (Say). Pl. V, fig. 13.

North Vernon, Jennings Co. Ind., 1903. In one specimen the last two-thirds of the body-whorl has been injured, resulting in a linear furrow running parallel with the suture. Another shell (fig. 13) was injured before forming the peristome which has two projecting points. The umbilicus is partly open.

Polygyra exoleta (Binn). Pl. V, fig. 14.

Mitchell, Ind., 1901. Peristome malformed, bisinuate.

Polygyra elevata (Say). Pl. V, fig. 15.

Mitchell, Ind., 1901. This specimen appears to have attained its growth, and apparently had partially formed the peristome, for at the proper place a parietal tooth was formed which is about one-half grown. The last half of the body-whorl was then broken off and a new peristome and parietal tooth were formed.

Polygyra hirsuta (Say). Pl. V, fig. 7.

Bass Lake, Starke Co., Ind., 1902.

Aperture malformed exactly as in the specimen of *elevata* described above, the last part of the whorl having been broken off after the parietal tooth was half formed, and a new peristome and tooth were developed a fourth of a whorl behind the original aperture.

Polygyra fraterna (Say). Pl. V, fig. 5.

Brookville, Ind., 1903.

This shell had formed a peristome and parietal tooth, then carried the shell forward five mm. and formed another peristome and parietal tooth, the last aperture being small. The original parietal tooth remains standing deep within. This shell belongs to the same class of monster described above for *P. profunda.* It is unlike most abnormal shells which are due to breakages, for here a second period of growth is superposed upon a normally formed and completed shell.

Holospira bilamellata Dall. Pl. V, figs. 8, 9.

Hacheta Grande Mts., New Mex., 1910.

This shell was normal up to the penultimate whorl, when without any apparent injury it contracts suddenly to three-fifths the former diameter. The last whorl enlarges again somewhat, and is nearly normal. A normal example is figured for comparison, fig. 9.

Holospira n. sp.

Dragoon Mts., Arizona, 1910.

A strongly ribbed new species which will shortly be described by Messrs. Pilsbry and Ferriss. The shell was broken at the end of the eighth whorl. The next $2\frac{1}{2}$ whorls are ribless; ribs then begin again irregularly, finally becoming normal on the last whorl.

Oreohelix strigosa depressa (Ckll) Pl. V, fig. 16.

Jacob's Canyon, Kaibab Mts., Ariz., 1910.

A scalariform specimen, the last whorl free in front. There are no signs of this shell ever having been injured in the least. Normal shells are shown in figs. 18, 19.

A reversed specimen from Powell Plateau, Grand Canyon, Ariz., 1909 is shown in fig. 17. J. H. Ferriss of Joliet, Ills., has a similar one.

Planorbis bicarinatus Say. Pl. V, fig. 4.

Lily Lake, La Porte, Ind., 1909.

This extraordinary specimen has the general shape of *Viviparus intertextus* with a small umbilicus and a deeply channeled suture. The aperture is oval with expanded peristome. So abnormal in shape is this specimen that it takes some study to see the common *Planorbis bicarinatus* in it.

Campeloma decisum (Say).

Pine Lake, La Porte, Ind., 1906.

While reversed specimens of *Campeloma* are not rare the novelty of this one is that it contained three young, all of which are reversed.

NOTE BY THE EDITOR.—Little has been published upon abnormal shells in this country, though there must be abundant material scattered among collections. The notes given above afford suggestions for a classification of teratologic shells into several classes as follows:

1. Shells in which growth has been resumed after the completion of a normal peristome, figures 5 and 12.

2. Shells in which broken parts have been repaired or restored.
   2*a*. Broken peristome repaired, figs. 13, 14.
   2*b*. A new peristome developed behind the original one after breakage, figs. 7, 10, 15.
3. Shells coiling abnormally.
   3*a*. Shell sinistral, fig. 17.
   3*b*. Shell scalariform, figs. 4, 16.
   3*c*. Shell irregularly coiled, figs. 1, 2, 8.

Part of these cases are always due to breakage; others, such as no. 1 and 3*a* to more obscure constitutional conditions. Scalariform shells may be due to several causes. The subject is of some importance from an evolutionary standpoint because the modifications denoted by classes 1, 3*a*, 3*b* and 4*c* have in some cases become actually characteristic of species or genera.

Collectors having material enlarging our knowledge of the subject as illustrated by Mr. Daniels are invited to make it known.

Figures 1 to 9 of plate V are about double the natural size.

NEW CUBAN LAND SHELLS: I. ANNULARIA AND CHONDROPOMA.

BY H. A. PILSBRY AND JOHN B. HENDERSON, JR.

The following species will be illustrated in connection with the second paper of this series.

ANNULARIA RAMSDENI n. sp.

The shell is openly umbilicate, wider than high with conic spire. Last whorl brownish-corneous or whitish, the spire light red, whorls $4\frac{1}{2}$, strongly convex, the last tubular. Suture narrowly channeled, the axial threads forming little points where they pass over the raised edge of the channel. Sculpture of closely crowded lamina-like axial threads, their intervals having one or two smaller threads. There are a few low spiral cords within the umbilicus and a stronger one near its opening. The first $1\frac{3}{4}$ whorls are smooth and well raised. Aperture oblique, circular, pale red within the lip. Peristome expanded, duplicated by a thin outer lamina on the columellar and upper sides, dilated in a forwardly concave lobe above, which is appressed to the preceding whorl.

Operculum of three whorls, each nearly covered by a white, level lamella arising from the inner margin, spreading nearly to the outer.

Male: alt. 11.8, diam. 13 mm.

Female: alt. 12.5 diam. 15 mm.

El Jique de la Argolla, Barrio de Rio Seco de San Antonio, 15 miles northeast of Guantánamo, under stones, Charles T. Ramsden 12–24-'11.

The cotypes figured were observed coupling Mar. 19, 1912.

This species is closely related to *A. pseudalatum*. It is larger, with more rapidly enlarging whorls, the aperture relatively larger, its diameter more than half that of the shell, while in *A. pseudalatum* it is less than half; and it differs especially in sculpture, the major axial threads being much more numerous and closer together. *Annularia* (or *Choanopoma*) *alatum* Pfr. is a more opague, solid shell with no sutural channel, coarser, unequal striæ, and a differently formed lip.

In both *A. ramsdeni* and *A. pseudalatum* the last whorl is free in front just before finishing the lip, but the additional thin lamina and superior appendage of the lip connect with the preceding whorl in the fully adult stage. This species is named for the discoverer, Mr. Charles T. Ramsden.

Dr. W. H. Dall has called attention to the fact that the generic name *Annularia* has precedence over *Choanopoma* for species of this type.

ANNULARIA PSEUDALATUM Torre, n. sp.

The shell is openly umbilicate, wider than high, with a low conic spire; corneous; whorls $3\frac{3}{4}$ to $4\frac{1}{3}$, very convex, the last tubular, barely in contact with the preceding at the aperture. Sculpture of straight, slightly retractive and very thin axial threads, separated by intervals in which there are several (three to five) delicate axial striæ. There are several weak spiral cords within the umbilicus, the lowest one stronger than the others. The first $1\frac{3}{4}$ whorls are smooth. Aperture oblique, circular, the peristome thin, narrowly expanded, duplicated by a thin outer lamina along the columellar, parietal and upper margins, this lamina expanding in a forwardly concave lobe above. Operculum of about 3 whorls, similar to the preceding species.

Alt. 7.2, diam. 10.2 mm.

Alt. 8, diam. 10 mm.; diam. of aperture, with peristome 4.5 mm.

Alt. 7, diam. 9 mm.

Farallones east of Hermitage, Charles Wright.

This form is described and figured from specimens determined by Professor de la Torre, for comparison with the preceding. It is not uncommon in the older collections, under the name "*Choanopoma alatum Pfr.*" Pfeiffer's species is larger, more solid, opaque fleshy-white, with the interior orange; the aperture is much more oblique and the sculpture entirely different.

Chondropoma carenasense n. sp.

The shell is perforate, tapering-cylindric, truncate, about $3\frac{1}{2}$ whorls remaining; very pale brown, with small often indistinct dark spots arranged in vertical series at irregular intervals, sometimes having 5 or 6 narrow, faint, spiral bands, with spots as above. Sculpture of many fine spiral cords, 10 to 14 on the penult whorl, crossed by more closely set vertical lamina-like threads which have small nodes where they cross the spirals, and finely crenulate the suture, where each stands separate. Whorls convex, the last ascending a little in front. Aperture ovate; peristome expanded, built out in a narrow flat margin, widest as the columeller margin and at the posterior angle, where it ascends in a very short lobe or obtuse point; parietal margin appressed to the preceding whorl. The operculum retracts about one-fourth of a whorl within, is light brown, with a spiral convexity just outside the suture.

Length 12.5, diam. 7.5 mm.; length of aperture with peristome 6 mm.

Length 12, diam. 7 mm.; length of aperture with peristome 5.7 mm.

Length 10.3, diam. 6 mm.; length of aperture with peristome 5 mm.

Cayo Carenas, in Cienfuegos Bay, Pilsbry, 1903, and Henderson 1910.

C. candeanum (Orbigny) tapers more, to a narrower truncation, and the sutural denticulation is more widely spaced. In *C. carenasense* every vertical thread denticulates the suture, none being united or dropped out there. It is an abundant snail on Cayo Carenas, a tiny islet now nearly covered with residences.

CHONDROPOMA WILCOXI n. sp.

The shell is perforate, ovate-twisted, truncate, 3 to $3\frac{1}{2}$ whorls remaining. Pale fleshy buff, indistinctly marked with spiral series of small brown dots. On the last whorl there are 6 such rows, three more distinct; on preceding whorls there are four rows of dots; the suture has a narrow whitish margin below. Sculpture of many subequal spiral threads intersected at right angles with narrower, slightly weaker, axial laminæ, producing an evenly latticed surface; on the last whorl there are about 40 spirals. The whorls are rather strongly convex; last whorl ascending slightly to the aperture. The aperture is vertical, ovate, somewhat acuminate above, brown inside. The peristome is white; it expands a trifle, and is somewhat thickened on the face, which is narrow and convex. At the upper angle it is a very little dilated.

Length 19, diam. 11, length of aperture $8\frac{1}{4}$ mm.

Rosario, east shore of Ensenado Cochinas, in forest. Henderson and Simpson.

STRANGE NAMES FOR OLD ACQUAINTANCES

BY C. HEDLEY AND H. A. PILSBRY.

It is pleasant to find that there was a time when the editor was not regarded as the natural enemy of his contributor. Thus when Dr. Johann Hermann, aforetime Professor at Strassburg, presents an essay on conchology to his friend the editor of "Der Naturforscher" he greets him cordially and at the conclusion takes affectionate leave.

Did this quaint old essay not disturb our nomenclature, we should not need to disturb the dust which for nearly a century and a half has gathered on this venerable periodical. But the names which Hermann gave have escaped the attention of most modern writers, yet they are presented regularly and precede several now in use. Like Marryat, the novelist, our author made but a single contribution to molluscan literature. In the professor we find a careful worker: not only did he fulfill the modern requirements of nomenclature, but he embellished his paper by a colored engraving illustrating five species.

Two species of Spondylus are contrasted and are named respectively *S. mediterraneus* and *S. americanus* (Der Naturforscher XVI, 1781, p. 51). The former is evidently *S. gœderopus* Linné. The latter must be the common spiny species of the West Indies commonly known as *S. americanus* Lamarck, for which Doctor Dall (Trans. Wagn. Inst. III, part iv, 1898, p. 759), after a consideration of the extensive synonymy, adopts the name *S. echinatus* Martyn, 1784, which apparently should now give place to *S. americanus* Hermann, 1781.

TURBO CANALICULATUS, Hermann.

Der Naturforscher, XVI, 1781, p. 52, Pl. ii., fig. 1.

This species is clearly the ordinary large *Turbo* of the West Indies, usually known as *T. spenglerianus* Gmelin. The latter author, though he cited Hermann's work (Syst. Nat., xiii, 1790 p. 3571), made the double error of using his name in another sense and of bestowing a subsequent name upon Hermann's species. Both the species termed *Turbo canaliculatus* by Gmelin and Kiener respectively, are Oriental.

NERITA CANCELLATA, Hermann.

Der Naturforscher, xvi, 1781, p. 56, pl. ii, fig. 8. 9.

Here again Gmelin has misunderstood the work of Hermann. After quoting (p. 3670) the above reference, he adds from Chemnitz accounts of two distinct shells, the one *Natica sulcata* Born, a West Indian *Natica*, the other *Vanikoro cancellata*, Lamarck. In his turn Lamarck was thus led to confound these species, a confusion which was unraveled by Deshayes, (An. s. vert. [2] viii, 1838, p. 644). Hermann's illustration well represents *Nerita sulcata*, Born (Index Mus. Caes. Vind., 1778, p. 416, Pl. 17, fig. 5, 6), which is maintained by three years' priority.

BUCCINUM SOLUTUM Hermann.

Der Naturforscher, XVI, 1781, p. 53, Pl. ii, fig. 3, 4.

Dr. W. H. Dall kindly gave us the benefit of his wide knowledge of this boreal group. Hermann's figure appears to him to represent a short form of *Chrysodomus heros* Gray (Proc. Zool. Soc. 1850, p. 14, Pl. VII) a species erroneously united by Kobelt, Conch. Cab. Fusus, 1881, p. 61, to *Neptunea fornicata* Gray non Gmelin. Hermann's name should therefore be maintained.

BUCCINUM STROMBOIDES Hermann.

Der Naturforscher, XVI, 1781, p. 54, Pl. ii, figs. 5, 6.

This has been recognized as a fossil from the Paris basin and was used by Mörch (Journ. de Conch. VIII, 1858, p. 43) as the type of the genus *Cryptochorda*.

MUREX DESPECTUS Linnê.

Hermann, Der Naturforscher, XVI, 1781, p. 54, Pl. ii, fig. 7.

Here Hermann professes to figure a sinistral *Chrysodomus despectus* but Dr. Dall points out to us that the illustration better suits a sinistral *C. antiquus*.

A NEW SPECIES OF ANCULOSA.

BY A. A. HINKLEY.

ANCULOSA MINOR, n. sp. Pl. 1, figs. 7, 8.

Shell small, depressed, thick, light olive in color, whorls five, those of the spire flat; periphery angulate or carinate; lines of growth fine and cut by fine revolving impressed lines, giving the surface of the shell a reticulate appearance under the microscope; peristome straight and thin; aperture ovate, columellar callus thick, wide, dark colored or white, extended across the parietal wall; umbilical region indented. Three specimens measured length 33, diam. 27, length, 30, diam. 24, and length 30, diam. 25 hundredths of an inch.

Habitat—near Florence, Alabama, in the Tennessee river, near the south bank, in a shaded situation where there was but little current and a good many leaves in the water and on the bed of the stream. A. A. H. Nov. 10, 1904. Cotypes in coll. Hinkley and A. N. S. Phila. (No. 105866).

This little species has been referred to *Anculosa carinata*, but I am sure they are distinct; compared with that species they are smaller, thicker, spire not as elevated, aperture more regular oval and the columellar callus thicker. From *A. costata* it differs in being more elevated, thicker, whorls flat and suture linear, aperture smaller.

The young of this species is very much like the young of *A. crassa* Haldeman, but are more elevated, have no bands and the impressed lines are more strongly marked; nor does *crassa* have the second carina. The young are quite close to those of *A. praerosa* and *tintinnabulum* (see the Nautilus vol. 21, page 110, plate 10, figures

1, 7, and 8) *A. minor* is a little more elevated than the former, has 5 flat whorls and the space between the carinæ is on the same plane as the rest of the base of the shell, the revolving impressed lines are more prominent than on either of the above species, the mature shell has more in common with *A. melanoides*, but that species is more elevated with no carina or impressed lines.

A NEW FLORIDIAN VIVIPARUS.

BY H. A. PILSBRY AND C. W. JOHNSON.

VIVIPARUS WALKERI n. sp. Plate III, figs. 6, 7.

The shell is perforate, moderately solid, globose-conic, dull yellowish-brown, lighter beneath, with three or four very indistinct purplish-brown bands, or dull purplish-brown without bands; faintly striated. Spire short, much eroded and whitish in adults, which retain about 3 whorls. Whorls very convex, somewhat flattened horizontally below the very deep suture. The aperture is large, slightly oblique, ovate, livid and showing the bands within by transmitted light, when any are present.

Length 26, diam. 22, length of aperture 16½ mm.

Length 26, diam. 21, length of aperture 15 mm.

Types no. 70,053 A. N. S. P., collected by Pilsbry and Johnson in Juniper Creek, Lake County, Florida, 1894.

This species differs strongly from other Viviparas of Florida by its broad, short form. It was taken in considerable numbers in the type locality, but was not found elsewhere. *V. waltoni* Tryon is the prevalent form in Lake George, and *V. georgianus* in the St. John's River. It is named in honor of Mr. Bryant Walker, whose work on fresh-water shells is well-known to readers of THE NAUTILUS.

NOTES.

WE have to record the recent death in Honolulu of the veteran Hawaiian conchologist, D. D. BALDWIN. A sketch of his life and work will be given later.

MR. CHARLES HEDLEY of the Australian Museum has been studying the museums of this country during June and July.

MESSRS. J. B. HENDERSON, JR., and GEORGE H. CLAPP have returned from a cruise in the western Bahamas on the yacht *Eolis*. DR. PAUL BARTSCH has also visited the Bahamas on the Carnegie Institution yacht *Anton Dohrn*.



TORRE: CUBAN UROCOPTIDÆ.
(× 2)

The Nautilus.

Vol. XXVI. SEPTEMBER, 1912. No. 4

NOTES ON THE VALIDITY OF HELIX (EPIPHRAGMOPHORA) OREGONENSIS LEA.

BY HENRY M. EDSON.

[The article following is probably the last written by Mr. Edson whose untimely death was recorded in The Nautilus for March, p. 132.]

This species seems to be recognizable, but has been buried for a long time in the synonymy of *Epiphragmorphora dupetithouarsi* Desh. It was described by Dr. Lea[1] in 1836 from specimens collected by Mr. Nuttall in Oregon; after a careful reading of the original description and comparing it with Pfeiffer's description of *mormonum* I am convinced that *Epip. mormonum* is synonymous.

The type locality of *mormonum* was Mormon Island in the American River in Sacramento Co., Cal; it has also been collected at Redding, Shasta Co., by McGregor,[2] in Tuolumne Co., by Hemphill,[3] and at Klamath,[4] Oregon. Dr. Lea's description of *oregonensis* follows, "Shell subcarinate, thin, smooth, reddish-brown, double-banded on the carina, above slightly convex, below somewhat inflated; Habitat, Wahlamet near its junction with the Columbia River; Diam., 6, Length .4 of an inch."[5]

Dr. Lea also states that the specimen was immature so that of course the description would be somewhat defective. "That part of the

[1] Lea Obs. Vol. II, 1836, p. 100.

[2] Nautilus, XIII, 1899, p. 64.

[3] Nautilus, XIII, 1899, p. 128.

[4] Washington State Museum, fide H. Hannibal.

[5] Diam. 15 mm., Alt. 9 mm.

specimen which has the epidermis presents in it, under the microscope, a remarkable waved appearance." He also says, "shell subcarinate," and his figure[1] shows it to be so. This feature alone would eliminate *Epip. dupetithouarsi*, for although the very young shells of *dupetithouarsi* are planulate the whorls are always rounded and especially in a shell of the size of Dr. Lea's specimen, which must have been over half grown. This taken with the subsequent collection of *mormonum* Pfr., at Klamath, Ore., which is close to the original locality of *oregonensis*, and the fact that Dr. Lea also described *Polygyra townsendiana*,[2] from the same locality at the same time, makes it pretty certain that *Helix mormonum* Pfr. was the shell Dr. Lea described, and not a specimen of *dupetithouarsi*; While Dr. Lea's description was based on an immature shell I consider it quite recognizable, as Nuttall was not likely to confuse his localities, and *dupetithouarsi* does not by any chance get so far north. Dr. Lea described *Helix oregonensis* in 1836; it thus has precedence of *mormonum* Pfr. by twenty-one years, as Dr. Pfeiffer did not describe *mormonum* until 1857.

The synonymy would stand as follows:

1836. *Helix oregonensis* Lea, Obs. II, p. 100, xxiii, fig. 85; Pfeiffer Mon. Hel. Viv., I, 1848, p. 428.

1857. *Helix mormonum* Pfr., Proc. Zool. Soc., 1857, p. 109; Mon. Hel. Viv., iv. p. 276; Fischer & Crosse, Moll. Mex. et Guat., 1870, p. 251.

1885. *Arionta mormonum*, W. G. Binney, Manual Amer. Land Shells, Bull. 28, U. S. N. M., p. 14.

1897. *Epiphragmophora mormonum* Pfr., Pils. Naut. xi, p. 48. Keep, West Coast Shells, 1911, p. 277.

Note.—As the type of *Helix oregonensis* Lea is now in the U. S. National Museum we requested Dr. Paul Bartsch to compare it with *mormonum* in view of the identification advocated by Mr. Edson. He writes as follows. "I have been looking over Mr. Edson's manuscript and also Lea's type of *Helix oregonensis*. I consider that *oregonensis* has nothing to do with *dupetithouarsi* Desh. The latter species in the young stage has a well rounded periphery, while in *oregonensis* we have a strong keel at the junction of the dark and

[1] Lea Obs. II, 1836, pl. xxiii, fig. 85.

[2] Lea Obs., II, 1836, p. 99.

light band. The anterior light band is about as wide as the dark. The posterior one is scarcely indicated, being represented by a mere line. The nepionic whorls of the two shells are also different: that of *oregonensis* being papillose while that of the *dupetithouarsi* is wrinkled. The young *oregonensis* is also much larger considering the same number of whorls in the two species. Then too, the sculpture of the adult whorls is quite different; in *oregonensis* we have a surface which is covered by a crinkly epidermis, the crinkles placed obliquely. The lines of growth are irregular and not well marked, while in the *dupetithouarsi* we have regularly impressed lines and none of the crinkled epidermis, the latter being smooth; *dupetithouarsi* has a well-rounded spire while in *oregonensis* it is flattened, etc. I believe that *oregonensis* comes nearer to the group of *E. mormonum* than *E. dupetithouarsi*, You have the types of *cala* and *buttoni*, so I will leave the comparison of them with you."

It seems from the above that we will have to write *oregonensis* in place of *mormonum*, but whether Lea's name applies to the typical middle Californian *mormonum* or to one of the described subspecies remains in doubt. Adult specimens from Oregon are needed to settle this question.

H. A. P.

LAMPSILIS VENTRICOSA (BARNES) IN THE UPPER POTOMAC DRAINAGE.

BY A. E. ORTMANN.

Some time ago the writer reported (Mem. Carnegie Mus., 2, 1906, p. 373) that a western species of crawfish, *Cambarus obscurus* Hagen, is found in the Potomac drainage, in Wills Creek, at Ellerslie, Allegany Co., Md., and Hyndman, Bedford Co., Pa. This fact was discussed on p. 445 (l. c.), and explained in general terms as an "artificial introduction by human agency." I may add that this crawfish apparently has increased in numbers since its first discovery (August 10, 1904, and May 8, 1905), for on September 7, 1909, I found it very abundantly at Hyndman (for about a mile in the creek above the tannery). And further, on May 6, 1912, this species was seen in the South Branch Potomac at Romney, Hampshire Co., W. Va. Although only a cast-off shell was found, this was of a male of the first form, and positively established the presence of this species at this locality.

A case parallel to this has now come to light: the presence, in the Potomac system, of a western mussel, *Lampsilis ventricosa* (Barnes). This is a form common, for instance, in the Ohio drainage in western Pennsylvania and West Virginia, but it is *not* found in any of the Atlantic river systems investigated by the writer (Delaware, Susquehanna, James). The following records are at hand:

September 4, 1909. Potomac River, Hancock, Washington Co., Md. Here it was in the smaller branch of the river in great numbers and in all sizes, from 41 mm. long upward. About two dozen were taken.

May 9, 1911. South Branch Potomac River, Southbranch, Hampshire Co., W. Va. About a dozen were found within the first mile of the South Branch above its junction with the North Branch. All were of medium size.

August 16, 1911. Shenandoah River, Harper's Ferry, Jefferson Co., W. Va. A single male was found, below medium size.

May 6, 1912. South Branch Potomac River, Romney, Hampshire Co., W. Va. About a dozen were found in a very small branch of the river. Some were rather large, but most of them were quite small (smallest 20 mm. long). These small ones were found all together in fine gravel at the head of a riffle, in shallow water, attached to small pebbles by their byssus.

In other parts of the Potomac drainage, thoroughly investigated, this species was *not found*, and this is especially true for the headwaters of the Shenandoah River and the northern tributaries of the master stream (Antietam, Conococheague and Great Tonoloway Creeks in Pennsylvania and Wills Creek in Maryland).

It is probable that this species will turn up elsewhere in the Potomac. The localities known at present are all to the West of the Blue Ridge Mountain, that is to say, within the Great Allegheny Valley and the Allegheny Mountains. It should be noted, that its representative Atlantic form, *Lampsilis cariosa* (Say), which is so common in the Delaware and Susquehanna, is not found in this region, but is present farther below in the Potomac: I have specimens from Cabin John, Montgomery Co., Md.

There is no question that we have to deal with a form of *Lampsilis ventricosa*. It is true that the Potomac form represents a peculiar type in so far, as all the specimens collected are rather uniform in shape, but the same shape is frequently met with in western speci-

mens, although there is a much greater range of variation in the latter. The Potomac specimens are rather high and short, and their size remains far below that of western *ventricosa*. My largest specimen (a male from Hancock) is 114 mm. long, 82 mm. high, and 58 mm. in diameter, while I have many individuals from western Pennsylvania, which are 150 mm. long and over. Further the posterior ridge of the shell is in the Potomac form a little more distinct than in the average of the western specimens, but again in this character there are western specimens, which agree with the eastern.

In all other particulars the two forms agree, and also the color and color markings (rays) of the epidermis are the same, and quite variable. But on account of the small size, the shorter and higher outline, and somewhat more distinct posterior ridge, we might regard the Potomac-race as a variety, which I propose to call: *Lampsilis ventricosa cohongoronta* (an old Indian name of the Potomac). I may add, that I have investigated the anatomy, of males, sterile and gravid females, and that there is no difference whatever. Also the glochidia are the same.

It is evident that this shell is, zoögeographically, out of place in the Potomac, and, considered together with the case of the western crawfish mentioned above, we must conclude that both are artificial introductions into this river, probably quite accidental, and the idea is suggested that *the shell might have been transported in the larval stage, as glochidium, during its parasitic life upon certain fishes.* Western species of fishes have repeatedly been transplanted into the Potomac system. H. M. Smith and B. A. Bean (Bull. U. S. Fish Comm. 18 (or 1898) 1899 pp. 179–187) mention no less than seven species (disregarding four European), and among them are the following: Rock Bass (*Ambloplites rupestris* (Raf.)), Large mouth Black Bass *Micropterus salmoides* (Lac.), and Crappie (*Pomoxis annularis* Raf.).

Now just these have recently been given by G. Lefevre and W. C. Curtis (Bull. Bur. Fisheries, 30, 1912, pp. 157 and 160) as most susceptible to infection with the glochidia of species of *Lampsilis*, among them *Lampsilis ventricosa*. The introduction of Black Bass was in the Shenandoah River in 1889; that of the other two fishes in various parts of the Potomac and the Chesapeake-Ohio Canal chiefly in 1894. I do not hesitate to attribute to this fact, the transplantation of certain western fishes, the accidental introduction of

glochidia of this mussel, which found in this region congenial environment, and thus the mussel became established and is propagating. In the western waters, *L. ventricosa* prefers similar ecological conditions: rivers and streams of medium or small size, with riffles and rapidly flowing water, and gravelly bottom.

Very likely the same cause accounts for the accidental introduction of *Cambarus obscurus;* in seining for bass, for instance, in western Pennsylvania, it would be quite impossible *not* to catch a number of this crawfish.

This is an interesting instance of the stocking of a stream with a mussel species by the help of fish. Lefevre and Curtis (l. c., p. 192) discuss this way as a possible means of mussel transportation; however, they express some doubt as to its practicability. But here we have, apparently, the demonstration that this *is* possible, and having happened once quite by accident, it is to be expected that it also might be successfully accomplished when intended and done with the proper care.

NEW ENGLAND NOTES.

BY REV. HENRY W. WINKLEY.

A careful search at Wood's Holl and Chatham, on Cape Cod, reveals more specimens of the species described as *Pyramidella* (*Sulcorinella*) *bartschi.* These have been compared with the type by Dr. Bartsch as well as myself, and they show it to be an *Odostomia.* The name is therefore changed to *Odostomia* (*Evalea*) *bartschi.* Mr. W. F. Clapp has also found it within the cape.

Short trips to a few points reveal *Odostomias*, and their distribution should be noted. The four species, *Odostomia trifida, bisuturalis, winkleyi* and *P. fusca*, are to be found at low tide in inner waters as follows: In a small creek in the marsh near the station at Rowley, Mass. (This same spot is the most northerly locality where I have found *Paludestrina salsa.*)

At old Newbury, Mass., in the Parker River, in eel-grass patches, is another colony. The third is in Great Bay, New Hampshire, at a spot half way between Dover and Portsmouth.

Last summer I obtained the best lot of *Pyramidellidæ* I have ever seen. My stay at Chatham was short, and confined to low-tide col-

lecting. One small area revealed excellent specimens of *P.* (*Syrnola*) *fusca* and *producta*, with one specimen of *winkleyi*. At Wood's Hole I found much material of species already recorded and a few of *Turbonilla verrilli*. I also got a good series of *Turbonilla sumneri*. *Eulima stenostoma* and *conoidea* occur at Wood's Holl, and a few specimens of *Tellimya ferruginosa* were found at the locality where Verrill located them years ago. While *Turbonilla* is ordinarily to be found only by dredging, it does occur at some places at low tide. A chance to observe leads me to think it is a burrowing shell, hence collectors will do well to sift sand in seeking this form. Another suggestion for collectors is, watch carefully for *Eulima*. I accidentally discovered in sifting that *E. stenostoma* after being sifted from the mud has a tendency to float on the surface of the water, as does also *Solenomya*. In working at low tide and sifting there is a temptation to float off dead eel grass, leaving the shells at the bottom of the sieve. Watch carefully, lest *Eulima* floats away.

A trip made the day before writing this revealed a few *Odostomia trifida* in the creeks back of Nantasket Beach, Mass. I cannot too strongly urge on field workers the duty of sweeping with a dip-net in places where eel grass abounds, and be sure to take at least an inch below the surface of the mud or sand in which it grows. Shells are abundant there, and occasionally rare forms, *Odostomia gibbosa*, for example, of which I obtained two more specimens last summer.

NEW CUBAN UROCOPTIS OF THE U. CINEREA GROUP.

BY DOCTOR CARLOS DE LA TORRE.

Urocoptis (*Gongylostoma*) *cinerea* Pfr. Plate vi, figs. 14, 15.

This species has hitherto been known only from the original description in the Conchylien Cabinet and its copy by Pilsbry in the Manual, xv, page 273. The exact locality of its occurence in Cuba has also been unknown, collectors apparently having missed it. In my recent conchological excursions through the central part of the island I obtained it both in its typical form and through a series of varieties and mutations which appear to define a natural group containing a number of species and subspecies.

Specimens agreeing closely with Pfeiffer's description generally

have 9–10 whorls; if the shell is entire (rare) there are 22–24 whorls, the two embryonic ones being smooth, corneous and somewhat bulbous. The radula formula is 7.2.1.2.7, there being 19 teeth in each transverse row; the inner two laterals on each side are much larger than the others, the third tooth becoming abruptly smaller than the second; the outer laterals are capped with a low ledge in place of cusps. The internal axis of the shell is encircled by a low thread below, which in the upper whorls is lamelliform and bears very delicate filament-like spines. This last character determines the position of this species in a group along with *U. wrighti*, *baculum* etc., although the radula frequently indicates the Section *Teterentodon*.

Central Cuba on walls and stones at Cerro de Guajabana near Caibarien, Santa Clara Province.

Urocoptis (*Gongylostoma*) *livida*, *n. sp.* Plate vi, figs. 21, 22.

Shell slenderly fusiform, somewhat swollen in the middle third, tapering to a narrow truncation above; livid-colored, fleshy above and bluish or purplish below. Surface glossy, sculptured with obliquely subarcuate riblets *not flattened on the lower whorls*. Remaining whorls 10–12, somewhat convex, the last free in a descending curved neck scarcely carinated. Aperture as in *U. cinerea* but paler within. The spiral thread of the axis is stronger and perceptibly spinose in the middle and upper whorls.

Longitude 16, diameter 2.7, aperture diameter 2.1, whorls 11½.

Longitude 17, diameter 2.9, aperture diameter 2.3, whorls 12.

Longitude 12.5, diameter 2.2, aperture diameter 2, whorls 11.

Central Cuba: On walls and large stones at El Palenque de Taguayabon and La Puntilla, near Remedios in the Province of Santa Clara.

This species differs from *U. cinerea* Pfr. by its more fusiform shell, livid color, glossy surface and by its regular sculpture; also by its stronger internal axis.

Urocoptis livida occulta n. subsp. Plate vi, fig. 23.

Differs from the type in sculpture and color; the surface is glossy and nearly smooth, the riblets being obsolete in the middle whorls but well raised on the free curved neck; the color is livid pinkish above and violet below with scattered corneous streaks, peristome white.

Longitude 15, diameter 2.9.

Longitude 12, diameter 2.3.

Longitude 13, diameter 2.5.

Central Cuba: Under stones at the sides of the road between Remedios and Taguayabon in the Province of Santa Clara.

Urocoptis (*Gongylostoma*) *pallidula* n. sp. Plate vi, figs. 18, 19, 20.

Belonging to the *cinerea* and *livida* group, it differs in its sculpture and color; the surface is obliquely subarcuately striate, the delicate striæ being closely spaced to the last whorl, where they become coarser and more widely spaced riblets; the color is fleshy to pale yellow, with scattered corneous streaks, sometimes bordered with an opaque white patch. In general shape, though slightly fusiform, it resembles typical *U. livida* with its descending curved neck and moderate aperture. The internal axis is somewhat sinuous, and the lamelliform spiral thread is well developed for this group, lying near the base in each whorl.

Long. 18.3, diam. 3, ap. diam. 2.6, whorls 13.

Long. 15, diam. 2.4, whorls 13.

Long. 15, diam. 2.2, whorls 20 (entire).

Central Cuba: On walls and stones at Mogotes de las Jumaguas, very near Sagua la Grande, in the Province of Santa Clara.

Urocoptis (*Gongylostoma*) *fortiuscula* n. sp. Plate vi, figs. 24, 25, 26.

Resembles *U. livida occulta* in color. The median whorls are finely striate, but it differs from the last in its stronger and rather swollen shell, the last whorl less or not at all descending, and the aperture being larger. The internal axis is slightly sinuous and the lamelliform spiral thread is near the base, as in *U. pallidula*.

Long. 17.5, diam. 3.8, ap. diam. 3, whorls 11.

Long. 14.5, diam. 3.2, whorls $9\frac{1}{2}$.

Some specimens from the same locality are much smaller:

Long. 10, diam. 2.6, ap. diam. 2, whorls 8.

Long. 12, diam. 3, whorls 9.

Central Cuba: On and under stones at Casimba de Buenavista, district of Malpaez near Sagua la Grande, in the Province of Santa Clara.

Urocoptis (*Gongylostoma*) *bacillaris* n. sp. Plate vi, figs. 7, 8.

Slenderly fusiform, somewhat swollen in the middle third, taper-

ing to a narrow truncation above; thin, pale corneous, profusely variegated with cream-white and having pale brown streaks. Surface glossy, sculptured with rather strong and oblique sinuous riblets, separated by wide intervals; suture subcrenulate. Whorls 12–15, somewhat convex, the last free in a moderately long descending curved neck. Aperture subcircular, a little oblique, the peristome white, sometimes brownish below, expanded and somewhat reflexed throughout. Axis encircled by a thread-like lamella, spinose in the median and upper whorls, and a low spiral cord above it; in the last two whorls the axis is simple and sinuous.

Long. 17, diam. 2.8, ap. diam, 2.3, whorls $14\frac{1}{2}$.

Long. 13.8, diam. 2.2, whorls $13\frac{1}{2}$.

Central Cuba: On stones at the Sierra de Matahambre y Tatibonico, and Veredas de Aguada y del Chorreron, in the mountains near Las Llanadas, district of Mayajigua, in the Province of Santa Clara.

Related to *U. contentiosa* Ar., but differs by its sinuous and more widely spaced riblets and by the free and descending last whorl.

Urocoptis bacillaris exilis n. var. Plate vi, fig. 10.

Smaller than typical *bacillaris;* slenderly fusiform, often entire, the upper two-thirds conspicuously attenuate to the bulbous apex, the lower part somewhat swollen; paler, variegated colored, sculptured with oblique and sinuous thread-like riblets, suture subcrenulated. Whorls 22–24 in the entire shell, 12–13 in the truncated specimens, a little convex. Axis twisted and encircled by a delicate spiral thread, spinose above.

Long. 14.7, diam. 2.1.

Central Cuba: On stones at Vereda le los Negros, a road between La Legua and Tatibonico, district of Mayajigua in the province of Santa Clara.

NOTE ON THE GENUS SEPTA PERRY (TRITON AUCT.).

BY WM. H. DALL.

A recent article in the Victorian Naturalist (Australia) by Messrs. Mathews and Iredale, on the hitherto unnoticed *Arcana* of George Perry, shows that another change is necessary in the nomenclature of this genus. It appeared that although Perry states in his "Con-

chology" that several of his genera are "for the first time published" in that work (the text of which probably was written first), yet some of them actually appeared in print a year earlier in his short-lived journal called the "Arcana." Among these was *Septa*, which appeared in connection with the single species *Murex rubecula* Linné, which of course must be taken as the type. Perry's idea of his genus doubtless was that it covered the species called *Triton* by Lamarck, and so, in intent, it is a synonym of *Cymatium* Bolten, 1798. On account of the character of the type however, it might be retained as a section for the species I formerly ranked under *Lampusia* Schumacher, 1817.

The first name available for the group typified by *Murex tritonis* L. seems to be *Nyctilochus* of Gistel, 1848, and if, in accordance with usage, we take this as the typical genus of the family, the family name would be *Nyctilochidæ*. The genus *Aquillus* Montfort, 1810, upon which the family name was based by Dr. Pilsbry some years ago, is unfortunately synonymous with *Cabestana* Bolten, 1798, and therefore cannot be used.

NOTES.

Polygyra palliata Say in New England.—The occurrence of this species in New England has for a long time needed verification. Repeated search in the Berkshires and Green Mountains by the writer has failed to reveal its presence, although the other species with similar distribution have been found. C. B. Adams, in Thompson's History of Vermont, p. 159, 1842, says: "It is seldom found in Vermont." W. G. Binney includes the species in the "Invertebrata of Massachusetts," but gives no locality. In July Messrs. G. M. Allen and W. S. Brooks while trapping for small mammals on Mt. Ascutney, Vt., obtained three specimens, one adult and two young; the latter were found feeding on the wet oat-meal used in baiting the traps. It is somewhat surprising that this species should turn up at last in this more isolated peak in the Connecticut Valley and not in the Green Mts. proper, although there is little doubt but that it will be found there with more thorough collecting.—C. W. Johnson.

The University of Michigan has conferred the honorary degree of Doctor of Science upon Mr. Bryant Walker, of the class of 1876, in recognition of his valuable work on mollusks.

Snail Trap.—"A woman whose garden was being overrun with snails wrote to the Bureau of Agriculture at Washington for a remedy. She was told to put large pieces of cut raw potato on the under side of a board, raised at one end a few inches from the ground. This she did, with the result that in one week a thousand snails had collected in that one spot."

PUBLICATIONS RECEIVED.

New Species of Land-shells from the Panama Canal Zone.—By William H. Dall. Smiths. Misc. Coll. vol. 59, no. 18, July 27, 1912.

Pleurodonte (*Labyrinthus*) *goldmani* and *Plekocheilus pirriensis* are described as new. The latter is a large and fine form related to *P. coloratus* of Colombia. Not having seen the specimens I may be wrong, but *P. goldmani* does not appear distinct from *sipunculata* Forbes, which I have ranked as a variety of *P. otis*. The specimen figured as *P. labyrinthus* Desh. seems to be what I described as *P. otis orthorhinus*. It is not *labyrinthus* Dh., which is the same as *P. otis* Sol. H. A. P.

The American Species of Sphyradium, with an inquiry as to their generic relationships. By G. Dallas Hanna. U. S. Nat. Mus. Proc. vol. 41, pp. 371–376. The type species, *S. edentulum* Drap. (*Vertigo simplex* Gld.) has been dissected by Mr. Hanna, who shows that by the direct ureter and genitalia it belongs to the *Pupillidæ*, closely resembling *Vertigo*. The jaw however is composed of 16 overlapping plates, as in *Punctum*. The classification proposed by Dr. Sterki and generally adopted recently will therefore give way to the old view that *edentulum* is a Pupoid snail. Until *Punctum* can be examined, its position must be held in suspense. If the foot has pedal grooves, it will remain in *Endodontidæ*. Otherwise it may be a helicoid form of *Pupillidæ*, but this does not seem likely. Mr. Hanna gives *S. alticola* Ingersoll specific rank, and describes a new form from the Pleistocene of Kansas as *S. hasta*. It is nearly 6 mm. long.

It may be mentioned in this connection that the name *Sphyradium* is not tenable for this group, which must be called *Columella*. The argument in the case will be given in a monograph of New York shells now in preparation.—H. A. Pilsbry.

The Nautilus.

Vol. XXVI. OCTOBER, 1912. No. 6

THE MUSSEL SHELL BEDS NEED PROTECTION.

BY WILLIAM T. BARRET.[1]

I began in the mussel shell business in the season of 1902 on the Ohio river at Henderson. Ky.; have been in it ever since that time, and have done practically all of my work on the Ohio. The first five years I never worked above Louisville, finding all the shells that we could sell in that territory, that is, from Louisville down. The beds were large and very productive. I have known from 50 to 60 boats to work on one bed, and some beds produced from 3000 to 5000 tons of shells before they were whipped out. What I mean by being "whipped out" is, that there are not enough shells left to pay wages for catching them. A mussel catcher could work faithfully for 10 hours and not catch over a box of shells on these beds. The price for catching is from 25 to 40 cents per box. The price depends on the quality of shells caught. These boxes will hold about 100 pounds of cooked shell.

In the early days the mussel catchers caught from 10 to 15 boxes per day on these beds. As this territory became less productive, it was necessary for me to find new territory where shells were more plentiful. I came to the upper Ohio, and with camps scattered along the river, have worked it from Louisville, Ky., to Parkersburg, W. Va., the last four years. There may be some wild beds left, but there are very few, if any.

[1] From a memorial addressed to the General Assembly of Kentucky.

The Ohio has been worked hard from one end to the other, and if the present system of working the beds continues, in two or three years, they will all be "whipped out." Under the present system you can work where you like, so naturally everybody likes to work where the mussels are the thickest, and the catchers keep right after them until they whip the bed out.

I remember when the bed at Shawneetown, Ill., was found, and it was one of the best ever found on the river. The mussel catchers working on the Wabash river in Indiana heard what a fine bed it was, and came out of the Wabash, floated down the Ohio to Shawneetown until there were over 100 boats at work on that bed. They worked it out in a few weeks. Since that time it has been worked some each year, but has produced very little. It was found seven years ago and produced that season about 2000 tons of shells, and the total production since that time will not exceed 200 tons in the seven years.

Over 25 per cent. of the mussel shells caught in the Ohio river today are practically wasted—they are so small that the cutters in the blank factories, who work by the piece, can't make anything cutting them, so naturally they throw them in the waste shell if the foreman of the shop isn't standing right by looking. In most shops they work as many as 100 cutters, and one foreman; so you see he can't see many.

My idea is that the State, through its Legislature, should take control and regulate the number of boats per mile that work these beds. I think that four boats to the mile would be about right and only allow them to catch 15 tons to the boat. In other words, confine the production to 60 tons to the mile of shell beds per year.

There are about 90 miles of shell beds in the Ohio river in Kentucky's boundary from Cattletsburg to Cairo, and equally as many in her tributaries. This would mean a production of from 5,000 to 10,000 tons per year. This production would save the mussel beds, and they would continuously consume the sewerage, thereby purifying the water, and would insure from 5,000 to 10,000 tons of raw product annually for the button factories of this country.

The Legislature could pass a bill granting the State the right to protect the beds by license for a term of years, say on a royalty basis of $1.00 per ton for all workable shells produced. This would give the state a revenue from this source of from $5,000 to $10,000 per

year, and it would be handled without any expense to the state. The State Auditor should be given the power to make a contract along the lines mentioned for their protection. The contracting party could be required to send an invoice of each shipment monthly, together with check for royalty to the Auditor, thereby saving the State any expense. The books to be always open for inspection, and a bond given the State that will guarantee this contract to be carried out to the letter.

The Legislature in making a law giving the State this right should embody in it all the things necessary to protect the mussels and mussel beds, viz.: That 60 tons of button shells to one mile of mussel bed is all that can be taken out in one year; this to apply to beds of less than a mile proportionately, one-quarter of mile, 15 tons; one-half mile, 30 tons, etc. No dredges of any kind, or oyster tongs should be allowed—nothing but the hook that catches the mussel as an individual. And the hook should not be allowed to work on any mussel beds when there is more than 20 feet of water on the nearest government gauge. When there is more water than this, the current is much stronger and drags the hooks over the beds so fast that they often catch into the mussel itself, thereby killing it and losing the shell entirely. I have seen boats working in high water that would pull up as many as 25 of these mussel meats at one time. This is ruinous to a mussel bed.

The catchers like to work on high water because there is more food in the water for the mussel at that time, and consequently more mussels come to the surface to feed, and they can catch them faster. There should be a fine of not less than $25.00 nor more than $50.00 and 30 days in jail for each offense for any one violating any of these laws. Each county should have legal jurisdiction to prosecute.

First Step Toward Protection.—Iowa is the oldest state in experience with the mussel-shell button industry. The first factory was started at Muscatine about 16 or 18 years ago. They got the shells from the upper Mississippi river, and for several years found sufficient quantity there to run their plants. As the business increased, it became necessary for them to hunt new fields. They have continued to do this until to-day the Mississippi river and all its tributaries have been worked and worked without any idea of protecting the mussel beds.

The button industry has made tremendous strides—9 years ago

consumption of shells by the button factories was from 10 to 12 thousand tons—to-day the consumption of a normal year will exceed 100 thousand tons. The State of Iowa has seen how necessary it was to do something to insure the raw product for the button factories. To this end through their Congressmen they got the Federal Congress at Washington to appropriate money for the protection of the mussel. With this money they have established at Fairport, Iowa, a mussel hatchery, where they are experimenting and trying to raise the mussel in ponds. They realize the necessity of producing the shells.

Instead of working the mussel beds until we are forced to hatch, cultivate and raise them, why not protect the wild beds that we have, and harvest the crop that they produce annually?

There is no doubt but this is the better way; and Kentucky has in its boundary the major portion of the Ohio river with 90 miles of shell beds. Protect them and save this raw product for the button industry. It is valuable to Kentucky and to the whole world. This is the right way—take the lead, and the other states will follow.

FRESH-WATER MOLLUSCA LIVING OUT OF THE WATER.

BY V. STERKI.

Aplexa hypnorum (L.) and *Sphærium occidentale* Pme. are well known examples of mollusks living at places where water stands only during part of the year, often for short periods in spring, and occasionally after heavy rains. The last named species appears to be especially well adapted to that kind of habitat, and even recently discharged young mussels have often been found living on apparently quite dry soil under a thin layer of dead leaves. It is also known that small *Lymnæa*, and *Pomatiopsis* are ofter found crawling out of water.

Fresh-water mollusks of almost all groups bury themselves in the soil, with the disappearance of surface water, and survive for shorter or longer periods. But quite a number of pulmonates and branchiates appear to remain not only alive but active and propagating for long periods or permanently. How much this fact has to do with

the evolution of terrestrial life from aquatic, and what physiological, and eventually anatomical and embryonic changes are involved, must be left to the scientists.

The following observations may be worthy of notice in this connexion.—In low woods, north of Geneva, Ohio, *Segmentina armigera* (Say) were found alive in company with *Sphær. occidentale*, and in the same woods *Physa gyrina* were found, then mostly dead shells, where they could not have been transported from any permanent body of water.

In damp woods at Kenmore, Summit Co., Ohio, among patches of various terrestrial mosses, a few dozen *Amnicola limosa* Say (*A. parva?*), living and of various ages, were found (late August), and with them a few puerile *Physa* apparently *sayi* Tappan. No water was standing in the woods, even after heavy rains, and moreover these snails or their parents must have survived e. g. the exceptionally hot and dry summer of 1911.

In July of that summer, in a marsh near Hudson, Ohio, which had been dry as a bone for weeks, like hundreds of others, Dr. Rush and I found several dozen specimens of *Lymnæa* apparently *lanceata* Gld., living and in good condition, clinging to stems and leaves of sedges (*Carex* and *Scirpus*), several inches above the ground. There is hardly a doubt that they had been gathering all the moisture they could from dew at night.

In woods of that vicinity we found *Pisidium abidtum subrotundum*, typical, full-grown to young, under dead leaves in a damp spot. Other *Pisidia* have been found at similar places, and also in swamps among mosses, out of water, and even on steep banks where water could never be standing.

Small *Lymnæa*, apparently *parva* Lea, are often found in greenhouses, on mossy flowerpots, sometimes in large numbers, evidently doing well and propagating.

Similar observations have doubtless been made by other malacologists.

NEW CUBAN UROCOPTIS OF THE U. CINEREA GROUP.

BY DOCTOR CARLOS DE LA TORRE.

(*Concluded from page 58*).

Urocoptis bacillaris albicostulata n. subsp. Plate vi, fig. 9.

This species differs from the type and its other varieties in the sculpture of stronger white thread-like riblets.

Long. 18.2, diam. 2.9, whorls 17.

Long. 12.9, diam. 2.1, whorls 13.

Central Cuba: On stones at Cueva Bonita and Rosa Perdida in the northeastern limit of the Santa Clara Province.

Urocoptis bacillaris sordidula n. subsp. Plate vi, fig. 11.

This is the smallest of the *bacillaris* race. It is slenderly fusiform and truncate above, corneous and whitish streaked but the surface is lusterless; sculptured with less oblique and sinuous thread-like riblets separated by less wide intervals. Whorls 10–41 a little convex, the last free in a shorter neck, with the aperture more oblique. Axis twisted and encircled by an inconspicuous spiral thread, minutely spinose above.

Long. 12, diam. 2.1, ap. diam. 2 mm., whorls 12.

Central Cuba: On stones along the road east of El Bano de Mayajigua in the Province of Santa Clara.

This may be regarded as a degenerate form of *U. bacillaris* living in the plain near the foot of the mountains where the typical form lives.

U. (*Gongylostoma*) *stricta* n. sp. Plate vi, figs. 16, 17.

Differs from *U. bacillaris* in its more slender shell which in this species is not swollen and is almost cylindric in the middle third; also differs by the sinuous, well-raised riblets, more regularly wide-spaced throughout which are somewhat tubercular at their upper ends. Whorls slightly convex. Axis encircled by a thread-like spinose lamella which in some of the whorls has an indistinct spiral cord above it.

Long. 19, diam. 2.2, ap. diag. 2, whorls $16\frac{1}{2}$.

Long. 13.7, diag. 2, whorls $12\frac{1}{2}$.

Central Cuba: On stones at Lomas del Purio near Calabazar de Sagua in the Province of Santa Clara.

This species resembles *U. wrighti* Pfr. though it is less slender and more strongly sculptured.

Urocoptis (*Gongylostoma*) *boqueronis*, n. sp. Plate vi, figs, 1, 2, 3.

Slenderly fusiform, tapering to a narrow truncation above; thin, corneous with some indistinct darker streaks; regularly sculptured with narrow thread-like whitish riblets which are slightly arcuate, somewhat oblique in the upper half, less so in the lower, and separated by intervals two or three times their width. Whorls 11–13, a little convex, the last free and produced into a rather short neck. Aperture subcircular, a little oblique, the peristome whitish, expanded and somewhat reflexed. Axis encircled by one spiral spinose lamella with a low spiral cord above it.

Longitude 13.2, diameter 2.4, aperture diameter 2, whorls 13½.

Central Cuba: On walls exposed to the sun at the entrance to El Boqueron, an underground passage of the river Tatibonico del Norte through the Sierra de Tatibonico, in the eastern limit of the Santa Clara Province.

This species differs from *U. bacillaris* and its varieties by its non-sinuous but slightly arcuate riblets, more uniform color, less produced neck, etc.

Urocoptis (*Gongylostoma*) *parallela*, n. sp. Plate vi, figs. 4, 5.

Slenderly fusiform, frequently entire, somewhat swollen in the middle, tapering to a narrow truncation or to the bulbous apex: pale brown in color; sculptured with strong wide-spaced riblets which are nearly straight; whorls 26–28 (entire), 12–14 in the truncate specimens, somewhat convex, the last one free and produced in a long curved neck. Aperture subcircular, slightly or not at all oblique, peristome whitish, expanded and reflexed as in the other related species. Axis encircled by one spiral spinose lamella and a well developed spiral cord above it.

Longitude 20, diameter 2.9, aperture diameter 2.5, whorls 26, (entire).

Longitude 15.2, diameter 2.9, aperture diameter 2.6, whorls 13½, (truncated).

Central Cuba: On walls exposed to the sun at the north side of

El Boqueron where reappears from underground the river Tatibonico de Norte, in the eastern limits of the Santa Clara Province.

This species resembles *U. boqueronis* but is easily distinguished from it by the larger size, uniform color, the straighter riblets which are also stronger and wider spaced.

Urocoptis acicularis, n. sp. Plate vi, fig. 6.

Cylindric-subulate, very slender, the upper third tapering to a very narrow truncation or to a bulbous apex; brownish with scattered dark corneous streaks bordered with white patches. Sculpture, delicate thread-like and wide-spaced riblets which are nearly straight and slightly or not at all oblique. Whorls 25–28 in the entire shell, 14–16 in the truncate specimens, nearly flat, the last free in a curved neck with a weak basal keel. Aperture subcircular, brownish inside, the peristome white, glossy, expanded and flatly reflexed. Axis encircled by one spiral, denticulate or spinose, well-developed lamella without a noticeable cord above it.

Longitude 17, diameter 2, aperture diameter 2, whorls 27, (entire).

Longitude 14, diameter 2.1, whorls 15½, (entire).

Longitude 15.2, diameter 2.3, whorls 15½, (entire).

Central Cuba: On stones and walls at Chinchilla and the Mogotes de las Tumaguas, near Sagua la Grande in the Province of Santa Clara.

Characterized and distinguished from the others of the group chiefly by its cylindric form and very slender shell.

Urocoptis (*Gongylostoma*) *camagueyana* n. sp. Pl. vi, figs. 12, 13.

Fusiform-cylindric, more swollen above the middle and slowly tapering to a wide truncation; cream-colored or pale corneous, a little darker below. Sculptured with very strong cord-like whitish riblets which are in part hollow but rarely broken, more inflated and closely spaced in the upper than in the lower whorls, where they become degenerate, though again becoming somewhat lamellose on the neck. The suture is denticulate by the projection of the ends of the riblets. Whorls 7–9, somewhat convex, the last one free and produced in a long, round, twisted neck. Aperture subcircular, slightly oblique, the peristome whitish, expanded and a little re-

flexed. Axis encircled by an inconspicuous spiral cord which is somewhat lamellose and minutely spinose above.

Long. 11.3, diam. 2.5, ap. diam. 2, whorls 8.

Long. 14.5, diam. 2.8, whorls 13.

Central Cuba: On stones at El Cercado, Cerro de Tuabaguey, and Los Paredones, a narrow pass across the Sierra de Cubitas, in the northern part of the Province of Camaguey.

This little species, notable for its wide truncation, strong cord-like riblets and the long contorted neck, belongs to the same group as the more elongated preceding species of the neighboring region of Santa Clara province, but it is also geographically and morphologically one of the most distant forms of this section. Notwithstanding, in the young specimens, when the shell is complete, it has the shape of *U. boqueronis* and *U. parallela*, its singular form being due to the mutilation suffered in the adult stage; the aperture and peristome are similar; the internal axis, though different at first sight, because it is encircled by a low cord, if attentively examined in the first whorls, appears to be somewhat lamellose, and bearing very minute characteristic spines at the edge; and finally the radula is of the same type as in *U. bacillaris*, etc., similar to that of *U. baculum* Pilsbry, and to Pilsbry's section *Tetentrodon:* 15 to 19 teeth; but in *U. camagueyana* the two last lateral teeth are so degenerated that only their places remain on the radula, so that the formula could be (2)5.2.1.2.5(2).

NOTES.

Pleistocene Fossils from Eastern Cuba.—Mr. Clarence L. Aman at my request investigated a deposit of clay exposed in an excavation at Guamo, on the Rio Cauto, Oriente Province, Cuba. He sent the following species to the Academy of National Sciences: *Marginella minuta* Pfr., *Bittium gibberulum* Ad., *Cæcum* (*Meioceras*) *nitidum* Stimp., *Leda acuta* Conr., *Polymita picta* Born, *Cepolis strobilus* Fér., *Cepolis lucipeta* Poey and *Pleurodonte emarginata* Gdl. This find is interesting as it shows that in Pleistocene times the sea extended far up the Cauto valley.—E. G. Vanatta.

Ashmunella thomsoniana.—I recently collected a dead shell of this species in a new locality, the steeper part of the trail between

the Field and Viveash ranches in the cañon of the Pecos River, New Mexico. The specimen is to be referred to *thomsoniana* proper, having the basal tooth simple, but it is the smallest on record for that species; diam. max. $11\frac{1}{3}$, min. a scant 10, alt. $6\frac{1}{3}$; whorls $4\frac{1}{3}$. Should this small size and number of whorls prove constant in the locality, the snail would I think deserve to be separated as a race.—T. D. A. Cockerell.

The Genera Cryptogirasia and Cryptosoma. In the Fauna of British India, Mollusca (vol. I, 1908), p. 203, Godwin-Austen adopts the generic name *Cryptogirasia*, crediting it to "Cockerell, A. M. N. H. (6) vii, 1891, p. 99 (no description)." The name *Cryptogirasia* was a manuscript one of mine, dating from 1891; but as a matter of fact it was printed in the plural form (*Cryptogirasiæ*) in the place cited. It was not wholly without description; this and the other genera and groups of "*Helicarioninæ*, sens. lat." were defined by a table. From the standpoint of generic nomenclature, *Cryptogirasia* must date from 1908, when it was adopted and defined by Godwin-Austen. The other genera credited to me in the same work were published in the correct form, either in P. Z. S. or Nautilus. They were defined in the table cited. *Cryptosoma* Theobald, 1857, applied to an oriental genus of the same family, is a homonym of the earlier *Cryptosoma* Brullé, used for a genus of crabs of the family Calappidæ. I have not found the exact date of Brullé's genus, but it is not later than 1848. The Molluscan *Cryptosoma* may take the name *Megaustenia*; type *Megaustenia præstans* (*Vitrina præstans* Gould, 1843).

T. A. Cockerell.

Unio (Nodularia) Putzeysi new name for *U.* (*Nodularia*) *subnigra* Preston.—Mr. Bryant Walker having pointed out to me that the name "*subnigra*" used by me (Ann. Mag. Nat. Hist., London, Ser. 9, vol. IV, 1909, p. 89) for a species of *Unio* from the Lower Belgian Congo is preoccupied, I now propose to substitute for this the name "*Putzeysi*" in honour of Dr. S. Putzeys of Brussels, whose work on the land and fresh-water molluscan fauna of the Belgian Congo is well known to every student of the conchology of that region.—H. B. Preston.

PUBLICATIONS RECEIVED.

THE MALACOLOGICAL SOCIETY OF LONDON has given variety to the last number of the *Proceedings* by adding a department of humorous fiction. In it they print a fantastic farce entitled "A synopsis of the recent and tertiary fresh-water mollusca of the Californian Province." A leading part in this piece is taken by a weird character called "Syntonia," or the "Syntonic Form"—a sort of djinn or genie, competent to produce any transformation desired with the aid of a pinch of magnesium salts, which the author obligingly supplies when it cannot be found otherwise.

That magnesium salts are "producing the mischief" with the snails is proved by an act with "balanced aquaria,"—conditions, results and laboratory where done not specified.

What the Syntonic Form cannot do is readily accomplished by an "accentuation of the evolutionary influences" which, along with other deviltry, are to blame for making *Lymnæa auricularia* out of the "so-called *peregra* and *catascopium*." Curious nobody ever thought of this perfectly lucid explanation before.

But what occult influence is responsible for subgenera based on forms ranked as subspecies of species belonging to other subgenera? Or putting a genus in one subfamily, its subgenera in another (done several times on p. 147)? The method of classification is blamed on Professor Hyatt, but we all know that Hyatt was a good naturalist. He was above all a believer in strictly phylogenetic classification. He had little use for such elementary trifling in taxonomy as grouping together genera of different phyla which have reached superficially similar or analogous evolutionary stages.

Time wasted in looking up type specimens might be saved if naturalists would get syntonic brains by judicious dosing with magnesium salts. Just see what has been done to the freshwater Mollusca of the Californian Province! Synonymy can be made by glancing over a few books; classification by drawing the names out of a hat after they have been well shaken up. The study of variation and inheritance may be be left to old fogies who want to know what really happens under known conditions.

We fear that the Malacological Society has started out too well in its department of humorous conchology. Merely ignorant or wrong-headed articles they may possible get, but where such another contribution to the gayety of nations as this?—H. A. P.

The Mollusca of Colorado, Part III. By Junius Henderson (Univ. of Colo. Studies, vol. IX, May, 1912). Contains numerous additional locality records (of which fourteen pertain to species and subspecies not included in former parts of the work), some rectifications, and evidence for the removal of a few species given in former lists. A faunal work is never finished, but the papers by Mr. Henderson form a good foundation for further work in Colorado. It is by far the most complete account of any Rocky Mountain mollusk fauna.

The West American Mollusks of the genus Eumeta.

The recent and fossil mollusks of the genus Cerithiopsis from the west coast of America.

The recent and fossil mollusks of the genus Bittium of the west coast of America.

New mollusks of the genus Aclis from the North Atlantic.

The west American mollusks of the genus Amphithalamus.

The recent and fossil mollusks of the genus Alvania from the west coast of America.

The west American mollusks of the genus Cingula.

These papers by Dr. Paul Bartsch published in the Proceedings of the National Museum for 1911–12 are well-illustrated monographs of the west American species of the genera named. All are small or minute shells, the identification of which has hitherto been of great difficulty owing to the brevity of the scattered literature concerning them, and the lack of illustrations. Dr. Bartsch has described the old and new species with a fulness of detail which all who work with these forms hereafter will appreciate.—H. A. P.

Report on Landshells collected in Peru in 1911 by the Yale Expedition under Professor Hiram Bingham. By William Healey Dall (Smithsonian Misc. Coll. vol. 59, No. 14, 1912). Extensions of the known range of a considerable number of species are given, but the chief interest of the paper attaches to the discovery that a group of slender Andean Bulimuli including *scalaricosta*, *infundibulum*, *umbilicatellus*, *tubulosus* and *spiculatus* have an internal lamella on the axis, somewhat as in *Holospira*. The new subgenus *Phenacotaxus* (type *umbilicatellus* Pils.) and the section *Ataxiellus* (type *pectinatus* Dall) are proposed to receive these forms. The color-variation of *Drymæus poecilus* Orb. is discussed.—H. A. P.

D. D. BALDWIN.

The Nautilus.

Vol. XXVI. NOVEMBER, 1912. No. 7

NOTES ON MAINE MOLLUSCA.

BY REV. HENRY W. WINKLEY.

For the purpose of making a more careful examination of the warm-water colonies in Maine, the writer spent two weeks at Wiscasset and the same time at Orr's Island in Casco Bay. General collecting at Wiscasset revealed the same species that I found on former visits. A careful search in sheltered places revealed several colonies of *Odostomia trifida* and *bisuturalis*, especially large and abundant at Edgecomb. A drive of five miles and work in a drenching rain at Sheepscote Bridge gave one or two specimens of *Odostomia bisuturalis* var. *ovilensis*, *Modiolus demissus* var. *plicatulus* and *Ostrea virginiana*. Unfortunately the refuse from a saw-mill has nearly exterminated life at the type locality for *ovilensis*. Ten miles east of Wiscasset, at Newcastle and Damariscatta, I had only one low tide, but I found *Modiolus demissus* var. *plicatulus*, *Crepidula convexa*, *Ilyanassa obsoleta*, *Haminea solitaria*, *Urosalpinx cinereus*, *Astyris lunata* and *P.* (*Syrnola*) *fusca*, the last two new to Maine, I believe. *Odostomias* have been found here by Mr. Wentworth, but I didn't have time to visit the locality where he found them.

At Orr's Island I found two more colonies of *Astyris lunata*, a few *Odostomia bisuturalis;* other forms of interest were *Cingula aculeus*, *Skenea planorbis* of a light color, *Retusa pertenuis* and *gouldii*, *Bela bicarinata* var. *violacea* (one specimen at low tide) and *Mysella planulata*, with of course the more common forms. I found only one land shell, *Zonitoides arborea*.

A day was spent in the famous Quohog Bay. I dredged the whole length of the bay and found it absolutely barren, except a very interesting colony at the head of the bay. The species found there were: *Ensis directus*, *Callocardia morrhuana*, *Macoma calcaria*, *M. balthica*, *Yoldia limatula* (the largest I have ever seen), *Venus mercenaria*, *Mysella planulata*, *Gemma gemma*, *Tellina tenera*, *Retusa pertenuis*, *Crepidula convexa*, *Haminea solitaria*, *Astyris lunata*, *Odostomia trifida*, *bisuturalis*, *winkleyi* and *gibbosa*.

A trip to the biological laboratory at South Harpswell enabled me to rob the cemetery in the next lot, result equal: *Vitrina limpida*, *Cochlicopa lubrica*, *Pupilla muscorum*, *Vallonia excentrica* and *Pyramidella cronkhitei* var. *anthonyi*.

The famous Brown Cow Island was visited. *Helix hortensis* banded and plain was there, but not so abundant as sixteen years ago. I also obtained a few of the beautiful wine-colored *Polygyra albolabris*, together with *Succinea ovalis*, *Pyramidula alternata*, *Vitrea hammonis*, *Vitrina limpida*, *Pyramidula cronkhitei anthonyi*, *Helicodiscus parallelus* and *Cochlicopa lubrica*.

On the whole a very satisfactory summer, with several species added to the marine list.

ON A COLLECTION OF MARINE GASTROPODA FROM ADEN WITH DESCRIPTIONS OF NEW FORMS.

BY MAXWELL SMITH.

(Read before the Brooklyn Conchological Club Jan. 29, 1912).

The collection which forms the subject of this paper is a portion of a series collected by Mrs. Blair, the wife of a Governor of Aden. The greater portion of this interesting collection is housed in the Royal Scottish Museum of Edinburgh. During the summer of 1910 Mr. Eagle Clarke, the Curator of the Department of Mollusca, placed in my hands a number of shells selected from the Blair collection. Upon reaching London I compared many of the species with the types in the British Museum and also examined the collections made by Messrs Yerbury and Baynham which are preserved in the same institution.

A brief résumé of the work that has already been done at Aden

may be of interest. One of the most exhaustive treatises dealing with molluscan life in these waters may be found in the proceedings of the Zoological Society of London, 1891, part iii. In this paper Mr. E. A. Smith lists the collections made at Aden by Major Yerbury and the Rev. Mr. Baynham. Mr. Smith also gives a review of the principal collections previously recorded from Aden. Signor Caramagna, Dr. Jousseaume and the Rev. A. H. Cooke reported on mollusks from the vicinity of Aden. After the appearance of Mr. Smith's paper Commander E. R. Shopland was apparently the next person who collected extensively at Aden. His field work was accomplished between 1892 and 1901. An account of the results may be found in the Proceedings of the Malacological Society of London (Vol. 5 page 171).

My best thanks are due Mr. Eagle Clarke of Edinburgh and Mr. E. A. Smith of London for their kindness in supplying data, without which it would have been impossible to prepare this paper.

Sepia rouxii D'Orb. According to Tryon this cephalopod inhabits the Red Sea and Indian Ocean. It is thus natural to find it in the Gulf of Aden, which connects the two.

Conus amadis Auct.

Conus betulinus L.

Conus cuvieri Crosse. As Mr. E. A. Smith writes, this species is not identical with *C. cervus*. In my opinion *cuvieri* is a distinct species.

Conus nussatella L.

Conus taeniatus Hwass.

Conus textile L.

Terebra caerulescens Lk.

Terebra cinerea Born.

Terebra clarkei n. sp. Plate IV, fig. 18.

Shell slender, shining, suffused irregularly with golden brown; whorls 9–10, somewhat gradually increasing, distinctly rounded; apex acute, darker in color; surface closely plicated, forming on the earlier whorls slightly crenulated sutures, plications extending to the base; distinctly three banded, the two lower bands closer together; aperture long.

This species comes in the same section of the genus as *T. nitida* Hinds. From *Terebra celidonota* Melvill & Sykes it differs in being more slender. The aperture is also longer and there are three spiral

bands in place of the one band, near the suture, which occurs in *celidonota*. In *clarkei* only one of the bands is visible on the upper whorls. I take pleasure in naming this species after Mr. Eagle Clarke. Type in the writer's collection.

Terebra lamarckii Kien. I agree with Mr. Smith upon the validity of this species. The examples before me are well defined and show little variation. Tryon and Reeve considered it a variety of *T. duplicata* L. In the latter the longitudinal striæ are farther apart. The style of coloration is also strikingly peculiar.

Pleurotoma tigrina Lk.

Murex palma-rosæ Lk. Not previously reported from Aden.

Murex ramosus L.

Murex rota Sowb. Fresh beach specimens were obtained of this beautiful species.

Murex tenuispina Lk. This and *M. palm-rosæ* may have been obtained at some distance from Aden.

Columbella picta Reeve. Pl. IV. fig. 13. Up to the present the locality of this species has been unknown. The specimen before me agrees perfectly with the type in the British Museum.

Columbella misera Sowb. var. *californica* Reeve. Plate IV, fig. 15. Dr. Pilsbry, to whom I submitted the single example, writes "this occurs in Japan—not in California, which was an error of Reeve's."

Columbella propinqua E. A. Smith. Plate IV, fig. 14. The type is in the British Museum.

Columbella terpsichore Leathes. Plate IV, fig. 12. Aden, as the first locality known for this species, was reported by Shopland. The type is in the British Museum.

Eburna valentiana Swainson. I am inclined to believe that this is only a stout form of *E. spirata* as pointed out by Von Martens.

Nassa albescens Dunker.

Nassa arcularia L.

Nassa bimaculosa A. Ad. From other examples in my collection I conjecture that *bimaculosa* is a valid species.

Nassa canaliculata Lam.

Nassa coronata Lam.

Nassa fissilabris A. Ad.

Nassa gemmulata Lam.

Nassa nodifera Powis.

Nassa pulla L.

Nassa pulla L., var. *minor*, n. var. Plate IV, fig. 17. The length of this shell is about the same as the aperture of the type. All the examples obtained are of the same size.

Vasum cornigerum Lam.

Ricinula ricinus L.

Rapana bulbosa Solander.

Cancellaria melanostoma Sowb.

Mitra costellaris Lam. Apparently the first record of the species in these waters. Shopland reports 50 species of *Mitra* as occurring at Aden. With the addition of two species taken by Yerbury and Baynham, and the present two species, the total number is brought up to 54.

Mitra cremulata Lam.

Mitra deshayesii Reeve. This species may be said to inhabit the triangle formed at the corners by Polynesia, Mauritius and the Red Sea. It has not before been reported from the Gulf of Aden.

Mitra plicata Lam.

Latirus polygonus Gmelin.

Harpa minor Rumph.

Harpa ventricosa Lam.

Oliva andamanensis Bdg. Plate IV, fig. 7. The four examples before me agree perfectly with those in the British Museum, which were taken at the Andamen Islands.

Oliva inflata Lam. From the number of individuals taken, this must be an abundant form.

Ancillaria acuminata Sowb.

Ancillaria albisulcata Sowb. A characteristic feature of this species is the long white columella.

Ancillaria castanea Sowb. I am inclined to believe that this species will prove to be a variety of *A. fulva*. The color is the only distinction.

Ancillaria fulva Swains. The columella of this species is considerably thickened at the base.

Marginella obscura Reeve. Plate IV, fig. 16. The locality of this species was for many years unknown. In 1891 Mr. E. A. Smith discovered the species among the Yerbury and Baynham Collections and included it in his list (P. Z. S. L.). When in London I examined these shells and also the type. The Blair shells are slightly larger, but otherwise similar to the type.

Cassis nodulosa Gmelin. The example before me is remarkably fine. There are 12 tubercles on the margin.

Lotorium pilearis L.

Natica lineata Lam.

Natica chinensis Lam.

Naticina papilla Gmel.

Cypræa clandestina L.

Cypræa cruenta coloba Melvill.

Cypræa erosa nebrites Melvill. Plate IV, fig. 9.

Cypræa erosa subalba, n. var. Plate IV, fig. 11.

This beautiful new variety of *erosa* comes nearest to *nebrites*. It differs from the latter in possessing milk white "specks" on a violet white ground. The ocellations are not so persistent as in the var. nebrites. The brown square spots, however, are similar. Type in the writer's collection.

Cypræa erosa carmen n. var. Pl. IV, fig. 10. Sides much swollen, pitted and furrowed above the callus on the left side. The base is white, having none of the stippling of *nebrites*, and the squarish spot at the side is nearly or quite covered by the basal callus. The back is pale lilac-pink with traces of white spotting at the sides only. Two cotypes measure 32 and 24 mm. long, in coll. A. N. S. P. (No. 106457) and my own.

Cypræa errones L.

Cypræa felina ursellus Gmel. This variety may easily be distinguished by the white base.

Cypræa helvola L. Pl. IV, fig. 8.

Cypræa helvola L. var. mascarena Melvill. Plate IV, fig. 6.

Cypræa isabella, L.

Cypræa pantherina Sol. Among the series are the vars. *theriaca* (one nearly black), *albonitens* and *syringa*.

Cypræa staphylæa L. This species has apprently not been reported from Aden.

Cypræa stolida L. This new record extends the western range beyond Ceylon.

Cypræa talpa L.

Cypræa tigris L. One individual is $4\frac{1}{2}$ inches long.

Cypræa turdus Lk. The Aden shells are large. I have in my collection three similar specimens which where taken at Muscat, Arabia. This species reaches its greatest development in the Per-

sian Gulf, where the shells are of the same size but greater in weight. The dorsal surface of the Persian Gulf examples I find white and the coloring also somewhat lighter.

Turritella columnaris Kien.

Mitrularia equestris L. var. *tortillis* Reeve.

Strombus columba Lam. Tryon when writing the Manual was uncertain as to the presence of this species in the Red Sea. Shopland's report and the present record confirm the presence of the species in these waters.

Strombus gibberulus L.

Strombus mauritianus Lam. Tryon and Paetal cite Mauritius as the home of this Strombus. I doubt if it was collected so far north.

Strombus variabilis Swains.

Rostellaria curvirostra Lam. A young specimen is of interest. There are 14 whorls present. The embryonic whorl is pearly, the following $10\frac{1}{2}$ whorls covered with the longitudinal plications, the remaining whorls smooth except the lower portion of the body whorl which is ornamented with spiral lines. The opercula of an adult is twisted and the under surface raised in the center.

Bullaria ampulla L.

Hydatina physis L.

Umbraculum indicum Lam.

NOTE ON A VARIETY OF POLYGYRA FALLAX.

BY H. A. PILSBRY.

The group *Triodopsis* in States of the Atlantic slope presents numerous forms which by their variations perplex the conchologist. *Polygyra* (*Triodopsis*) *hopetonensis* (Shuttlw.) in its various varieties is a widely-spread species. The variety *obsoleta* from Newbern, N. C., probably belongs to it, but it needs further investigation with more specimens.

Polygyra vannostrandi (Bld.) is distinct by its closely coiled whorls, but the aperture is like that of *P. fallax*, both having large teeth, the basal tooth inclined to be double and the outer lip-tooth wide and deep-seated. The two species are not always easy to tell

apart, but are doubtless distinct. *P. v. alabamensis* and another variety, with decidedly wider umbilicus, but still unnamed, replace typical *vannostrandi* in Alabama. Neither is so distinct as to be startling.

Polygyra fallax is common in southeastern Pennsylvania, and extends south to the Carolinas at least. Specimens taken by Messrs. Henderson, Walker, Clapp and myself at Smith's Island, Cape Fear, are much smaller than Pennsylvanian shells. In the western part of the Ozark region the closely related and still smaller *P. cragini* (Call) occurs, being widely separated from all parts of the range of *P. fallax*.

When collecting fossils in Florida in 1900, Mr. C. W. Johnson found a peculiar variety which seems referable to *fallax*, and which I think is worthy of a name.

Polygyra fallax goniosoma n. subsp. The shell differs from *P. fallax* by being distinctly or strongly angular in front, the angle being situated high on the whorl. Sculpture of rib-striæ stronger above the angle, the striæ becoming smaller below it, and usually more numerous by intercalation of striæ. Aperture as in *fallax*, the outer lip retreating, outer lip-tooth broad and deeply placed, basal tooth buttressed on the columella side, parietal tooth angularly bent.

Alt. 7, diam. 12.4 mm.; whorls $5\frac{1}{2}$.

Alt. 6.2, diam. 10.3 mm.; whorls $5\frac{1}{3}$.

Blountstown, Calhoun Co., Florida, under oak logs in oak and pine woods, collected by C. W. Johnson, 1900. Cotypes no. 77948 A. N. S. P.

This form may possibly be referable to *P. vannostrandi*, but in the number of whorls it agrees better with *P. fallax*.

MOLLUSCA FROM NORTHERN NEW MEXICO.

BY JUNIUS HENDERSON.

In 1910, while engaged in ethno-zoological work for the School of American Archæology, at its summer camp in the canyon of El Rito de los Frijoles, about thirty-five miles northwest of Santa Fe, New Mexico, and also in the near-by Jemez Mountains, near Valle Grande, I collected the following species of mollusks:

Ashmunella ashmuni Dall. Abundant at Rito. A few immature specimens from the mountain locality may possibly be the variety *robusta* Pils. The Rito is not a great distance from Bland, the type locality of *ashmuni*, and *robusta* was described as from "Jemez Mountains near Bland, New Mexico, at higher elevations than *ashmuni*."

Oreohelix strigosa depressa Ckll. Three dead specimens at the mountain locality.

Pupilla muscorum L. One at the Rito.

Pupilla blandi Morse. Abundant at both the Rito and the mountain locality.

Bifidaria pellucida parvidens Sterki. One at the mountain locality. Identified by Prof. Cockerell.

(*Vertigo coloradensis basidens* P. & V. Though the type locality is Bland, N. M., none were found at the Rito.)

Cochlicopa lubrica Müller. Common at the Rito.

Vallonia cyclophorella Ancey. Abundant at the Rito and common at the mountain locality.

Vitrina alaskana Dall. Three dead ones at the mountain locality and four at the Rito.

Euconulus trochiformis alaskensis Pils. Common at the Rito and one at the mountain locality.

Zonitoides arborea Say. Abundant at the Rito and common at the mountain locality.

Pyramidula shimekii cockerelli Pils. Only two at the summer camp at the Rito, but abundant two miles above.

Succinea avara Say. One at the Rito.

With the exception of *Ashmunella* and *Bifidaria*, it will be noticed that this is the fauna common in the mountainous portions of Colorado, in similar situations. It would be interesting to know how far north *Ashmunella* extends, which could be determined by an examination of the Rio Grande Valley and adjacent mountains from Buckman northward into Colorado.

University of Colorado, Boulder, Colo.

DAVID DWIGHT BALDWIN.

D. D. Baldwin, the veteran Hawaiian conchologist, died at Honolulu, June 16th, at the age of eighty years and seven months. Burial was in the old Makawao Cemetery, Maui.

Mr. Baldwin was born at Honolulu November 26, 1831. His childhood was spent at Waimea, Hawaii, and his boyhood at Lahaina, Maui. From 1844 till 1851 he attended the Punahou School, where he was prepared for college under the tuition of the Rev. Daniel Dole. In 1852 he sailed for the United States by way of Cape Horn. He entered Yale College in 1853, and graduated there with honor in 1857, taking the astronomical prize. During his college course he became acquainted with Miss Lois Gregory Morris, whom he married October 7, 1857. Returning to the Islands he became principal of the Lahaina School, which he conducted for seven years. Latterly he engaged in the cultivation of sugar cane. From 1865 to 1872 he was manager of the Kohala Planatation, where he introduced the so-called Lahaina cane, to which that plantation owed its success. He then visited New Haven, Conn., where he remained over a year. On his return, in 1874, he was appointed vice-principal of Lahainaluna Seminary.

In 1877 Mr. Baldwin was appointed Inspector-General of Schools, holding that post for eight years. During his administration, and largely by his influence, the number of schools in which the English language was the basis of instruction increased from five to one hundred. He also drew up the first course of study for such schools.

In October, 1885, he retired from the arduous position of Inspector-General to take his former position as vice-principal of Lahainaluna Seminary. In 1891 he removed to Haiku, Maui, and finally retired from educational work in January, 1905, having been connected with the Department of Public Instruction for thirty-eight years.

While Mr. Baldwin's energies were largely devoted to education, he was deeply interested in the natural sciences, especially studying shells, ferns and mosses. In conchology he was esteemed the chief Hawaiian student, and almost the only one to publish original work on *Achatinellidæ* between the period of Newcomb and Gulick and the present time. Baldwin's Catalogue of Land and Fresh-water Shells of the Hawaiian Islands (Honolulu, 1893), though brief and

unpretentious in form, has been most useful to other investigators for its judicious and reliable synonymy and its locality records, the fruit of extensive collecting and inquiry.

Many beautiful new species of *Achatinellidæ* were described by Baldwin in articles published in the Proceedings of the Academy of Natural Sciences of Philadelphia (1895) and in THE NAUTILUS. Among them are several of exceptional interest to those tracing the relationships of the species and genera.

Mr. Baldwin was known extensively among conchologists in America and abroad as a kind and generous correspondent. He supplied many of the new Hawaiian species described by Ancey, Sykes, Pilsbry and others.

His extensive collection and scientific library remain the property of his son Mr. E. D. Baldwin, of Honolulu.

Species of the genera *Amastra*, *Leptachatina*, *Tornatellina*, *Philonesia*, *Endodonta*, *Cecilioides*, *Succinea*, *Helicina* and *Melania* have been named in honor of Mr. Baldwin, and Ancey dedicated to him the group *Baldwinia*, a section of *Partulina*. These, with his good work on *Achatinellidæ*, will keep his memory green.

GEORGE W. TAYLOR, D. D.

The Rev. Dr. George W. Taylor died at his home on Departure Bay, Vancouver Island, August —, ——. Dr. Taylor was well known as a student of Pacific coast shells, especially those of British Columbia. A keen collector in conchology and entomology, he was also well posted in many other branches of natural history.

When a biological marine station was established on this coast, he was put in charge by the Dominion government, remaining in that position until his death. He was largely responsible for the establishment of this station, chose the site on Departure Bay, near Nanaimo, Vancouver Island, and had the building of the station left entirely in his hands.

Dr. Taylor leaves a very large and valuable collection of shells, particularly rich in *Limpets* and *Unionidæ*. In the Transactions of the Royal Society of Canada, 1895–96, he published a Preliminary Catalogue of the Marine Mollusca of the Pacific Coast of Canada, with notes upon their distribution; also a Supplement to the same a

year or two later. He was engaged upon a new Catalogue for publication when I visited him two years ago, which owing to poor health was never completed. Dr. Taylor was a Fellow of the Zoological Society and a Fellow of the Royal Society of Canada. He also published lists of the land and fresh-water shells of Vancouver Island and of British Columbia.—A. W. Hanham.

NOTES.

Land Shells of Carthagena, Colombia.—During a brief stop at this port Dr. A. P. Brown collected the following snails on the hill La Popa, about a mile from Carthagena: *Euglandina cumingi* (Beck), *Oxystyla maracaibensis* (Pfr.) and *Bulimulus krebsianus* Pils. The hill is chiefly composed of recent corals and sea shells. H. A. P.

Viviparus contectoides in Philadelphia.—Mr. J. Baily, Jr., discovered *V. contectoides* Binn. in a lily pond near Memorial Hall, Fairmount Park, Philadelphia, in June, 1909; Mr. S. Jacobs collected some at the same place in 1910, and Mr. H. L. Mather found it again on April 20, 1912, showing that the species is well established there. E. G. Vanatta.

Lymnaea (Radix) auricularia Linn. in Colorado.—Mr. F. R. Warren, the well-known mammalogist, has transmitted to me for examination five specimens of this species, which were collected alive in a pond on Monument Creek, in Colorado Springs, Colorado, by C. F. Fraker. I wrote to Dr. F. C. Schneider, of Colorado College, which is located in that city, and asked him to look for more. He replied: "We went to the pond in Monument Park for them, only to find it recently cleaned. As a result we took only one specimen." There is no reason to doubt that the species has been accidentally introduced there, and it will be interesting to note how it withstands the climatic conditions of Colorado, at an altitude of nearly 6,000 feet.

Junius Henderson,
University of Colorado, Boulder, Colo.

The Nautilus.

Vol. XXVI. DECEMBER, 1912. No. 8

CONCHOLOGICAL CHAT FROM LONDON.

BY CHARLES HEDLEY.

Like so many kindred institutions the Natural History Museum at South Kensington has suffered from its architect. Space and funds have been lavished on fantastic decoration which would have gone far to improve laboratory accommodation. Our subject has suffered with others, perhaps more than others, from cramped space and poor illumination.

The hall devoted to conchology is about a hundred and forty feet long and forty broad, and is lit by windows in the roof. From the ceiling hangs two huge American models of Cephalopoda with outstretched tentacles. These fearsome things are the principal attraction for the casual visitor.

Around the walls special series are displayed. The wants of the local students are satisfied by an excellent exhibition of British Mollusca; the slugs and minute shells being shown by drawings.

General morphology is admirably illustrated by models, figures and specimens, both whole, sectioned and dissected. The operculum for instance is traced through various modifications. Branchiæ and other organs are shown in large models. There are micro-photographs of lingual ribbons. A movable frame imitates the torsion of the prosobranchiate viscera. Monstrosities have a case to themselves. Application of shells to arts and industries, to wit, ostreaculture, cameo cutting and the pearling industry are appropriately

displayed. Parts of perforated hulls or piles exemplify the damage inflicted on the works of man by the shipworm.

In this high latitude and damp climate the colors fade but little. Protection in this respect is afforded by drawing the window blinds on a bright day.

The floor of the hall is occupied by about two hundred glass-topped double-deck cases standing waist-high above chests of locked drawers. These run in four rows down the room and are divided by broad gangways.

A modern policy in museum administration is to divide collections into a general exhibit and a special study series. The former presents an epitome of the subject for the use of the public or the beginner in science, the latter to contain types and full series for the study of the specialist. Such special collections are kept in rooms from which the public is excluded, where books, microscopes and the like are available and where drawing or dissecting can be done. This arrangement is current in other departments at South Kensington but not in that of conchology.

Practically the whole of the museum collection of shells is exhibited or stored in the public hall. Shown in the glass cases to the general public are named examples of almost all the shells of the world.

When a specialist consults the collection, an attendant unlocks the drawers or cases indicated and waits beside the student till the case is to be locked up again. To the public an open case and a conchologist at work are part of the show. So he is tormented and distracted by rustics who try to read what he is writing in a note-book, by children who want to finger the priceless types he is examining, or by beginners who want his help in naming shells. If the public were excluded it still would be sufficiently difficult to make observations and comparisons with a pocket lens in a dim-lit hall and write with a fountain pen upon the cases for a table.

Recent acquisitions to the collection are mounted in black, square, glass-topped boxes, and labeled both under the lid and on the bottom of the box. The bulk of the collection is, however, gummed on paper-covered wooden tablets. Labeling consists of the separate specific name and author, written or printed on white paper in the lower center, the locality in the left lower corner. Where "type" occurs it is at the left top corner. The generic name in larger print

occupies a separate tablet. On the under surface of the tablet, H. C. in the corner, means that it was a part of the famous Hugh Cuming collection. Besides, the under surface sometimes has a wealth of scientific information ; critical notes by those who have worked upon the collection. Here the specialists gather a harvest of synonomy, references to literature, comparisons, collected notes on habitat, and so forth. The register number, including the year of accession, when it occurs, is given here. But the student may glance over the face of the collection and miss the information jotted down by his predecessors, but concealed below. Frequent reversal of the tablets helps to dislodge the specimens. Types are not invariably thus distinguished. I noticed several " Challenger " shells, which though evidently types, were not so marked. There is more strictness in this respect of late years than of former.

After a lapse of time the gum becomes brittle, so that the opening and shutting of a drawer or the lifting of a tablet may unfasten a shell. The loose shell may readily be returned to a wrong tablet and thus originate error. For this reason it is to be hoped that the whole collection will be ultimately transferred to boxes. Again in boxes they are easier to examine, far safer from dust and from accidental crushing.

In former years no clear distinction prevailed between the range in space of a species and the source of a particular specimen. So " localities " were " added " rather recklessly and perhaps sometimes erroneously to old specimens not provided with a " habitat." Even upon the original tablet of a type subsequent specimens have sometimes been intruded. Again in the union of names, a species was apt to be submerged under the name of another species with which it was thought identical. Varietal names were disallowed, so that an absolute synonym and a geographical race were treated alike in elimination.

On the whole, the nomenclature is well up to date. Recent suggestions by Dall and Pilsbry, and the restored names of neglected authors like Bolten and Perry, are incorporated.

Compared with other large museums the staff is weak. Instead of two or three or four naturalists that we find elsewhere, there is but a single scientific officer. He is aided by a couple of attendants who perform such mechanical work as labeling, registering or packing. There is no departmental artist.

The maintenance of so large a collection is sufficient occupation for the time of a single officer, and the increase of the collection is therefore slow. To attempt any field work is contrary to the traditions of the institution. Englishmen, both at home and abroad, loyally contribute to their great museum. But the casual efforts of amateurs do not give the results of professional collectors. So that additions are sporadic and usually consist of chance gifts from the friends of the museum and small purchases of dealers.

Unlike other museums of the front rank, South Kensington produces no official conchological publication and lacks the organization and inspiration which such would afford.

It follows that the London collection, which was once the largest in the world, will cease to be so in the near future, if indeed it has not already lost this supremacy.

This year the veteran Mr. E. A. Smith retires to enjoy a well-earned pension. For half a century he has helped the conchological workers of the world. And we from the ends of the earth, alike with these in the center of civilization, cannot find expressions to convey our appreciation of his patience, his geniality and his learning. We are like the Fijian chief who exclaimed, "Would that we were dogs, that by the wagging of our tails we could show what we feel."

TWO NEW AMERICAN LAND SHELLS COLLECTED BY MESSRS. HEBARD AND REHN.

BY H. A. PILSBRY.

The form described below was taken at Garrison, Montana by Messrs. Morgan Hebard and J. A. G. Rehn during their entomological expedition of 1909. On superficial examination it was at that time referred to *O. alpina* Elrod. A few specimens had been picked up also at Garrison by Mr. E. M. Kindle in 1907 and transmitted to me by Mr. J. H. Ferriss. Both lots consist of "dead" shells only. Doubtless where the "bones" are so abundant as to attract the attention of non-conchological naturalists, there are plenty of living snails to be found hard by.

OREOHELIX CARINIFERA, n. sp.

The shell is lenticular, carinate, umbilicate, the width of umbilicus between a fourth and a fifth that of the shell, whorls $4\frac{1}{2}$, slowly increasing, the first $2\frac{1}{2}$ strongly convex, obliquely striate, the striæ finer on the embryonic portion. Subsequent whorls are strongly convex around the upper (inner) part, becoming concave near the outer edge; the striation is rougher, and some weak traces of spiral striæ appear in places. The last whorl is noticeably concave above and below the peripheral keel; it descends very slightly or not at all in front, and on the base there are very inconspicuous, well spaced spirals composed of granules. The oblique alt. and the diameter of the apeture are equal, and there is a slight angle at the termination of the keel.

Alt. 5, diam. 9.4 mm.; width of umbilicus 2 mm.; oblique alt. and diam. of aperture 4 mm.

Garrison, Montana. Type and paratypes No. 99253 A. N. S. P.

Oreohelix alpina Elrod, which comes from high elevations (8,500–9,000 ft.) in the Mission Mountains, is about the size of this snail, but the whorls are less convex, the convexity of the later ones is simple, while in *carinifera* there is a concavity above the periphery. *O. alpina* has no spiral sculpture or granulation, and the keel is less pronounced. I regard such similarity as exists as due to convergence rather than to actual relationship. *O. hemphilli* Nc. differs in the sculpture and shape of the embryonic whorls.

O. carinifera is one of the smallest Oreohelices, the dimensions given above being those of the largest shell out of about 20 in the two lots seen. A very similar small form was taken with *O. haydeni* in the Wasatch Mountains many years ago by the Wheeler Expedition, but I think it will prove to be distinct. Its exact location in that range is not known, and it was not among the forms taken by Hemphill there.

HOLOSPIRA MESOLIA, n. sp.

The shell is cylindric-fusiform, of a delicate pink-white tint, the upper part white with blue stains; dead shells white throughout. Whorls 14, the first $2\frac{1}{2}$ smooth, forming the mamillar embryonic shell, the first whorl rapidly increasing, second swollen, next half whorl very narrow. Subsequent whorls are nearly flat, rather finely but strongly striate, regularly increasing to the 8th or 9th whorl,

where the shell reaches its greatest diameter. After that the whorls are nearly smooth, rather glossy, and the shell diminishes slowly in diameter to the base. The last two whorls have retractive axial ribs which gradually increase in strength, and are strongest on the straight part of the last whorl and base. The last whorl is somewhat compressed laterally and projects, carrying the aperture well forward. Aperture is very shortly ovate, almost circular, light brown within. Peristome very broad, flatly reflexed, white. The internal axis is smooth throughout, rather slender, tapering downwards, with a diameter of about 1 mm. in the widest part.

Length 23.5, greatest diam. 6 mm.; whorls 14.

Length 22.6, greatest diam. 6.3 mm.; whorls $13\frac{1}{2}$.

Length 20, greatest diam. 5.8 mm.; whorls $12\frac{3}{4}$.

Sanderson, Terrell Co., Texas, on a low limestone ledge along the railroad. Elevation 2800–2900 ft. Types no. 107001 A. N. S. P., collected August 25, 1912 by Messrs. Morgan Hebard and J. A. G. Rehn.

This handsome *Holospira* is most closely related to *H. semisculpta* Stearns, which was described from a canyon above San Carlos, Chihuahua, a place on the Mexican side of the Great Bend of the Rio Grande. Dr. Dall and Dr. Bartsch have kindly compared specimens with the type of *semisculpta*, and report that the new species "differs in the profile, which in your shell is more contracted toward the base, rendering it spindle-shaped, while the former is more cylindrical. The ribs in yours do not extend over so many of the basal whorls, and the expanded peristome gives it a very distinct appearance. It is doubtless a distinct species."

With the *Holospira* were found specimens of *Polygyra texasiana texasensis* and a *Succinea*.

SHELLS OF SOUTHEAST MISSOURI.

BY F. A. SAMPSON, COLUMBIA, MO.

During a late trip through southeast Missouri I had a few hours' time for collecting shells, and the result, combined with some former collecting, is here given.

Fern Glen, St. Louis County.

This is on the Merimac river, twenty-one miles from St. Louis. The rock exposures are of Chouteau and Burlington limestone, but none of the shells were found in abundance.

Polygyra zaleta Binn. These cannot be distinguished from Ohio specimens, and are of about the same size, up to 29 mm. diameter. From other places in Missouri I have the *P. zaleta ozarkensis* of 19 mm. or less diameter, contrasting quite sharply with these.

Polygyra thyroides Say. I found only dead shells, but doubt not that on the low ground living ones could be found.

Polygyra clausa Say. Dead shells not differing from others in the state.

Polygyra pennsylvanica Green. A single shell that looks like a postpliocene fossil, but I think it is not such. I have found it in the postpliocene at Providence, Boone county, and living as noted below.

Polygyra elevata Say. Typical shells.

Polygyra appressa Say. With rather heavy striation, and from $17\frac{1}{2}$ to 25 mm. diameter.

Polygyra fraudulenta Pils. Rather more numerous than any other kind, and not found in Missouri except in localities noted on this trip.

Polygyra inflecta Say. Three dead shells. The umbilicus of two was not covered. Similar specimens are noted from Shepard's Mountain and from Rivermines.

Polygyra hirsuta Say. A single shell. Pilsbry says that evidence is wanting to show that this is found west of the Mississippi and south of Kansas and Pettis County, Missouri whence I reported it some years ago. The shell from this place and those from two other places, hereafter noted are of this species; they certainly are neither *blandiana* nor *uncifera*.

Polygyra fraterna Say. With quite small umbilicus.

Polygyra dorfeuilliana Lea. This is the farthest north that I have found this species.

Pyramidula solitaria Say. A single shell, which like the *pennsylvanica* looks like a postpliocene fossil but I do not think it is such.

Circinaria concava Say. A number of dead shells up to 29 mm. diameter.

Bifidaria armifera Say.

Pupoides marginata Say.
Gastrodonta ligera Say.
Vitrea indentata Say.
Physa gyrina Say. In a small pond by the side of the railroad track.

From a pile of mussel shells left by a pearl-hunter I noted seventeen or eighteen species.

Kemmswick, Jefferson County.

Polygyra appressa Say. A search by the railroad and on the bluff above gave a single dead shell of this species, but no other.

Washington County.

Here we are getting away from the limestone formations, and shells are scarce.

Polygyra fraterna aliciæ Pils. Mineral Point.
Bifidaria armifera Say. Potosi.
Pupoides marginata Say. Potosi.
Zonitoides arboreus Say. Mineral Point.
Zonitoides minusculus Binn. Potosi.
Vitrea indentata Say. Potosi and Mineral Point.
Physa gyrina Say. In a spring branch at Potosi. In a large stream close by no shells were found. A swift-flowing clear stream at Mineral Point did not show any shells.

Ironton, Iron County.

Polygyra albolabris alleni Weth. A single dead shell was found on Shepard's Mountain adjoining the town. The other land shells were found at the same place.
Polygyra inflecta Say. The umbilicus of the single shell found was not covered.
Polygyra dorfeuilliana Lea.
Zonitoides arboreus Say.
Physa gyrina Say. In a spring branch. In a larger stream close by no shells were found.

St. Francois County.

Polygyra thyroides Say. Farmington and Rivermines, the latter 19½ to 25 mm. diameter.

Polygyra appressa Say. Farmington. One of the shells did not have the umbilicus covered.

Polygyra inflecta Say. Rivermines. In three of the shells the umbilicus was not covered.

Polygyra monodon Rack. Rivermines. The small, widely umbilicated shell, not distinguishable from the northern shell, except that the color is not uniform, but usually more or less in stripes with the lines of growth alternately light and dark-colored. A shell from Jackson, Cape Girardeau county, shows the same stripes.

Polygyra fraterna aliciæ Pils. Farmington and Rivermines.

Pyramidula solitaria Say. Farmington.

Pyramidula alternata Say. Farmington and Rivermines.

Pyramidula perspectiva Say. Farmington and Rivermines.

Bulimulus dealbatus Say. Rivermines.

Pupoides marginata Say. Rivermines and Farmington.

Bifidaria armifera Say. Rivermines and Farmington.

Zonitoides arboreus Say. Rivermines and Farmington.

Vitrea indentata Say. Rivermines.

Helicodiscus lineata Say. Farmington.

Cape Girardeau County.

Three places in this county were visited: Allanville in the White-water river bottom, Cape Girardeau on the Mississippi river, and Jackson on higher ground between.

Polygyra albolabris alleni Weth. Cape Girardeau.

Polygyra thyroides Say. Allanville and Cape Girardeau, the largest from the former, $28\frac{1}{2}$ mm. diameter.

Polygyra clausa Say. Jackson.

Polygyra pennsylvanica Green. Jackson. This is the only place in the State where living shells have been found.

Polygyra fraudulenta Pils. Allanville. Somewhat larger than those from Fern Glen, $17\frac{1}{2}$ mm. diameter.

Polygyra appressa Say. Cape Girardeau, Allanville. Those from Cape Giradean vary from the small light colored, very finely striated shells to the larger, dark colored, coarsely striated, but not the same as large, dark reddish colored ones from Allenville, which Pilsbry reports similar to some Arkansas and Illinois shells. They are an unusual variety. An interesting specimen shows that by some injury a part of the last whorl was broken away, and a new

lip was formed more than a half-whorl short of the original one. The original parietal tooth is more than a half-whorl beyond the new one.

Polygyra inflecta Say. Cape Girardeau.

Polygyra leporina Gld. Jackson.

Polygyra monodon Rack. Jackson. See remarks under St. Francois County.

Polygyra fraterna imperforata Pils. Cape Girardeau.

Polygyra hirsuta Say. Jackson and Allanville, the latter the larger.

Pyramidula alternata Say. Jackson and Allanville.

Pyramidula perspectiva Say. Allanville and Jackson.

Omphalina friabilis W. G. B. Cape Girardeau. Pilsbry and Ferriss distinguish this species from *fuliginosa* largely by the appearance of the apical whorls, these being smooth, whitish-corneous and unworn, while in southwestern *fuliginosa* the summit is invariably worn, the cuticle removed from the earlier whorls. The shells from Cape Girardeau are very fine, measuring 28 mm. diameter, and the apex not worn in the least.

Vitrea indentata Say. Allanville.

Zonitoides arboreus Say. Cape Girardeau and Allenville.

Helicodiscus lineata Say. Allanville.

Succinea sp. Cape Girardeau.

Physa gyrina Say. Allanville and Jackson. From a pond on the west side of the Whitewater river at Allanville. The shells were small; from one on the east side a single larger one and of short spire.

Sphaerium solidulum Prime. Jackson.

Musculium elevatum Hald. Jackson.

Musculium transversum Say. Jackson.

St. Genevieve, St. Genevieve County.

Polygyra appressa Say. Found more abundantly than any other, and more like those from central Missouri than others herein noted.

Polygyra fraterna Say.

Polygyra fraterna imperforata Pils. Of ten *fraterna* found six were imperforate, and the others almost so.

Polygyra inflecta Say.

Pupoides marginata Say.

Bifidaria contracta Say.
Bifidaria procera Gld.
Lymnæa obrussa Say. In a spring branch these and the two following were found.
Physa gyrina Say.
Pisidium sp.

SPHÆRIIDÆ, OLD AND NEW, II.

BY V. STERKI.

PISIDIUM PERALTUM St. THE NAUTILUS, XIV, p. 5, 1900.—Specimens have been received from some other places: near Douglas Lake, Cheboygan Co., Mich., collected by Mr. H. B. Baker, and Cedar Lake, Lake Co., Ill., by Dr. F. C. Baker. But those from Ky., Ill. and Ia., formerly referred to *peraltum*, are distinct and range with the following:

PISIDIUM FRAUDULENTUM, n. sp.—Mussel of medium size, barely longer than high, somewhat oblique, rather well inflated; superior margin curved, passing with angles into the adjoining; supero-anterior slope well marked, steeply oblique, straight or nearly so, anterior end a rounded angle situated near the ventral side, inferior margin rounded, posterior truncate at right angles to the longitudinal axis; beaks rather large, rounded or somewhat flattened on top, moderately projecting over the upper margin; surface dullish or with a silky gloss, rather smooth, with fine, crowded, somewhat irregular striæ; color corneous to yellowish; shell opaque or subtranslucent, rather strong, hinge strongly angular, curved, very stout, its whole surface rugulose; right cardinal tooth angular, enclosing a deep excavation for the left anterior, often emarginate in the middle, its anterior part thin, the posterior very thick and usually grooved; left anterior rather short and massive, strongly curved upward, with apex pointed, the posterior short, steep oblique, slightly curved; ligament and resilium rather short, the latter strong.

Long. 4.5, alt. 4.3, diam. 3 mm. (100 : 95 : 66).

Habitat: Va., Ky., Ill., Ia., Mo., Miss.; ditch on the Cameron Run, west of Alexandria, Va., No. 602, types, and Roach's Run, Va.,

opposite Washington, D. C. (No. 628), both in tide water, collected by myself, Oct. and Nov., 1896; Bowling Green, Ky., Habana, Ill., Iowa City, Ia.; as mentioned (under *P. peraltum*, l. c.); creek below Elizabethtown, Ill., collected by Mr. A. A. Hinkley, 1894 (received 1904); Flat Creek, Pettis Co., Mo., by Mr. F. A. Sampson, and from the same place in Mr. Bryant Walker's collection; northeastern Mississippi, collected by Mr. A. A. Hinkley.

This differs from *P. peraltum* as follows: the superior margin is somewhat less curved, the angle formed by it and the supero-anterior slope is more projecting; the mussel is larger but somewhat less inflated; the color is corneous to yellowish, that of *peraltum* is pale corneous to nearly colorless or grayish, and the shell is more translucent; the surface of *P. fraudulentum* is more dull. As written, these differences may not appear striking, but they are constant so far as known; *P. fraudulentum* is remarkably uniform.

It appears also that our species ranges near some forms of *P. compressum*, and is of the same group; but the mussel is not so oblique, the beaks are broader and without ridges; moreover, the two were found associated at several places, and distinct.

(*To be Continued.*)

NOTES.

Cincinnatia in the Delaware Drainage.—On October 1, 1912 Mr. Delos E. Culver collected some pond-weeds at the settling pond on Darby Creek near Addingham, Delaware Co., Pa., which contained living specimens of *Cincinnatia cincinnatiensis* Anth. and the following other species of shells: *Planorbis antrosus* Conr. *parvus* Say; *dilatatus* Gld.; *Lymnaea columella* Say; *Physa heterostropha* Say; *Campeloma decisum* Say (one specimen is sinistral); *Amnicola limosa* Say; *Lyogyrus granum* Say; *Musculium transversum* Say; *Pisidium noveboracense* Pr.; *variabile* Pr.; *compressum* Prime. So far as I know, this is the only record of *C. cincinnatiensis* in the Delaware drainage, or anywhere between New York state and the Potomac River.—E. G. Vanatta.



1-3. CYPRÆA HIRASEI ROBERTS.
4. OVULA HIRASEI PILS.
5-8. CYPRÆA HUNGERFORDI KIIENSIS ROBERTS.
9. OVULA NIPPONENSIS PILS.
10, 11. TRIVIA EOS ROBERTS.

The Nautilus.

Vol. XXVI. JANUARY, 1913. No. 9

NEW CYPRÆIDÆ.

BY S. RAYMOND ROBERTS.

Recently there was sent to me for examination by Mr. Y. Hirase, Kyoto, Japan, a small but extremely interesting lot of shells, mostly belonging to the *Cypræidæ*, two species of which I believe to be new to science.

CYPRÆA HIRASEI, n. sp. Plate 7, figs. 1, 2, 3.

Shell pyriformly ovate, extremities produced, base convex; only moderately calloused laterally, the callus extending well up the dome on the left side, but not on the right. Dorsal surface longitudinally and spirally faintly ridged or malleated in irregular spiral series,[1] this sculpture obsolete towards the anterior end. Aperture nearly straight, the upper fourth a little curved; inferior third a little dilated; teeth small, about 27 on each side and well within the aperture. Ground work of dorsal surface creamy white, three banded with flesh color, over which there is a tracery or imperfect reticulation and near the borders of the lateral calluses a few blots of yellowish brown; topped with a conspicuous irregular blotch of a darker shade. Extremities pink, base white, sides, especially the left side, faintly spotted pinkish brown.

Length 51, diam. 32, elevation 26 mm.

Habitat, Tanabe, Kii Coast, Japan (Hirase).

[1] This malleation, while conspicuous enough in the shell does not show in the photographic figure. It resembles the malleation of a rough *Lymnæa palustris*.

The sculpture and pattern of the dorsal surface of this beautiful shell resembles somewhat that of some ovate examples of *C. subviridis* Rve., it differs however in having the spire produced, instead of sunken; the aperture is less curved posteriorly, the teeth more numerous and the sides faintly spotted, the form of the columella within the aperture is also entirely different, being broad and excavated, with a second series of about 6 teeth deep within.

But one specimen was obtained which belongs to the collection of Mr. Hirase, in whose honor it gives me great pleasure to dedicate the species.

Mr. Hirase sent me also two specimens of a Cypraea which after as exhaustive a study as I have been able to make, I am inclined to consider a strongly marked variety if not sub-species of *C. hungerfordi* Sowb. The following description is therefore offered.

CYPRÆA HUNGERFORDI KIIENSIS, n. subsp. Pl. 7, figs. 5, 6, 7, 8

Shell pyriform, dorsal surface creamy white, closely freckled with light chestnut-brown spots and indistinctly banded with pale purplish-brown clouds or diffuse spots; extremities, sides and base salmon, the sides more or less spotted with blackish-brown, along the edges of the lateral calluses. Aperture nearly straight, teeth in outer lip rather coarse, 17 or 18 in number, those on columella marginal, white, somewhat smaller and extending well within the aperture.

Length 31, width 19¼, alt. 15½ mm.

Length 31, width 19, alt. 15 mm.

Habitat, Kii coast, Japan (Hirase). The dorsal surface of a specimen not fully mature is faintly ornamented with a broad, brownish band, the spots on the sides are more numerous and run together (fig. 9).

Compared with the description of the typical *C. hungerfordi*, a shell which I have never seen, it has fewer teeth and the margins are *not thickened.* This is obvious in figs. 5–7, which represent a shell certainly adult. The coloration is also quite different and were it not for the locality from which it was obtained, I might claim for it recognition as a full species.

There were also in the collection, much larger specimens of *C. cernica* Sowb. than I had before seen, and *Trivia insecta*, Mighels, *grando*, Gask., *oryza*, Lam. and *pilula* Kien. This last shell, which heretofore I had placed in the synonymy of *Trivia globosa* Gray, is

to my mind undoubtedly entitled to specific distinction, which more recent observers have accorded it.

Trivia eos, n. sp. Plate 7, figs. 10, 11.

Shell globosely ovate with rounded base, white, rather finely ribbed. Dorsal line indicated by a narrow smooth area from which ribs diverge in every direction, a few being interposed on both sides; posteriorly they pass directly over the spire, which projects well, showing five whorls. On the ventral side the spiral ribs have an oblique trend and extend well into the aperture, where they decrease in number and become coarser. The aperture is rather narrow, but slightly wider anteriorly, outer lip thickened outside, strongly ribbed transversely, with about 29 ridges forming teeth on the inner margin; retracted anteriorly. The inner lip bears a very small callous projection above, and near the base there is a rounded ridged lobe projecting inward.

Length 25.7, diam. 20, alt. 17 mm.

Habitat, Boshu, Japan (Hirase).

This magnificent *Trivia* is named for the "Sunrise Kingdom." Type in the collection of Mr. Hirase. In size it is approached only by *T. oniscus* Lam. of South Africa, which however differs by its pink color, larger smooth dorsal area, fewer ribs, shorter spire, and especially by its much wider aperture, as in other members of the South African group of Trivias, to which *T. eos* has but little affinity. It should be noted that the irregularity on the left side of fig. 10 is wholly an effect of light in the photograph, as that slope of the shell is perfect and evenly curved.

A COLLECTOR IN WESTERN CUBA AND THE ISLE OF PINES.

BY REV. H. E. WHEELER.

A mid-winter vacation is of such rare occurrence in a minister's career that the generosity of a congregation that made such a thing possible deserves special mention. It was my purpose to join my genial correspondent, Mr. Walter F. Webb, at Baracoa on the North Coast of Cuba, for which port he sailed early last January, but it turned out that we were separated by the length of the island, and my excursion to the Oriente Province was abandoned. Enroute

for Havana I had the unexpected good-fortune to be on the first regular train which passed over the sea-going extention of the Florida East Coast Railway, completing the line from Knight's Key to Key West.

Shortly after my arrival in Havana, I met Dr. Carlos de la Torre, Curator of the Zoological Department in the Academy of Sciences, Havana, and Professor of Biology and Zoology in the University of Havana, and before I could protest I found myself in the toils of characteristic Cuban hospitality. The Doctor's contributions to the literature of Cuban malacology are too well known to readers of the NAUTILUS to need any reference here, but it may not be so well known that he was the favorite pupil of Felipe Poey, the most brilliant naturalist which Cuba has produced.

To see his collection is worth the expense of a trip to the Antilles. So far as the described Cuban fauna is concerned it is almost complete. Here are specimens of nearly every species, and for the display of a variable shell like *Polymita picta*, Born, or *Coryda alaudù*, Fer., many ample drawers are required. You may see specimens of *Licina percrassa* (Wright) Pfr., rarest of Cuban mollusks, and of which no living examples have ever been found. Its habitat is Luis Lazo, Pinar del Rio Province, in one of the highest peaks of the Sierra de Los Organos.

In fact a drawer of Cuban shells calls up many a vivid scene and many an honored name among the students of molluscan life. Poey and Gundlach and Pfeiffer seem to spring out of the past, speaking with such versatility that you give them respectful audience; in your imagination Wright and Arango are on the trail of a hundred new species which unaccountably have been overlooked; the Count Morelet is on the high seas again exploring regions that no naturalist ever saw before him; and Férussac is rampant with chagrin that some oft-handled snail is crawling about with a name wholly unworthy of its race.

Once on Cuban soil, under the spell of matchless evening skies, the dormant or overworked energies aroused by all that a tropical winter can be, why should one sleep? A single rainy day at Ceiba Mocha, at Rangel, or on any limestone mountain-side, and how shall arid plains or denuded hillsides, where only shells of somber hue and homely feature care to dwell, claim any longer a collector's enthusiasm? For here is *life* and color and health, not only in the

lowly kingdom of crawling things, but for the student also who searches out their haunts.

It was quite natural that the first shells collected by the writer should have been those introduced by commerce, namely, *Rumina decollata*, Linn., and *Subulina octona*, Brug., but surely I must have seemed, both to natives and tourists, more like an escape from some asylum than a would-be naturalist, engaged, as I was, in turning over unceremoniously the flower-stands in the Parque de Colon, right in the heart of Havana.

It was 1878 that Arango published a catalogue[1] of Cuban shells which included also those from the Isle of Pines. In it 578 land forms alone are enumerated, but since that list was prepared more than a hundred species and sub-species have been described. But this is only a numerical evidence of the richness of the fauna.

Arango's collection is at the Academy of Sciences, and Gundlach's "Museum" may be seen at the Institute. His (Gundlach's) observations were published only after the most painstaking study; birds, reptiles, insects, and shells all claiming his interest. With a conscience for accuracy which is the naturalist's only passport in the scientific world, he would delay the taxidermy of a mammal or bird until he understood its anatomy and had become thoroughly familiar with its habits.

Though I have hardly a bowing acquaintance with ornithology, there were some birds which interested me. Among them was a Phæthon,[2] the nearly extinct "guacamayo" (*Ara tricolor*), and the "carpintero real" (*Picus bairdi*). The stately flamingo (*Phœnicopterus ruber* Linn.) is common enough, I suppose, but of irresistible plumage.

I was attracted by a turtle (*Emys rugosa*), which I saw in great numbers near Camoa, and my attention was called to a curious fresh-water fish[3] which has survived the evolutional changes of nobody knows how many thousand years. Here also could be

---

[1] "La Fauna Malacologica Cubana, par Rafæl Arango y Molina," Havana, 1878.

[2] *Phæthon flavirostris* Brandt, the only representative of the *Phæthontidæ* which visits our Eastern seaboard.

[3] *Lepistosteus* (*Atractosteus*) *tristœchus*, Block, a species still living in Mexico and the Southern United States.

studied that rare insectivorous mammal, which from its channeled teeth, is called *Solenodon* (*S. cabanus* Peters). There is a related species[1] in Santo Domingo, but nowhere else, I believe, except in Madagascar, is there an animal, the *Centetes*, at all like it. The Hutia, a large rodent, which in war times was hunted for food, is represented by three species,[2] but one of these, as well as the *Solenodon*, belongs to the fauna of Eastern Cuba.

At the University of Havana nothing is more touching than the regard in which Felipe Poey is held. His statue, a magnificent work of art, is the chief memorial in the quadrangle, and his library with many interesting relics, and many of his types have their repository in the spacious Museum. The present University, by the way, represents only one of the many far-seeing services rendered the Cuban people, during the American intervention, by Gen. Leonard Wood. He converted the old ordnance factories into University buildings which now fittingly crown the Vedado.

At the time of my visit the Professor's assistants were mounting a recently captured ray (*Lebisa*), a fish well known by name only to Poey and mentioned in his several works, but which has never been described.[3] This curious inhabitant of the Caribbean Sea is from one to two meters in diameter, having a rough spiny skin which has been used by the natives as a dish rag. It has a long, slender, sharply-pointed tail, near the base of which are erected two obliquely-set and forbidding spines, which look much like the stoutly grown thorns of a healthy locust tree.

For one who is interested in botany the notes of A. Richards in Ramon de la Sagra's "*Historia Fisica Politica y Natural de Cuba*," will be found instructive, and other works, such as the "*Flora Cubana*," of Sauvalle, the "*Flora Havanensis*," of Gomez de la Maza, and the "*Flora of the British West Indian Islands*," of Grisebach, have an honorable place among the older authorities.[4]

---

[1] *Solenodon paradoxus* Brandt.

[2] *Capromys fournieri*, Desmarest; *C. prehensilis*, Poeppig; and *C. melanurus*, Poey.

[3] The description will appear in Poey's "*Ictiologia Cubana*," which Dr. de la Torre is now editing for publication.

[4] This painstaking catalogue has the advantage of being written in English. Its author, A. H. R. Grisebach, M. D., was Professor of Botany in the University of Göttingen, and the first edition was published in London in 1864.

Of characteristic trees one should mention the gigantic *Ceiba*,[1] the "West-Indian God-tree," with its strangely distorted and buttressed trunks; the *Almacigo*,[2] whose maroon-red bark peels off much after the fashion of our sycamore; and a heavy-foliaged evergreen, the Indian laurel[3] everywhere abundant. The deep holes which often corrugate the trunk of the last-named species are favorite retreats for *Liguus fasciatus*, Müll., and the common *Eurycampta bonplandi*, Lam. On the Güiness road near Camoa, I took sixteen specimens of this noble shell from the "crevices" of a single tree.

At Matanzas I saw the *Salvadera*,[4] a rather large tree whose fruit, when fully ripe, discharges its wedge-shaped seed from a large spherical pod with a violent explosion. When such a catastrophe overtook one of my specimens in my room at the hotel, I am not sure but that the other guests had good cause to suspect a suicidal termination of an unclassified American.

On my trip to Camoa I collected a shrub, the leaflets of which strongly reminded me of our native holly. That evening my Cuban host, seeing what I had in my bag, held up his hands in horror, crying, "*Guao! Guao!*," and immediately calling some tongs into requisition, unceremoniously disposed of my beautiful plants. I had collected one of the most poisonous plants[5] on the Island. A terrible pain and swelling usually attends infection, often fever, and some natures are so sensitive to it that they cannot pass under its shade without being poisoned. But though it was prophesied that I would pay dearly the penalty of my ignorance, no appreciable inconvenience was suffered. Being already immune to its relative, our common poison-oak,[6] I am tempted to experiment next with that unnamed plant which Professor Simpson so utterly abhorred.[7]

---

[1] *Ceiba pentandra* Gaertn. —*Eriodendron anfractuosum D. C.*

[2] *Bursera gummifera* L., of the older authorities, now called *Elaphrium simaruba* (L.). This tree is common in Florida, there being known as the "*gumbo-limbo.*" [3] *Ficus nitida* L.

[4] *Hura crepitans* L., of the family *Euphorbiaceæ*, called in English-speaking countries "*sand-box.*"

[5] *Comocladia platyphylla.* Another member of this genus which occurs in the Western provinces of Cuba is *C. dentata.* The (*Hippomane mancinella* L.), however, is a much more dangerous plant, and immunity from one class of poisons does not justify such a prospect in respect to others.

[6] *Rhus radicans* Linn. [7] See "Collector's Journal," Vol. I, No. 4, in loco.

Ceiba Moche, a little village of thatched cabins near Matanzas, is a famous place for shells. Though the season was unusually dry, I collected the following species there in a few hours:

Oleacina oleacea, Fér.
Oleacina solidula, Pfr.
Thysanophora vortex, Pfr.
Cysticopsis debilis, Pfr.
Cysticopsis cubensis, Pfr.
Helix (Microphysa) stigmatica, Pfr.
Jeanneretia multistriata, Desh.
Eurycampta bonplandi, Lam.
Thelidomus auricoma, Fér.
Liguus fasciatus, Müll.

Urocoptis scaeva, Gund. One of the two sinistral representatives of this genus found in Cuba, the other being Urocoptis coronadoi, Ar., from Marianao, near Havana.

Urocoptis fortis, Gund.
Urocoptis gracillima, Poey.

Urocoptis philippiana, Pfr. This beautiful species has not been previously reported from this locality. It had chosen a piece of wall unoccupied by other Urocoptids, and seemed to have developed a prosperous colony.

Megalomastoma apertum, Poey.
Ctenopoma rugulosum, Pfr.
Chondropoma dentatum, Say.

Chondropoma pictum, Pfr. This species is distinct from *pfeifferianum*, Poey. The animal differs by having red tentacles, and in the shell the peristome is simple. The sexes are separate, the male being always about one-third smaller than the female. On the female shell is generally to be seen a white mark caused by the abrasive contact of the male.

Eutrochatella sloanei, Orb.
Helicina adspersa, Pfr.
Alcadia hispida, Pfr.
Emoda submarginata, Gray.
Alcadia nitida, Pfr.

The *Jeanneretia multistriata* calls to mind a bit of history. Ch. Jeanneret was a Swiss watch-maker who, in 1868, was living at Santiago de Cuba. He discovered many important species, and was a valued correspondent both of Gundlach and Poey. But being taken for a spy during one of the political uprisings he was shot from across the river. His collection was lost. The subgenus *Jeanneretia*, erected by Pfeiffer in his honor, is a genuine Cuban group, and is remarkable for its spiral ridges. Its most typical representative is *Jenneretia sagraiana*, named by A. d'Orbigny in honor of Don Ramon de la Sagra, the author of the famous "*Historia fisica, politica y natural de la Isla de Cuba*," 10 volumes, Paris, 1840–1850.

Another species which bears the name of an almost forgotten naturalist is *Helix parraiana*, Orb. D. Antonio Parra was a native of Spain, and has the distinction of having written the first work on Cuban fishes. This was published in 1787, the engravings being made on wood by his son. His expedition was furnished by the Museum of Madrid, but his work on the fishes and crustaceans was printed in Cuba.

The Loma de Camoa[1] is a locality which has been thoroughly explored, but as it can be reached in an hour from Havana on the electric railway, it is impossible to pass it by. Here I collected the following species:

Oleacina oleacea, Fér. Polygyra paludosa, Pfr.
Oleacina solidula, Pfr. Helix (Microphysa?) stigmatica, Pfr.
Oleacina subulata, Pfr. Cysticopsis cubensis, Pfr.
Thysanophora vortex, Pfr. Jeanneretia multistriata, Desh.
Thelidomus auricoma, Fér. Eurycampta bonplandi, Lam.

Liguus fasciatus, Müll. At Cardenas this species is white, generally devoid of all markings, save occasionally a few green lines on the last whorl. But at Matanzas—less than twenty-five miles to the west—it is richly adorned, and like Joseph's coat, with many colors, and the apex is almost invariably a rosy hue.

Urocoptis humboltiana, Pfr. Urocoptis cyclostoma, Pfr.

Urocoptis ischna, Pils. I am convinced that ischna is only a form of cyclostoma. The smooth form is generally found in the stone walls at the base of the hill, and on either side of the provincial road, but with it is a costate form that links it with the ribbed variety. This (ischna) is usually collected on the exposed cliff above where the heat of the sun would supposedly develop such a protective sculpture, and a series of specimens can be laid down showing the imperceptible transition from the typical and original form down to the elaborately ribbed variety.

Succinea sagra, Orb. Eutrochatella sloanei, Orb.
Cistula catenata, Mke. Eutrochatella babei, Arango.
Chondropoma dentatum, Say. Helicina adspersa, Pfr.
Chondropoma pfeifferianum, Poey. Alcadia nitida, Pfr.
Proserpina depressa, Orb., a species without operculum.

[1] *Loma* = hill; otherwise known as "Jamaica," and for that reason a frequent occasion for confusion of localities.

The following species are also known from this locality:

Hyalinia gundlachi, Pfr. Macroceramus unicarinatus, Lam.

Microceramus turricula, var. perconicus, Pils.

Melaniella acuticostata, Orb. Urocoptis camoensis, Pfr. and

Vertigo pellucida, Pfr., which I found abundantly at Mariano. In a small "laguna" near Camoa I found *Ampullaria conica*, Wood, a *Planorbis*, *Ancylus havanensis*, Pfr., and *Physa cubensis*, Pfr.

At Guanabacoa in the Havana Province there is a vigorous colony of *Carocolus sagemon*, var. *semiaperta*, Mart., which has a curious history. The species belongs to Eastern Cuba. Some fifty years ago Dr. Nicolás Gutierrez, the founder of the Academy, and one of Cuba's most distinguished scientists, put some young *sagemons* in his garden at S. Francisco de Paula, near Guanabacoa. They were typical *Caracolus arangiana* in different colors, some with and some without umbilicus. But all are now umbilicated, and correspond precisely with the variety named by Martens, *semiaperta* (not in Arango), which came originally from the Cauto river near Bayamo.[1]

Cardenas and Matanzas are well worth a visit whether one has visions of snails or not. The former city—the fifth in size on the Island[2]—is 106 miles from Havana by rail, and its commerce in honey, mahogany and asphalt has been augmented greatly by American enterprise. Indeed Cardenas has been called an American city, though I saw fewer Americans and heard less English there than in other western ports. However, the whole city turned out on Sunday to witness the increasingly popular American game—baseball.

I heard a missionary, whose church joined hard against a blacksmith shop, patiently compete for a hearing with its busy anvil, and just across the narrow street rang deafeningly and incessantly the electric gong of a moving picture show, which form of amusement has become as fully identified with Cuban life as it has outrun our American taste for cheap entertainment. The beautiful Bay is very ample but shallow, and lies over extensive beds of asphalt. The water is almost fresh from the constant upward flow of some mighty subterranean river, or perhaps from many streams flowing through the limestone rock of this region.

Here I found flourishing churches, well disciplined schools, and

---

[1] Prof. Pilsbry considers *semiaperta* only a variety of *arangiana*.

[2] Its population is about 25,000.

an extensive Museum, whose director, Sr. Francisco Blanes, is an ardent conchologist as well as a devout Christian. In fact, the Museum owes many of its choicest collections to this generosity, and in token of their appreciation the Trustees have given him a life pension and made him honorary curator.

In the court of the new Museum building *Bulimulus sepulchralis*, Poey and *Ennea bicolor*, Hutt., the latter introduced within the past decade, had found a congenial home.

At Matanzas I was the recipient of many courtesies. Dr. Luis A. Cuni, Professor of Natural Sciences in the Instituto de Matanzas, gave me an automobile excursion through the beautiful Yumuri Valley, which lies under the protecting care of the Palenque de Matanzas, and its twin "loaf" the Pan de Matanzas.[1] We ascended the San Juan river in a trim launch getting a magnificent view of the verdant meadows and the palm-crested mountains. At the Bellamar caves our guide was an old Cuban who claimed to be their discoverer fifty years ago, and who seemed to be veritably a part of their grotesque formations. With Dr. Valdez graciously serving as interpreter, Dr. Cuni and I compared notes on Natural History, and parted the best of friends.

To the west of the city rises the Loma de Simpson. On its summit is a hermitage—the Ermita de Montserrate—which preserves a cherished memorial of Spanish ecclesiastical history. In this revered situation I collected two interesting *Urocoptids*—*U. poeyana*, Orb., and *U. elegans*, var. *auberiana*, Orb.,—and many beautiful *Liguus* from the thorny hedges of the *Euphorbia*.[2] I reached the crest of the hill in the early morning, and I shall not soon forget the panorama of La Cumbre, the mountains set against the deepest of blue skies, the Valle de Yumuri, a mighty garden of Royal Palms and tropical flowers, nor the beautiful city itself, whose buildings in dazzling white seemed to step down to the bay with stately tread. The Bay of Matanzas is an æsthetic challenge; not until you see for yourself, my gentle reader, will you believe that it has mastered the science of color and perfected the art of display.

---

[1] Both of these mountains are famous for their shells. Nothing is more attractive in the cabinet than a well selected series of *Chondropoma presasianum*, Gundl., and *Chondropoma irradians*, Shutt., which can be collected on both "pans."

[2] *Euphorbia lactea*, Han.

The Matanzas fauna was studied and described by Pfeiffer. Dr. L. Pfeiffer came to Cuba from Germany with Dr. Juan Gundlach, the ornithologist, and Edward Otto, the botanist, in 1839. Pfeiffer and Otto left after a residence of two months, but Gundlach remained, and for more than fifty years he devoted his talents to the exploration of Cuba's Natural History, becoming a worthy co-laborer of Poey.

(*To be concluded.*)

NOTES.

Dr. Pilsbry will sail from San Francisco Jan. 7th, for the Hawaiian Islands, where he will spend several months in studying *Achatinellidæ* for the Manual of Conchology. Articles and notes for publication in THE NAUTILUS should be sent, up to April 1st, to Mr. Johnson.

FEEDING HABITS OF ARIOLIMAX.—Dr. C. Hart Merriam who has been summering in the redwood forest of Marin County California sends some interesting notes about the feeding habits of *Ariolimax columbianus*. He observed it taking the fallen drupes of the California laurel (*Umbellularia*) which have a tough skin covering a layer of pulp outside of a hard kernel. The snail took the fruit by the end which had been attached to the stem and was able, by protruding the buccal mass, to eat out the pulp between skin and kernel for about half the surface of the fruit without breaking the rind. Subsequently it was observed to gnaw off the base of a toadstool about 3 inches in height with a stem a quarter of an inch in diameter, and make a meal of the stem. It is probable notwithstanding its "carnivorous" dentition that these slugs live chiefly on such vegetable matter. W. H. DALL.

MUSSELS IN A WATER MAIN.—The London Metropolitan Water Board officials have just made an extraordinary discovery. Recently it was found that a 36-in. water main at Hampton-on-Thames was showing signs of becoming choked. It was opened, and 90 tons of fresh-water mussels were taken out of a quarter of a mile of the main. The mussels were so tightly massed that the bore was reduced from 36 in. to 9 in.—*Popular Mechanics*, Nov. 1912.



NEW CUBAN UROCOPTIDÆ.

THE NAUTILUS.

VOL. XXVI. FEBRUARY, 1913. No. 10

TWO NEW CUBAN UROCOPTIDÆ.

BY H. A. PILSBRY AND J. B. HENDERSON, JR.

UROCOPTIS (GONGYLOSTOMA) LONGA, n. sp. Pl. VIII, figs. 9, 10.

The shell is cylindric or pillar-shaped, a trifle widest in the middle, tapering slightly to the rather wide truncation; whitish or pale brown, indistinctly mottled or rather distinctly streaked with brown; glossy, smooth, the last whorl thread-striate, contracted and angular above the origin of the neck, which is free and descends shortly. The aperture is somewhat oblique, subcircular. The peristome is rather broadly expanded, reflexed, white, usually brown-tinted below. Axis slender, encircled with a small, sub-basal lamella which is minutely spinose in the upper whorls.

Length 14.2, diam. 2.1, aperture 2 mm.; $13\frac{1}{2}$ whorls.

Length 15.5, diam. 2.1, mm.; $13\frac{1}{2}$ whorls.

A small rocky hill at Zaza del Medio, at the junction of the Sancti Spiritus branch with the Cuban Central R. R., Province of Santa Clara, collected by H. A. Pilsbry, April 15, 1904.

This species belongs to the group of *U. wrighti* as defined in the Manual of Conchology, vol. XV, p. 263. It differs from the species there described by its smooth surface. It is also different from the several new forms recently found by Professor de la Torre, all of which have been compared.

Urocoptis (Gongylostoma) cara n. sp. Plate VI, figs. 27–30.

The shell is very slender, pillar-shaped, of nearly equal diameter throughout, or tapering slightly towards the truncate summit; whitish, copiously mottled and streaked with brown; glossy, smooth (or rarely showing very weak traces of widely spaced striæ), the last whorl regularly thread-striate. The whorls are slightly convex, the last compressed towards its base, which is angular above the origin of the neck, last half whorl free, tubular and descending, rounded or having a weak keel below. The aperture is subcircular, white within, rather oblique; peristome broadly expanding, reflexed, white. The axis is encircled by two small, nearly equal lamellæ, the upper one more oblique, the lower thinner and shortly spinose. In the penultimate whorl the axis becomes noticeably gyrate and the upper lamella disappears.

Length 16.5, diam. 2.1, aperture 2.1 mm.; 14½ whorls.

Length 22.3, diam. 2.3, aperture 2.3 mm.; 20 whorls.

Length 13, diam. 2 mm.; 12½ whorls.

"San José rocks" about 2 miles northeast of Sancti Spiritus, Province of Santa Clara, Cuba, H. A. Pilsbry, April, 1904.

Several hundred specimens were taken from the vertical rocks where they clung in abundance, though in the main lodged in crevices and holes, like most of the slender Gongylostomæ, they vary widely in size and number of whorls. No complete individuals were found; and it appears that about 12 or 15 whorls are lost in large adults.

U. cara differs from species of the *U. wrighti* group by having two equal axial lamellæ. It is probably related to the unfigured *U. imparata* (Arango) which is described as fusiform-cylindric, whitish, with entire spire and 17 whorls.

Urocoptis hidalgoi cabrasensis, n. var. Pl. VIII, fig. 16.

Differs from *U. hidalgoi* by the sparse, weakly developed sutural nodules, shorter, less closely costate neck, and by having an intermediate columellar lamella between the two large ones in the antepenultimate whorl. The shell is corneous with white streaks and maculæ, smooth except on the last whorl. Length 13½, diam. in the middle 3 mm.; 9½ whorls remaining. Cerra de Cabras, Pinar del Rio, collected by J. B. Henderson, Jr., 1909.

A COLLECTOR IN WESTERN CUBA AND THE ISLE OF PINES.

BY REV. H. E. WHEELER.

(*Concluded from page* 108).

The last few days of my vacation were spent on an excursion to the Isle of Pines, and I was fortunate in having Dr. de la Torre for a travelling companion. Looking at the map one sees that in outline it is very like the profile of an ancient volante, the typical Cuban carriage, and this comparison has been handed down by those who travelled in that curious and now almost forgotten vehicle.

Baron von Humboldt and Aimé Bonpland visited the Island early in the nineteenth century, making important geological and botanical observations. The first conchologist to explore its shores was D. Jose M. Velasquez. This was sometime in the early thirties, but he published no description of the species collected. A little later Count Arthur Morelet made an expedition to the Antilles, sailed around the Isle of Pines, landing near the present capital, Nueva Gerona, on the Casas River. His book, entitled "*Testacea Novissima Insulæ Cubanæ et Americæ Centralis*," describes species collected here as well as in Cuba and Central America. The species described from the Isle of Pines were:

Oleacina folliculoris (= *O. subulata*, Pfr?).

Oleacina paragramma (= O. solidula, Pfr.).

Cylindrella (Urocoptis) pruinosa.

Cyclostoma semicanum, a very rare species from the South Coast.[2]

Cyclostoma pupoides.

Cyclostoma disjunctum (a synonym of C. moreletiana, Petit, disjunctum being preoccupied).

Helicina (Priotrochatella) constellata,[3] and Helicina scopulorum.

Poey described *Trochatella stellata*[4] from Velasquez' manuscript. This species is not found on the Sierra de Casas, the home of its nearest congener, *Trochatella constellata*, but on the Sierra de

[1] Published 1849-1851.

[2] At Carapachi Bei, meaning "Turtles' Bay," a name given to it by the fishermen of Cayman's Islands.

[3] See Revue Zool., 1847, p. 147.

[4] See Poey, Mem. I., pp. 117, 447; and Jay, Cat. 1850.

Caballos, a mountain some three miles to the East of Las Casas and across the Casas River.

Gundlach visited the Island about the year 1854 and collected all the species previously described and some others which were described by Poey and Pfeiffer. These were:

Chrondropoma dissolutum, Pfr., a species so near Chondropoma pictum that it may be worthy of a varietal position only.

Trochatella luteo-apicata, Poey. Trochatella callosa, Poey.

Helix (Jeanneretia) pityonesica, Pfr., which so far as studied does not appear to be distinguishable save, in size from the common multistriata.[1]

Helix (Cysticopsis) comes, Poey (near cubensis, Pfr.).

Glandinella (Bulimus) poeyanus, Pfr., together with a new genus, Pineria, which is represented by four species only two of which were described from the Isle of Pines. These were: Pineria beathiana, Poey, and Pineria terebra, Poey. The latter is not only one of the rarest, but also one of the most curious, of land shells.

The only addition to the fauna of the Isle of Pines since Gundlach is *Cerion pinerium*, Dall, and a variety of this species is claimed by its discoverer to be the smallest of the genus.

It was Dr. de la Torre's hope to find some new thing in the Sierra de la Canada in the interior of the Island. But this range proved to be a quartz formation covered with pine trees,[2] which surely have the right of way, so far as native vegetation is concerned, on the Island. The common grass of this region is *Hypericum galioides*, Lam. At the foot of the mountain, however, we collected

Oleacina oleacea, Fer. Subulina octona, Brg.

Helix (Jeanneretia) pityonesica, Pfr. Thysanophora vortex, Pfr.

On the Sierra de Casas we collected all the known species from that locality, *Urocoptis pruinosa*, Mor., rather plentifully. There seemed to be two forms of this species, one which was stout and large, the other much more slender, but fully as long. Near Nueva

---

[1] The generic position of *multistriata* is still problematical, and can only be definitely established by an examination of the soft parts.

[2] *Pinus caribæa* Morelet. This tree grows also in Western Cuba, Florida and the Bahamas, and until recently (1904) has been confused with *Pinus cubensis* Griseb., the pine of Eastern Cuba.

Gerona we collected two fresh-water species, *Ampullaria conica*, Wood, and *Physa cubensis*, Pfr.

Everywhere we saw great colonies of ants[1] which are there so destructive to vegetation. One of the most dread pests, however, is the *Jejene*[2] (pronounced hay-hay-nie), a tiny fly, smaller than a gnat, whose sting, while not very painful, raises innumerable little bumps on the skin which smart unpleasantly. They travel in swarms and are not barred by ordinary mosquito netting, and, their favorite time for attack being in the night, one can only escape their ravages under a canopy of closely woven goods. When the Cuban wishes to take off a conceited man, he says: "He knows also where the jejene lays his eggs," something, by the way, which even the Naturalist has not yet discovered.

Of the parrots the *Ara tricolor* is about extinct, but the "*Catarra*" (*Chrysolis leucocephala*), the white-headed parrot, is still abundant, as well as the "*Perico*" (*Conurus evops*), a small paroquet with a long tail. Some six thousand of these birds are exported annually for pets, but to be trained to talk must be taken from the nest. The rare "*Cayama*," known to us as the Wood Ibis (*Tantalus loculator*, Linn.), seems here to adorn itself with all the glory it can borrow in a tropical environment.

The giant among the trees of this Island is the *Yagruma*,[3] whose dead leaves infold themselves making a good retreat for snails and wasps alike. One of the abundant wild fruits is the *Icaco*[4] used in preparing a sweet jelly, but this is found also in the coasts of Cuba. At the foot of the Sierra de Casas we saw that peculiar *Anacardium*[5] which produces a huge seed, shaped like a pear, externally to the fruit. The bottle palm,[6] called in Spanish *Barrigona*, having a bulge in its trunk like an Indian club, could not fail to attract attention, and another common palm was the *Miraguana*,[6] with slender trunk, whose leaves filled out a circle as perfectly as a daisy.

---

[1] Spanish "Bibijagua," *Atta cephalotes*.

[2] *Œcacta furens*, Poey, not to be mistaken for the larger and less vivacious sand-fly.

[3] *Cecropia peltata*. See Prof. Simpson's reference to this and other Cuban trees in the "Collectors' Journal," Vol. I, Nos. 3, 4 and 5. His paper is entitled: "Notes from a Collecting Trip in Cuba."

[4] *Anacardium occidentale*, L. [5] *Colpothrinax wrightii* Griseb. and Wendland.

[6] *Colpothrinax miraguano*.

What of commercial opportunities in the Isle of Pines? Opinions will vary. But some indication may be had in the fact that out of a total population of eight thousand, five thousand are Americans. The land itself is by no means fertile. There is nothing specially inviting in the scenery, excepting the North Coast, and educational advantages are quite meager. Schools and churches there are, and many cultured people, but morals are at a low ebb generally. The climate is the Island's greatest asset, and its chief products will always be the citrus fruits.

NEW JAPANESE OVULIDÆ.

BY H. A. PILSBRY.

OVULA (NEOSIMNIA) HIRASEI, n. sp. Plate 7, fig. 4.

The shell is fusiform, its greatest diameter contained 2.8 times in the length; thin, bluish white, the ends pale ochre colored for a distance of 4 or 5 mm.; glossy, under the lens showing distinct very fine longitudinal striation, and about 6 oblique, low and well separated spiral cords at the lower end, one or two more than this at the summit. The rest of the surface has a microscopic spiral striation, so minute as to be hardly noticeable except under the compound microscope. The two ends are about equally produced, pointed. Outer lip thickened outside and within, retracted at both ends, very indistinctly angular near the lower end, columella straight and simple. Near the summit there is a low, indistinct spiral swelling very obliquely encircling the axis. Length 28, diam. 10 mm.

Province of Tosa, Japan, Y. Hirase.

This species is longer and less swollen than *O. sowerbyana* Weinkauft (= *Ovulum spelta* Sowb., Thes. p. 480, pl. 100, f. 63, 64, not of Linné). It is decidedly more inflated than *O. carpenteri* Dkr. The axial fold above is quite weak in *O. hirasei*, and the coloration is characteristic.

OVULA NIPPONENSIS, n. sp. Plate 7, fig. 9.

The shell is oval, angular at the ends, glossy, bright flesh-pink gradually becoming paler towards the lip; very indistinctly 4-banded with brown, maculate with brown on the back and behind

the ends of the outer lip, which is fleshy yellowish on the face. Sculpture of irregular growth-lines and weak, microscopic spiral striation, scarcely visible except under the compound microscope. At each end there are about seven spiral grooves, which are widely separated except close to the ends. The outer lip is very convex on the face, somewhat thickened outside, incurved and armed with about 28 unequal, not very strong teeth. The grooves between the teeth extend as far as the middle of the face of the lip. The columella bears a strong white fold, terminating outwardly in a vertical callus on the ventral side of the base of the shell. There is scarcely any anterior notch. Posteriorly the callus of the peristome extends downward from the apex and towards the aperture. Deep within the aperture there is a longitudinal callous ridge on the parietal wall of the aperture. Length 13.5, diam. 9 mm.

Fakara, Awaji, Japan, Y. Hirase.

This belongs to a rather numerous group of species, among which it is nearest to *O. bulla* Adams and Reeve (Zool. Samarang p. 21, pl. 6, fig. 5). The new form is less narrow anteriorly and differently colored. It is blunter at the ends than *O. punctata*.

A NEW LYMNAEA FROM MONTANA.

BY FRANK C. BAKER.

GALBA MONTANENSIS, n. sp.

Shell of medium size, rather thin, translucent, ovate-turreted; periostracum light horn color; surface shining, lines of growth distinct, crossed by very fine wavy spiral lines; whorls six, convex, the body whorl somewhat obese; sutures deeply impressed; spire acute, longer than the aperture; aperture ovate, outer lip thin; inner lip wide, somewhat triangular, reflexed over the umbilical region; there is no distinct axial plait, but the inner lip is slightly indented where it touches the parietal wall; the umbilical chink is narrowly open. There are two rest period marks on one adult specimen.

Length 14.00; width 6.50; aperture length, 6.80; width 4.00 mill. adult.

Length 9.00; width 5.00; aperture length, 5.50; width 3.00 mill. juv.

Length 14.25; width 8.00; aperture length, 7.00; width 4.00 mill. adult.

Length 10.00; width 5.25; aperture length, 5.00; width 3.00 mill. juv.

Type locality. Hayes Creek, near Ward, Montana, in the Bitter Root Mountains, altitude 3825 feet.

Types in collection of L. E. Daniels.

This Lymnæid belongs to the *palustris* group and is related to *Galba vahlii* ("Beck" Möller). It differs from that species in being more solid, the periostracum more polished, and the aperture rounder. The greatest difference is in the inner lip, which is roundly and evenly reflexed over the umbilicus, lacking the twist so noticeable in this group, and resembling in this respect such species as *Galba bulimoides cockerelli* and *Galba caperata*. Half-grown specimens might easily be mistaken for *Galba bulimoides*, the spiral lines and narrow umbilical chink, however, easily separating them. *Vahlii* has not been found south of the 55th parallel, and there is, therefore, a wide gap in the range of the two species.

The specimens at hand show some variation, one specimen being very obese; while another, which is believed to be typical, has a rather long, turreted spire. The young, half-grown specimens are always obese. The specimens were collected by Mr. L. E. Daniels in April 1912.

RADIX AURICULARIA AT CAMBRIDGE, MASSACHUSETTS.

BY W. F. CLAPP.

About four years ago, a dam was built across the Charles River at Boston. Above this dam, nearly four miles of the river, which was formerly salt and full of marine life, is now fresh water. In the NAUTILUS, Vol. 25, p. 132, Mr. C. W. Johnson writes of a colony of *Modiolus demissus* v. *plicatulus* Lam., which until a few years ago, lived about two miles above the dam. I collected a very fine set of *Paludestrina salsa* Pils., and a few *Odostomia bisuturalis* and *trifida*, in 1893, nearly three miles above where the dam now stands. But records of the marine molluscan fauna of this part of the river seem to be very scarce.

In August, 1912, Dr. Walter Faxon brought me three specimens of *Radix auricularia* (Linn.), which he had found very near the place where I had collected *P. salsa*, and *O. bisuturalis* and *trifida* in 1893. In October, 1912, Dr. Faxon and I collecting again at this spot, secured several more specimens, varying in age from very young to fully mature. All we found were on *Chara*, and in more or less deep and exposed parts of the river. I examined quantities of *Potamogeton* without securing a specimen, but nearly every clump of *Chara* yielded at least one. In the shallow sheltered coves and ditches we found great quantities of *Galba palustris*, *Planorbis trivolvis*, *Planorbis albus*, *Physa heterostropha* and *Ancylus parallelus* but these species all avoided the deep water of the river as surely as *R. auricularia* kept away from the shallow places. I have never found *Galba palustris*, *P. trivolvis*, or *R. auricularia* in that part of the river above Watertown which has always been fresh water. The common species there are *Pseudo succinea columella* and *Planorbis bicarinatus*, neither of which have yet been found in the river at Cambridge.

There are numerous explanations for the sudden appearance of exotic species in unexpected localities. A very popular, and probably in many cases a true one, is that they have escaped from some nearby greenhouse. In support of this theory, I will admit that there are several greenhouses in Cambridge, and that from some of them to the river would be but a short walk, but a comparatively long crawl.

SPHÆRIIDÆ OLD AND NEW, III.

BY V. STERKI.

P. COLUMBIANUM,[1] n. sp.—Mussel of medium size, strongly inflated, outlines (along the valve edges) oval to elliptic without any angles, beaks somewhat behind the middle, large, prominent, rounded or slightly flattened on top, or even calyculate; surface more or less uneven from lines of growth, somewhat shining, with fine irregular striæ, color light corneous to yellowish, often in alternating zones, shell thin, sub-translucent; hinge rather slight, cardinal teeth small, the right curved with the posterior end thicker, left

[1] The name is not strictly in conformity with the rules, but appears preferable to *columbiense*.

anterior with apex pointed, posterior quite short; ligament and resilium rather short and slight.

Long. 4.2, alt. 3.6, diam. 3 mm. (100 : 86 : 71).

Long. 3.3, alt. 2 6, diam. 2.3 mm. (100 : 79 : 70).

Hab.: British Columbia, apparently widely distributed and common, and rather variable with respect to size and shape (no doubt, also in Washington, etc.). Vicinity of Esquimalt, collected by Mr. Taylor over twenty years ago, sent by Justice F. R. Latchford, No. 6362 cotypes with the lot in Mr. Latchford's collection, and lots simply marked "B. C." at least some of them from that vicinity, are in various collections; Chilliwack Creek and Lake, B. C., collected by a member of the Canada Geol. Surv. staff, sent by Mr. Whiteaves; a marsh, Duncans, B. C., received from Mr. A. W. Hanham. The first specimens were received in 1895, and the species has been regarded as distinct ever since.

P. furcatum, n. sp.—Mussel oblique, medium inflated, superior margin little curved, posterior and inferior forming one continuous curve to the rounded-angular anterior end, or the posterior margin slightly subtruncate; supero-anterior slope at an angle with the upper margin, well marked, straight or little curved, at an angle of alt. 45° to the longitudinal axis; beaks slightly behind the middle, rather large, rounded or slightly flattened on top; scutum and scutellum slightly marked, small; surface polished, with slight, shallow striæ; color plumbeous around the beaks, straw along the margins, the two shades not sharply defined, shell subtranslucent in the plumbeous area, rather opaque in the light zones; hinge moderately strong, right cardinal tooth curved with the posterior part bifid, the outer, or posterior, shank separated by a groove even above, straight or curved with the convexity below, left anterior short with apex rounded, posterior oblique, slightly curved, the space between them wide posteriorly; outer laminæ of the right valve small; ligament and resilium short, moderately strong.

Long. 4, alt. 3.6, diam. 2.3 mm. (100 : 90 : 57).

Hab.: Green Lake, Seattle, Washington, collected by Mr. P. B. Randolph, with *P. idahoense* Roper and others; types acc. No. 3896 b, part, of the Carnegie Museum, Pittsburgh. Probably none of the specimens are full-grown. The mussel resembles some forms of *P. variabile* Pme., but is more oblique, the color is different, and the cardinal teeth are markedly different; named from the strongly bifid, or bifurcate, right cardinal.

The mussel also somewhat resembles *P. randolphi* Roper, from Seattle, Washington, but is much more oblique, the beaks are larger, and the surface striæ are slighter, more shallow, and more distant.

P. (*furcatum* var.?) *rhombicum.*—Mussel smaller, more oblique, rhombic in outlines; beaks small, color more yellow; the posterior part of the right cardinal tooth is thin and plain and correspondingly the space between the two left ones is quite narrow; ligament short and slight. Long. 3, alt. 2.6, diam. 1.8 mm. (the largest specimen). Also probably none of these are full-grown. Hab.: with the preceding, in the same lot. Apparently of the same group with *P. furcatum*, to judge from the shape; more material will prove whether the two are distinct or forms of one species.

F. W. BRYANT.

Mr. F. W. Bryant, of San Diego, Cal., while visiting Hawaii, died of heart failure on October 23, 1912. Born in Ohio about 70 years ago, he was an old friend of the late Prof. A. G. Wetherby, with whom he often collected. He described several species of Californian land shells in THE NAUTILUS (vol. XIII, pages 122 and 143, 1900 and Vol. XVI, page 70, 1902). *Pyramidula bryanti* Harper, was named in his honor.

NOTES.

LAMARCK'S COLLECTION OF SHELLS.—We take the liberty of quoting from a letter to one of the editors written by Mr. Charles Hedley from Geneva, Oct. 18, '12. "I have just been through Lamarck's types and have been rewarded for my trouble by recovering two or three lost species. The Lamarekian collection was as you know bequeathed by Delessert to the Geneva Museum. It is in excellent preservation, kept locked up apart from the general collection in four cabinets. The shells are gummed on the usual tablets and in the case of the bivalves the original label in Lamarck's writing is pasted to the under surface of the tablets. Each tablet has a colored margin to indicate geographical division; thus Australasia dark blue, Indian ocean yellow. "There is also a copy of the

Animaux sans vertebres annotated by Lamarck himself. I supposed that this neat mounting was done by Chenu, but the present custodians did not know. The director and his assistants pay great regard to this collection but they have not a conchologist on the staff. One part of a memoir illustrating by photography the shells of Lamarck has already appeared. I saw proof of the next part which should be issued this year, and will include the bivalves.

"This Museum has also the complete collections of Brot, the specialist on *Melania*, who was a Geneva citizen. Also the collection of Bourguignat, who left it here."

Banded Polygyra mobiliana: Some months ago I received from Herbert H. Smith a few *Polygyra mobiliana* Lea from near Mobile, Ala. Most of the shells were immature but I noticed that three adults were distinctly banded and some of the young also showed a faint band. On bringing the matter to Mr. Smith's attention he examined all that he had remaining and found that the majority of them were banded. Since then Mr. L. H. McNeill of Mobile has collected numerous specimens and finds most of them banded. I have just received 17 specimens from him of which 15 are banded. The band is brown, much darker than the balance of the shell, and is located just above the periphery. Shells collected at Foley, Baldwin Co., are bandless and Smith says that he has seen no banded shells of this species except those collected near Mobile. Mr. McNeill informs me that he got a great many of this species by "sweeping" the grass at night with an insect net.—Geo. H. Clapp.

Murex fulvescens Sowerby: Some handsome specimens of this species, pure white in color and measuring about seven inches in length, were recently obtained by Mr. J. H. Holmes in the Gulf of Mexico, off Wakulla Co., Fla., at a depth of 125 feet.—C. W. J.

Pseudogalba, new name for Simpsonia.—Mr. Bryant Walker has called my attention to the name *Simpsonia*, which is preoccupied by Rochebrune (1905), for the Naiad previously known as *Chamberlaina duclerci*. The name *Pseudogalba* may be used in place of *Simpsonia* (Lymnæidæ of North and Middle America, page 236). It is perhaps expressive of the relationship of the group to *Galba*.—Frank C. Baker.

The Nautilus.

Vol. XXVI. MARCH, 1913. No. 11

COLLECTING IN PANAMA.

BY EUGENE W. PRESBREY.

In 1850, C. B. Adams, Professor of Zoology in Amherst College, spent six weeks collecting in the waters of Panama Bay. In 1852, under his own direction, he published a complete catalogue of species collected, with careful notes on synonymy, stations and number of each species taken.

The writer recently spent two weeks covering carefully exactly the same territory, from Old Panama to Taboga Island, about fifteen miles. It may interest the readers of The Nautilus to know what changes in numbers, stations and characteristics may occur in sixty-three years. Many changes have occurred, due probably to disturbed environment. Adams, in six weeks, took 41,835 specimens, 516 species! ! Of these there were 38,920 Gasteropoda, 376 species! ! In the same season of the year, from the same rocks and sands, with the same tides—eighteen to twenty-four feet—the writer, in two weeks, under most favorable conditions, was able to find only 112 species of Gasteropoda, and about 1000 specimens. Some of these species were not found at all by Adams, who was a careful collector, letting nothing escape him, not even numbers. But the peculiar and interesting fact is that few of the species found in great numbers by Adams are plentiful now, and many of those species are not to be found at all. In many stations other species have taken the place of those found by him.

Adams collected 4500 *Oliva volutella.* These are not there now. Two dead on the beach, and one alive, were all that I found. Of

Oliva porphyria, Adams found only a broken fragment. I hunted in vain for even that. I doubt if Panama is now its natural habitat, as the shell was scarce in 1850 and is not found there now. Adams took 1500 *Nassa panamensis* on the reef that juts out from the old sea wall. Not a single Nassa is to be found there now. Of *Nassa luteostoma*, Adams took 330 near the old sea wall. They are not to be found there now, they having migrated two miles across the bay where a bushel could be taken at one tide. Adams found fourteen species of Nassa, I found four. Adams found incredible numbers of Nerita, two species, took 3200 specimens. Of these only 400 were *Nerita scabricosta*, the only species to be found in considerable numbers now on the high-tide rocks. Adams found 29 species of Columbella, 3000 specimens ! ! Four good specimens of *C. harpaeformis*, brought up by hermit crabs, and half a dozen other species, beach worn, rewarded persistent search.

Monoceros cingulatum. Adams found only 75, most of them young. Of these one may get a thousand adults in one tide—but not at Panama, at Taboga Island. Adams took 3900 specimens of Litorina, 12 species. Now, only one species to be found in considerable numbers. Adams found 110 specimens of *Ricinula reeviana*, I found one.

Conus, then and now: *brunneus*, 4—0; *gladiator*, 70—1; *mahogani*, 17—0; *nux*, 2—12; *princeps*, 8—4; *purpurascens*, 12—1; *regularis*, 1—0; *mus*, 0—25. I was fortunate enough to find two specimens of *Conus panniculus* Lam. at Taboga, one alive, and both splendid specimens. Has this or any other of the Texti been recorded from the Panama Province? A fine specimen of *Conus fergusoni* Sby. was found, a species not recorded in the Conch. Iconica, and considered doubtful by Tryon. This specimen, after removal of the olive-green, persistent epidermis, shows perpendicular stripes of delicate yellow alternating with lavender, from rounded shoulder to base where the colors merge into pale ashy brown, columella and interior violaceous white. All other specimens that I have seen are dead white. The figure in Tryon's manual is not *C. fergusoni* either in form or color. *Fergusoni* has the rounded shoulder and heavy form of *Conus betulinus* or *C. glaucus*, depressed accuminated spire, whorls eight, rounded and canaliculate.

Purpura seems to be more plentiful than whem Adams was there. He found 8 species, 600 specimens. Two miles from Panama one

might take a thousand each of four species in a single tide. These seem to flourish under conditions that have driven other species away. I am unable to recognize many of Adams' species.

Many Marginellas dead in the sandy mud, few alive. Adams saw only fragments of *Strombus galea*, easily found now, both young and old. All the Strombi attributed to this province are found in good condition but, like other species, washed ashore. They are more plentiful than when Adams was there, but found at Taboga only. *Ranella caelata*, Adams took 190 specimens, alive, at half tide. None alive from high to low tide now. Crabs bring them up in great numbers and good condition.

Adams found Murex, 9 species. Found them alive, from half tide to low. Not now. Hermit crabs and diving black boys are the collectors now. Many beautiful and absolutely perfect specimens, too, are washed up dead, to half tide. I found six varieties.

Adams names only one species of Fusus (bellus). I took four species at Taboga, two species alive at half tide. Hermits could have furnished a peck. *Cancellaria* about the same but not alive. *Pleurotoma* abundant, but in no such numbers as indicated by Adams. Few alive. *Cerithium*. Adams found 14 species, 3400 specimens! ! Half a dozen species now brought up by hermit crabs.

Turritella banksi. Adams took 350 specimens, one-third of them alive. They have gone to deeper water but crabs bring up fine specimens. Adams found 90 specimens of *Pedipes angulata* under Panama sea wall, and 650 *Auricula panamensis* in same station. Not a specimen of anything to be found there now except Purpura in small numbers and Nerita out of water. Adams indicated scarcity of *Pecten* and *Pectunculus*. Now most plentiful decoration of the beaches. Adams found *Cypraea cervinetta, arabicula, punctulatis* and *pustulata* on the Panama reefs. Not one of these there now. There is no coral. To be found in small numbers at Taboga where there are small patches of coral to furnish food.

Recapitulating: for four miles either side of the Canal mouth in the Bay all but the coarser, sturdier forms have disappeared. Chitons, Purpura, Siphonaria and Fissurella are there, but few others, and all in small numbers. There is no food on the reefs. Everywhere, in the region covered by Adams, the stations and habits have materially changed. It might be interesting to compare individual specimens of to-day with those of same species taken by Adams

sixty-three years ago. Forces capable of changing habitat and habit might change form and characteristics.

My method in collecting is to traverse the beaches and from dead or broken fragments of shells, observation of tide currents, eddies, and prevailing winds, mark out probable stations at low water or beyond. The method is seldom wrong and leads to live specimens, either for wading or dredging, in shortest time.

Some species in Panama Bay have gone to other stations. Some have disappeared. Nearly all have sought deeper water and stay there.

Very solid volcanic rock, some of it nearly pure iron, is continuous from two miles west of Panama City to Taboga Island, fourteen miles. The Canal runs through part of it, and over the rest of this strata. Heavy dynamite blasts on the main land and on the islands in the Bay where Canal fortifications are to make another Gibraltar, can be felt distinctly at Taboga, too far away to be heard, sometimes. These concussions are frequent, occurring many times a day. A breakwater extends from the mouth of the Canal to the island fortifications. This breakwater is two miles long, and all day long trainloads of dynamited rock are dumped into the waters of the Bay on either side. It is possible that four years of concussion and waters more or less poisoned by nitroglycerene, plus the sewerage of Panama, may have destroyed food and driven the species to new stations, to deeper water, or worse, destroyed many of them. The volcanic reef ends abruptly a mile out and the water suddenly deepens to thousands of feet. At Taboga the conditions are more favorable and there is superb collecting ground, but even there low tide does not expose the live species—to any encouraging extent. The inference that all Panama species near the Canal have sought deeper water is justified.

There is a big suction dredge at Balboa near the mouth of the Canal. This dredge takes up everything from bed rock to top sand and sends a twelve-inch stream of water, mud, and millions of shells (seldom alive), a mile or two inland to make new land. Most of these shells have not been dead long, some are semi-fossil. There is fine collecting at the end of that pipe. But specimens taken there indicate wholesale destruction of molluscan life at the mouth of the Canal.

The most interesting "station" however on the Isthmus is in

Culebra Cut, 150 to 260 feet above mean tide level. There, in squeezed-up marine strata, easily recognizable marine species may be found in considerable numbers. Some are, perhaps, as old as the Tertiary period, but many of them can be duplicated, alive, on the Pacific beaches. Coral is found 262 feet above present mean tide. Fifteen miles away in the Chagres basin marine deposits are also found. Sometime, not so very long ago, there was an open strait where the Isthmus now is. Is the Isthmus younger, very much younger, than scientifically supposed? How much younger? Is there any evidence—not of its age—but of its youth, to be found in the five tons of shell-bearing material sent to Washington by the diggers in Culebra Cut? I don't know. I'm only a gatherer of shells—with an imagination and some disposition to ask questions.

ON SOME CUBAN UROCOPTIDÆ.

BY H. A. PILSBRY.

The following notes relate to new or rare species collected by the writer in 1904. My journey was a rather rapid one, undertaken with the object of seeing something of the mollusk fauna of the central part of the island, as nearly all the Cuban shells I had studied had been taken in the relatively far richer and more frequently explored regions from Havana Province west, and in Santiago Province, or Oriente as it is now called.

My route was from Havana to Cienfuegos by rail, thence to Casilda, the Port of Trinidad, by the Menendez steamship line, thence to Tunas de Zaza on the south coast; by rail then to Sancti Spiritus, eastward to Majagua in Camaguey Province, and return by way of Matanzas. Collecting was done at the places mentioned as well as at many places along the route, and others within a day's journey on foot or mule from those named.

Around Havana, Matanzas, etc., various well-known *Urocoptidæ* were taken which call for no special notice, and also several forms of the *U. elegans* group—a very difficult series, not yet worked up, and extremely abundant in the environs of Matanzas and in Havana Province. *Urocoptis cara* P. & H., *U. longa* P. & H., and the following species may be mentioned among the new forms taken.

Urocoptis (Cochlodinella) mediana, n. sp. Pl. VIII, figs. 1, 2, 3.

The shell is cylindric-fusiform, the upper half or third tapering slowly to the truncate summit; thin; whitish-corneous with some indistinct clear corneous maculation. Sculpture of thread-like rib-striae, equal to or narrower than their intervals, nearly straight, slightly oblique, and somewhat irregular, *as though a thread were here and there omitted.* Close behind the lip the striation is crowded. Suture distinctly denticulated by the rib-striæ. The whorls are slightly convex, the last very shortly free in front, with a cord-like basal keel. The aperture is oblique, circular; peristome expanded and shortly reflected. The axis is slender and simple, as in *U. poeyana.*

Length 18, diam. 4, diam. aperture 3 mm.; 11½ whorls.

Length 16.5, diam. 3.4, diam. aperture 3 mm.; 11 whorls.

Length 13.5, diam. 3, diam. aperture 2.4 mm.; 10 whorls.

Zaza del Medio, Province of Santa Clara, on a small rocky hill close to the Cuban Central R. R., under stones.

This species is closely related to *U. poeyana*, from which it differs by the coarser striation, which is moreover irregular in places, while in *U. poeyana* the sculpture is close and even.

Urocoptis (Gongylostoma) canteroiana (Arango). Pl. VIII, figs. 4, 5.

Cylindrella canteroiana Gundlach MS., Arago, Anales Real Acad. Ciencias Habana, XII, p. 284, 1876.

A translation of the very brief and unsatisfactory description of this species may be found in my Manual of Conchology, XV, p. 254. In 1904 I collected it on the fine hill "La Vigia," which forms the background of the town of Trinidad, and where Gundlach had collected the types a half-century before. I found it on the slope looking out over the Caribbean Sea, on the west slope, and in a cave near the foot. Also in the foothills of the Trinidad Mountains on the west side of the valley of Trinidad, and along the trail in these mountains about two miles from San Joan de Letran.

The shell is cylindrical, the upper half or third tapering to a rather narrow truncation; uniform light brown, with lighter riblets; somewhat glossy. Sculpture of even, rather strong rib-striæ, nearly straight except on the last whorl or two where they are slightly

curved, as wide as their intervals. Sutures rather deep, whorls slightly convex, the last rounded below, very shortly free in front. The aperture is somewhat oblique, shortly oval, a little longer than wide. Peristome continuous and free, very little expanded, somewhat thickened, narrowly reflected. Axis bears one thin, minutely crenulated spiral lamella.

Length 14, diam. 3 mm.; 13 whorls.

Length 12, diam. 2.6 mm.; 13½ whorls.

UROCOPTIS (GONGYLOSTOMA) CIENFUEGOSENSIS, n. sp. Pl. VIII, figs. 6, 7, 8.

The shell is cylindrical, the upper half or third tapering to a rather small truncation; white, mottled and streaked with corneous-gray, as in *U. variegata*; glossy. Sculpture of rather low, coarse rib-striæ, closer and sharper on the tapering whorls and on the neck behind the lip. Whorls slightly convex, the last rounded below, very shortly free in front. The aperture is slightly oblique and a little longer than wide. Peristome white, expanded and reflected, continuous and free. Axis encircled by a small, thin, minutely serrate lamella.

Length 15, diam. 3.1 mm.; 13 whorls.

Length 12.5, diam. 3 mm.; 12 whorls.

"Lagunilla," a place about 6 miles due north from Cienfuegos, where there are several small brackish lagoons.

This species has much external resemblance to *U. poeyana lacteoflua* Pils. It is most closely related to *U. canteroiana* Arango, from which it differs by the variegated coloration, weaker ribs and smaller axial lamella. It is the only species of *Urocoptis* I found in the environs of Cienfuegos. The small lagoons at Lagunilla are slightly salt to the taste. The rocks and sticks are thickly coated with *Congeria*, and there is also *Neritina virginea*.

MICROCERAMUS SANCTISPIRITENSIS, n. sp. Pl. VIII, figs. 11, 12, 13.

The shell is shortly rimate, oblong-territe, tapering from the last or the penult whorl in a long cone which is slender and straight-sided above. Ground-color brownish-corneous, becoming gray-corneous on the last whorl, the apex whorl or whorls brown or gray subsequent whorls with sculpture of rib-striæ which are white except

where interrupted by narrow patches or streaks of darker ground-color. On the base there are often several spiral corneous bands, leaving one to three spiral zones of long white granules. The rib-striæ are nearly as wide as their intervals, weaker in the corneous patches, and *continue upon the base.* Whorls rather convex, the upper ones more convex ; last whorl rounded, with a very weak spiral cord defining the base, sometimes almost obsolete. The aperture is oblique ; peristome well expanded, white, the columellar margin built forward, in a plane with the outer lip.

Length 12.7, diam. 5.5 mm.; whorls 10.

Length 11.3, diam. 5 mm.; whorls 9½.

Length 9, diam. 4.3 mm.; whorls 9.

Near Sancti Spiritus, Province of Santa Clara, at a place called "San José rocks," about two miles northeast of the city.

This is a larger shell than *M. petitianus* (Orb.), also known as *M. tunicula* (Pfr.), with stronger, closer striæ continuing over the base ; it is not so wide and the spire is less attenuated above. The smallest shell found in a large series is 9 mm. long. Nearly all are over 10 mm.

San José rocks, conspicuous from Sancti Spiritus as a white ridge, stand upon a low ridge of gneiss which runs north of Sancti Spiritus. The rocks cover only a small area, but form a labyrinth of fissures, caves, tunnels, deep holes and knife-like crests, the whole clothed with trees and vines. Deeper and deeper I went, getting big Megalomastomas and Helicea in the cool rock chambers. The charming new *Urocoptis cora* hung on the cliffs. Operculates quite new to me were turning up. Finally I was quite lost in this absurdly diminutive jungle. I had dropped down several cliffs of eight or ten feet in gaining the heart of the thicket, and there was nothing to do but go on in as straight a course as I could steer. After a hard scramble of half an hour I emerged on the lower side, wet with perspiration and scratched by the sharp rocks, but happy in the possession of a fine bag of snails.

The rocks of San José are apparently a remnant of the Oligocene deposits which must once have heavily blanketed the gneiss in this region. Little trace of these beds remains in the country around Sancti Spiritus.

I found a variety of this species at Zaza del Medio, where on a small rocky hill or on the adjacent slope to the river I picked up two

lots, of 13 and 5 specimens respectively.[1] In one lot the shells are more conic than the types, with the rib-striæ more delicate and sharp, obsolete on the base. Specimens measure:

Length 10.5, diam. 5 mm.; 9½ whorls.

Length 10, diam. 5 mm.; 9 whorls.

Length 9, diam. 4.2 mm.; 9 whorls.

In the other lot the shells are larger but also sculptured with delicate, thin rib-striæ, much narrower than their intervals. Length 13, diam. 6 mm., 10 whorls. This race may be called var. *tenuistriata* (pl. VIII, fig. 14).

MICROCERAMUS PERCONICUS TRINIDADENSIS, n. var. Pl. VIII, figs. 15, 17, 18.

The shell is imperforate, the last two whorls forming a broad, shortly cylindric part, those above rapidly tapering in form of a straight-pointed cone, slender above; rather glossy; corneous-brown, the last 4 or 5 whorls nearly covered with patches on which the striæ are opaque white. Whorls convex, the earlier ones more strongly so. Sculpture of regular, closely set, retractive striæ, as wide as their intervals, every second or third one a little enlarged at the upper end, forming *an irregular series of subsutural papillæ.* The last whorl is rounded, or there may be an inconspicuous very low cord upon which the striæ enlarge, defining the base. The aperture is oblique; peristome scarcely expanding, slightly thickened and white within, the interior elsewhere light brown. Columella vertical, straight, projecting forward a trifle near the insertion, the margin dilated above and adnate, not built forward.

Length 7.3, diam. above aperture 3.2, including aperture 3.75 mm.; whorls 8¾.

Length 7.7, diam. including aperture 3.9 mm.; 9 whorls.

Length 5.8, diam. including aperture 3.2 mm.; 8 whorls.

West side of "La Vigia," Trinidad, under stones, etc. Also on the northwest side.

M. perconicus trinidadensis stands near *M. angulosus* ("Gundl.," Pfr.) which it resembles in sculpture and in the structure of the columella, but it differs by the much more obese shape constant in a large number collected, and by the stronger sutural denticulation.

[1] Most of the small shells I got at Zaza del Medio were lost through an accident. These two lots were saved by being in bottles carried in my pockets.

The shape, indeed, is almost exactly that of the west Cuban *M. denticulatus*—a species less closely striated and having a rimate base and built-forward columellar margin. *M. petitianus* has narrower, more separated striæ and no denticulation below the suture; moreover, it has a shortly rimate base and forwardly-built columellar margin.

M. perconicus (which I described as a variety of *petitianus*) is doubtless a species distinct from *petitianus*, differing in sculpture and the columella, which is not in the least built forward. It is more strictly conic than *trinidadensis*, without sutural denticles, otherwise the same. *M. palenguensis*, which I have not seen, seems to be broader with a larger aperture.

The specimens recorded as *M. turricula* Pfr. from Trinidad were probably *trinidadensis*. I have not seen the form reported from Cabo Cruz.

MACROCERAMUS CANIMARENSIS (Pfr.).

A single specimen, typical in form and sculpture but white with a corneous base, I found among rocks in a precipitous, shaded place on the northwest side of La Vigia. It has already been reported from Trinidad.

MACROCERAMUS CANIMARENSIS ROTUNDIBASIS, n. subsp.

The shell differs from *canimarensis* by having only a very weak trace of the subperipheral keel (which is strongly developed in *canimarensis*); the base being much more convex than in *canimarensis*, and not so strongly striated.

"San José rocks" near Sancti Spiritus.

I did not find this species living, but dug it from a bed of clay formed by disintegration of the limestone.

A LIST OF MOLLUSCA FROM THE MUSSELSHELL VALLEY, MONTANA.

BY S. STILLMAN BERRY.

Except for the brief lists by Squyer (NAUTILUS, Vol. VIII, pp. 63–65, 1894) and Elrod (Bull. Univ. Montana, No. 10, pp. 170–174, 1902), there are very few molluscan records extant for the entire

state of Montana, so that the following list of species from the upper Musselshell Valley, in the central part of the state, is of interest from a distributional standpoint. The exact locality is Winnecock, Meagher County, where the writer has done a little collecting during August of two succeeding years (1911–1912). The fresh-water species were taken in the various springs and overflow pools bordering the Musselshell River, and the terrestrial species occurred under sticks, leaves and stones in the more moist places among the surrounding cottonwoods. No species were taken in the swift-flowing waters of the river itself, and no land forms were found at a greater distance from the river than a few rods. All except *Vertigo binneyana*, *Zonitoides minuscula* and *Planorbis callioglyptus* were of fairly frequent occurrence. The elevation of Winnecock is about 4000 feel about sea level.

Vallonia costata montana Sterki. The specimens appear to correspond very well with Dr. Sterki's diagnosis, and specimens sent to Mr. George H. Clapp were similarly determined by him. They are possibly the same as the "*Vallonia gracilicosta* var., close to *costata* Say," of Squyer's list.

Vitrea radiatula Alder (*hammonis* Ström.).

Zonitoides arborea (Say).

Zonitoides minuscula (Binney).

Euconulus fulvus (Müller).

Agriolimax campestris (Binney) var. The specimens are perhaps to be referred to the *montanus* of Ingersoll.

Pyramidula cronkhitei (Newcomb) near var. *anthonyi* Pilsbry.

Succinea avara Say.

Succinea retusa Lea.

Lymnæa palustris Müller.

Lymnæa caperata Say.

Planorbis parvus Say.

Planorbis callioglyptus Vanatta. A single poor specimen is thus somewhat doubtfully determined.

Planorbis antrosus Conrad (= *bicarinatus* Say).

Physa gyrina Say var.

Aplexa hypnorum Linné.

CHARLES W. GRIPP.

Mr. Charles William Gripp, of San Diego, Cal., who is known to many of our readers as an indefatigable collector and generous correspondent in exchanges of shells, died as the result of a serious accident in San Diego, January 3, 1913, aged forty-eight years. Mr. Gripp was a native of Odeshög, Sweden, and came to America in 1875. Since 1897 he was connected with the San Diego post-office. The funeral services were held on January 6th, and interment was made in the Masonic cemetery. Mr. Gripp was especially interested in the Californian, and particularly the San Diego mollusk fauna. He made various excursions beyond the entrance of the harbor in fishing boats, and by careful attention to bottom samples and stones brought up on the roots of kelp, discovered in this way a number of new species and increased our knowledge of the old ones. His collection was very complete in his specialty, and arranged and labeled with that neatness and care more common with European than American collectors. Personally he was agreeable, generous and considerate, and will be much regretted by his correspondents. He left a widow, who was, at the time of his death, seriously ill in a hospital, thereby adding to the pathos of the circumstances under which he died. W. H. D.

NOTES.

Cannibal Snail.—A carnivorous mollusk called *Glandina* was imported into France from Mexico some months ago, and at a meeting of the Academy of Sciences in Paris, Professor Bouvier asserted that its cannibal propensities were conferring immense benefit on market gardeners.

Snails commit serious depredations on French cabbages and salads, but since the arrival of the *Glandina*, which lives on snails and slugs, the latter have largely decreased, and agriculturalists are able to send their produce to the Halles in perfect condition.

When the Mexican mollusk encounters a French snail, it thrusts forward its head, and, penetrating to the innermost recesses of the enemy's shell, devours its prey alive. The eggs of the *Glandina* are as large as those of some birds.

The experiments conducted with a hundred giant snails from Mexico have proved so satisfactory that arrangements are being made to supply 2000 more to the market gardeners round Paris.—*Daily Mail.*

The Nautilus.

Vol. XXVI. APRIL, 1913. No. 12

NOTES ON A FEW BRITISH COLUMBIA MARINE SHELLS.

BY A. W. HANHAM, DUNCAN, B. C.

The following notes do not refer necessarily to rare, or even uncommon species and are in some cases only a comparison of common species occurring at different localities, sometimes not very widely separated.

Victoria is on the south end of Vancouver Island, inside the Straits of Juan de Fuca; Maple Bay and Departure Bay lie on the East or inside coast of Vancouver Island, the former South of Departure Bay and distant about 30 miles; both are sheltered or more or less land-locked; Salt Spring Island lies along outside the entrance to Maple Bay, and in the channel outside this entrance the water is very deep.

Terebratella transversa Sowerby. This species is common on the underside of rocks at Maple Bay, and may be found not far below high water mark where the shore shelves steeply and the rocks are well piled up.

Pecten hastatus Sowerby and *Pecten rubidus* Hinds. These species occur together, and in the channels off Nanaimo and Departure Bay I have seen them brought up in the dredge by the pailful. Both are sometimes coated, even on both valves, with a native sponge, bright yellow in color when living, named *Myxitis parasitica*, Lambe.

Hinnites giganteus Gray. Very fine specimens may be taken among the rocks at Maple Bay at low tides, but the big fellows are hard to detach without breaking or chipping the lower valve. A nicely grown specimen makes a good ash-tray.

Axinaea septentrionalis Midd. Reported by the late Rev. G. W. Taylor as being not uncommon on the Northern and Western Coasts of Vancouver Island, but not observed by him at Victoria or Departure Bay, so that I was all the more surprised this summer, to pick up a specimen in good condition at Maple Bay lying on the beach near high water mark. Through the kindness of Mr. Taylor I possess a good set of the species from Northern waters.

Psammobia rubroradiata Nuttall. The late Rev. G. W. Taylor in his List of the Marine Mollusca of the Pacific Coast of Canada, refers to this as being a rare species, stating that he had not seen living specimens. At Maple Bay this appears to be one of the species occurring there in numbers, broken shells are in abundance, and quite good ones are often washed up, many still retaining the animal, and of quite a fair size. This shell seen lying in the water is rather a beauty, but in the cabinet a very ordinary shell, as the color seems largely to disappear.

Semele rubropicta Dall. Quite a rare species at Maple Bay, and while it might be captured by dredging, half a dozen specimens or odd valves represent the catch for almost as many seasons.

Volutella pyriformis Carpenter. This is a little beauty, and while Taylor refers to it as a common species, I have only been fortunate enough,—with the exception of a tiny juv. specimen at Maple Bay, —to find it once, and that was at Foul Bay, near Victoria; here under a stone in a cavity under a large boulder, I secured a small colony.

Purpura lima Martyn. (= *canaliculata* Duclos). This is the only Purpura that I have taken in any of our inside coast waters since leaving Victoria; there, *P. crispata*, Chemnitz, and *saxicola*, Val., occurred in perhaps equal numbers with Lima: this absence seems rather strange. The *P. lima* of Maple Bay is very common, a large shell, solidly built, heavy, and sometimes brightly colored and banded.

Cerostoma foliatum Martyn. I took only two specimens of Cerostoma all the time I lived in Victoria, and those at extreme low tide; at Maple Bay this is a fairly common species among the rocks between tides, and juv. specimens are quite abundant.

Eulima micans Carpenter. This species is usually only to be obtained in our Northern waters by dredging in from ten to forty fathoms, but I have met with it rarely at Maple Bay at extreme

low tide buried in a sponge on the underside of large flat rocks on which this sponge sometimes grows in masses.

Bittum filosum Gould. The species is abundant at Maple Bay on rocks at low tides among the sea weeds, and many of the living shells are white in colour.

Isapis obtusa Cpr. Since my notes on this species in the February Number of Vol. xxlv, a further search the following August in the same locality, showed the shell to be quite plentiful on its same host, *Tapes staminea* Conrad.

Natica clausa B. & S. and *Lunatia pallida* B. & S. A few small specimens, living and dead, fell to my bag on one of my few dredgings in Departure Bay.

Velutina laevigata Linn. At Maple Bay very occasional at extreme low tides among the rocks, but not, I believe, a rare species when dredging in from 10 to 30 fathoms.

Acmaea patina Esch. This and allied species are of course plentiful both at Victoria and on all our rocky coasts, but none of the shells found in our inside waters are of any size, miserable specimens when compared with those taken at Victoria, from which locality I have some beauties.

Acmaea mitra Esch. The same remarks apply to this species, except that at Maple Bay it is in addition quite a rare shell, and those taken have been very small and depauperate-looking, while in Departure Bay I have not yet met with it.

Fissuridea aspera Esch. This is one of our common species at Maple Bay between tides, and generous in size, like those I have from near Victoria, and certainly much more abundant than there.

Leptothyra bacula Carpenter. A rare species, I believe, in northern waters, and the only specimen that I have taken—except on Southern California beaches—was a small one out of a root of kelp washed up after a storm. This root was still attached to some very large barnacles, and was rather a "gold mine" for small species, as *Alvanias*, etc.

Margarita lirulata Carpenter. At Maple Bay by far our most abundant "top;" swarms in certain seaweeds on the rocks at extreme low tides, and stray specimens may even be taken on the under side of stones between tides, associated with small *Littorina*. As Taylor has pointed out, this is a very variable species in our waters.

Margarita helicina Fab. Reported not uncommon between tides around Victoria; not yet discovered at Maple Bay.

Solariella peramabilis Carpenter and *Solariella cidaris* Carpenter. I have a few nice examples from dredgings in or near Departure Bay, the latter though, only from channels outside the bay.

Note. While I have been able to do much more collecting at Maple Bay than at Departure Bay, dredging has as yet been attempted only at the latter locality.

"Errata." NAUTILUS, Vol. xxiv, page 114, line 10, read "tiny" instead of "fine."

SPHÆRIIDÆ, OLD AND NEW, IV.

BY V. STERKI.

Pisidium overi, n. sp.—Mussel rather small, oblique, with the anterior part somewhat longer, medium to well inflated; superior margin short, nearly straight, posterior subtruncate to rounded, inferior well curved; supero-anterior slope beginning with an angle, close to beaks, rather long, straight to slightly curved; anterior end situated well below the median line, rounded angular; beaks rather small, rounded; surface dullish to somewhat glossy, with very fine subregular striæ; shell thin, translucent, nearly colorless to pale corneous; hinge medium stout, plate narrow; cardinal teeth rather long, moderately curved to nearly straight, the right not, or little thicker in its posterior part, left anterior with its apex rounded to somewhat pointed in old specimens; ligament and resilium rather short and slight.

Long. 3.5, alt. 3, diam. 2.4 mm. (100 : 91 : 68).

Long. 3.3, alt. 2.7, diam. 2.1 mm. (100 : 82 : 64).

Hab.: Clear Lake, Deuel Co., S. Dakota, collected by Mr. W. H. Over, in several lots, in 1906 and 1908; types No. 5752 and corresponding with those in Mr. Over's collection, in whose honor the species was named (in 1906). Apparently the same *Pisidium* is from Cannon Lake, Rice Co., Minnesota, collected by Mr. L. E. Daniels in 1905 (No. 5595); the two specimens in the collection are somewhat larger and more inflated, with the beaks higher.

P. overi somewhat resembles *walkeri* St. in shape and appearance,

but is smaller, the anterior part of the mussel is much smaller, and the cardinal teeth are markedly different, as seen in quite a number of specimens.

Pisidium nevadense, n. sp.—Mussel inequipartite, oblique, higher in front of the beaks than behind, medium inflated; superior and posterior margins little curved with a rounded—somewhat over right—angle between them, inferior rather long, somewhat descending toward the rounded anterior end, supero-anterior slope somewhat curved; beaks nearer the posterior end, medium large, rounded, slightly prominent over the hinge margin; scutum and scutellum barely marked; surface somewhat shining, with slight irregular striæ and a few slight, impressed lines of growth; color: straw, shell subtranslucent, rather strong; hinge curved, medium stout, plate rather broad; cardinal teeth rather large, the right with the anterior part somewhat thickened, the posterior much thicker, with a deep groove, nearly bifid, left anterior long, angular-curved, posterior long curved, oblique, extending forward well beyond the apex of the anterior; laminæ little projecting inward, cusps short, pointed, especially those of the left valve; ligament and resilium rather long, moderately strong. Long. 6, alt. 4.8, diam. 3.3 mm. (100 : 80 : 55).

Hab.: Nevada Co., California; types no. 56278 in the National Museum, and my collection no. 2726; of the former, one specimen is 6.4 mm. long. For my specimens I am indebted to the late R. E. C. Stearns, being identical with those of the National Museum, which were received from (and collected by ?) J. J. Rivers. Evidently distinct from all described species, but the specimens were shelved in the hope of getting additional material. This *Pisidium* has the best-formed cardinal teeth of any seen.

P. (*nevadense* var., or sp.?) *modicum*, n.—Mussel of similar appearance, but smaller, less oblique, not higher in front of the beaks than behind, beaks somewhat broader, color more whitish; long. 4.3, alt. 3.5, diam. 2.6 mm.

Hab.: White Pine, Nevada. Specimens are in the collection of Mr. B. Walker, from the Wetherby collection, and my collection no. 5813, part of the original lot; more are probably in other collections.

P. glabellum, n. sp.—Mussel small, somewhat oblique, rather well inflated; superior margin nearly straight, bounded by angles in front and behind, supero-anterior slope nearly straight, anterior end

rounded, posterior and inferior margins forming one curve; beaks slightly posterior, rounded or somewhat flattened on top, moderately prominent; surface polished, with very fine, slight striæ, color corneous, occasionally with a slight reddish tinge, to clear amber, shell transparent to translucent; hinge stout, plate rather broad; right cardinal curved, its posterior part thicker, left anterior with apex pointed, posterior rather short, oblique, well curved; ligament and resilium short, strong.

Long. 2.2, alt. 1.9, diam. 1.4 mm. (100 : 87 : 64).

Hab.: New England to Pennsylvania and Minnesota. Types No. 324 from Hess Lake, Newaygo Co., Mich., collected by Mr. L. H. Streng, received in 1895, then regarded as distinct and named. No. 78977 of the Philadelphia Acad. Nat. Sciences, from Point Pleasant, N. J. (F. H. Brown, Aug., 1890), conform with the types. From various places there are specimens of nearly the same size and appearance, but more or less different in shape: superior margin more curved, beaks narrower and more elevated, the anterior end more angular, e. g. from Graham's Creek, Nepean, Ont., collected by Justice F. R. Latchford in 1911, No. 6812. They appear to represent a variety.

P. glabellum has some resemblance to small forms of *P. splendidulum* St.; the latter is widely distributed, common and very variable, but from all its forms, *glabellum* proved distinct. The hinge is markedly strong and somewhat like that of *P. variabile* and *compressum;* the mussel is somewhat like a miniature *P. variabile*, and it appears to range with that group.

P. lermondi, n. sp.—Mussel rather small, subequipártite; outlines nearly elliptical along the valve edges, well inflated around the beaks, impressed along the margins, with cutting edges; beaks a little posterior, rather small, somewhat prominent over the margin; surface polished, with shallow, irregular striæ; color deep corneous to somewhat olivaceous, shell transparent to translucent, thin; hinge slight, plate quite narrow; cardinal teeth little curved, the right with its posterior end slightly thicker, left posterior short, quite above the anterior; laminæ rather short and slight, cusps in the left valve pointed; ligament and resilium slight.

Long. 3.4, alt. 3, diam. 2.5 mm. (100 : 88 : 73).

Hab.: Duck Pond, Warren, Me., collected by Mr. N. W. Lermond (in whose honor the species was named), Sept., 1909. Types No.

6364, and in the collection of Mr. Lermond. Also at other places in the vicinity of Warren, Me.

The shape of the mussel is quite peculiar, especially in front or rear aspect, and if there were only one or a few specimens, they might be regarded as abnormal. Some forms of *P. medianum* St. are of somewhat similar shape, in lateral view, but they are of light color, the shell is not transparent, the surface dullish, and the hinge is quite different. Young and half-grown specimens may be mistaken for *P. splendidulum*.

Sphærium acuminatum (Pme.), *Cyclas acuminata* Prime, Proc. Boston Soc. Nat. Hist., iv, p. 155, 1851, was made a synonym of *S. striatinum* (Lam.) in Mon. Corb., p. 37, by Prime, but it is distinct, as evidenced by the specimens (a few valves) in the Prime collection.[1]

Cyclas albula Pme., l. c., is probably the same, to judge from the specimens—two poor, washed left valves in the Boston Society Natural History (No. 8086)—but not from the description. There are no specimens of *C. albula* in the Prime collection (Mus. Comp. Zool.), and for the above reasons it appears proper to use the name *acuminata*, though *albula* has precedence, yet both are on the same page.

The species is common in the great-lake region, and rather variable.

S. ohioense n. sp.—Mussel somewhat inequipartite and elongate, moderately inflated, the valves rather flat toward the margins, superior and inferior margins slightly curved, anterior margin rounded or obliquely subtruncate above, posterior obliquely truncate to subtruncate, end rounded; beaks small, slightly elevated; scutum and scutellum rather long, narrow; surface dull, with rather fine striæ—or sulci; color straw to yellowish corneous to grayish in old specimens; shell rather thin, hinge slight, plate narrow; cardinal teeth small, short, posterior part of the right thick, nearly cubical, laminæ slight; ligament and resilium rather long, slight.

Long. 11.5, alt. 8.2, diam. 5.3 (100 : 73 : 45).

Soft parts not examined.

Hab.; Ohio River and tributary. At Cincinnati on Nov. 1, 1895, I collected a good number of specimens, young to adult, in com-

[1] "Origin," probably from Lake Superior, also two valves in the Boston Society of Natural History (No. 8085).

pany with *S. striatinum* and *stamineum*. Types, No. 1172. Ohio River at Cannelton, Ind., collected by Mr. L. E. Daniels, Sept. 23, 1904; Elk River at Shelton, Clay co., W. Va., collected by Dr. A. E. Ortmann, July 10, 1911. *A. Sphaerium* almost exactly like these, and apparently identical, was collected at Joliet, Ill., by Mr. Jas. H. Ferriss, in 1896, (No. 1354).

Musculium orbiculare n. sp.—Mussel rather small, inequipartite, nearly circular in outlines with the postero-inferior part slightly extended, moderately and evenly inflated; beaks quite small, narrow, little prominent, calyculate, somewhat inclined forward; surface somewhat shining, smooth, with slight, microscopic striæ; color pale corneous to grayish with narrow straw colored zones along the margins: shell thin, hinge slight, cardinals small, the right angular with the posterior part about as thick as long; the whole tooth is little projecting, inclosing a triangular excavation for the left anterior; laminæ rather long, slender; ligament and resilium rather long, slight;

Long. 83, alt. 7.5, diam. 3.8 (100 : 90 : 48).

Hab.: Ill., Ind.; Athens, Ill., Nason collection in the Illinois State Lab. Nat. History seen in 1911. Types No. 6796. Hammond, Ind., collected by Mr. L. E. Daniels, Aug. 3, 1904 (No. 4400).

This species is markedly different from all others seen and appears to be distinct. It has some resemblance to *M. elevatum* Hald., but is much smaller and slighter, less inflated, the beaks are much smaller, and it differs from half grown *elevatum* of about the same size more than it does from adults.

There are still large numbers of *Sphæriidæ* on hand, of all three genera, which cannot be referred to any of the described species. Yet it is precarious to establish new species on them, considering the great variation of many species. On the other hand, haphazard lumping would do more harm then good, and is unscientific. More good material from all over the continent is desirable, and will help ascertaining their real relations. Also a large number of well marked varieties and subspecies, partly very interesting in several respects, systematically and geographically, are on file, and will be published later.

SOME CRITICISMS ON DR. F. HAAS' MONOGRAPH OF THE UNIONIDÆ.

BY L. S. FRIERSON.

The well known Monograph of Unionidæ of H. C. Kuster is now being continued by Dr. F. Haas, and 35 plates of fine figures have been received. Dr. Haas is a prolific maker of new genera, he having added already 14 new ones to the tremendous total, although three plates only deal with the groups of eastern Asia. The new genera are fully as heterogeneous as are many of those of his predecessors.

Shells so close as *Nodularia jourdyi* Morelet and *douglasiæ* Greg. are widely separated, and yet such diverse material as *jourdyi* and *asperula* Lea are placed together. The *Physunio crossei* and *Nodularia rusticus*, by Simpson's classification, are placed by Dr. Haas in the same genus, etc., etc.

The specific determinations are equally open to criticism. A few of the more striking are as follows: On plate 33, figs. 7, 8 and 9 are given as *P. micropterus* Morelet, and *Unio pulcher* Tapperone-Canefri is given as a synonym. While these species bear considerable resemblance in their external facies, yet they are so diverse as to warrant their being placed in different genera, as was properly done by Mr. C. T. Simpson. *P. micropterus* has very small but well-defined beak cavities, while *pulcher* has none. *L. pulcher* has a row of several dorsal muscle scars, while *micropterus* shows none (*i. e.*, hidden beneath the interdention). More especially, *micropterus* has a "third muscle scar" over the anterior adductor, not shown by *pulcher*, and the latter has the anterior adductor and protractor scars *widely separate* (which is shown in Haas' figure), while in *micropterus* they are confluent. Other differences might be noted, the whole forcing us to place *micropterus* in *Physunio*, while the *pulcher* belongs to *Lamellidens*, as Mr. C. T. Simpson properly placed them.

Figures 11 and 12, plate 31 are said to be figures of the *Harmandia somboriensis* Rochebrune. If these figures are from authentic specimens, then *Harmandia* must lose its place as a genus, for figures 11 and 12, are young *Hyria*, from South America. Even more singular is the treatment of the genus *Trapezoideus*, on plates 32 and 33. It is impossible that the sundry figures given as *Trapezoideus*

misellus Morelet can represent a single species. If Fig. 4, plate 33, is not an abnormal or diseased shell it must be a *Solenaia*, probably the *Solenaia* (*Mycetopus*) *rugatus* Sowerby.

The figures 3, and 4, plate 32, are equally impossible. They are supposed to be *Trapezoideus foliaceus*, Gould, but are really an undescribed shell, not far removed from the *N. dimotus* Lea (figured in the work, on plate 19). I have had for several years, a set of these, purchased, with a label "*Trapezoideus foliaceus*, Gould, Ex. Berlin Museum, *Birma*," and possibly a part of the same lot figured by Dr. Haas. Not being able to match them with Gould's species, I submitted them to Dr. Dall, who placed them near to *dimotus*, I, therefore propose for the species as figured by Haas, (fig. 3, and 4, plate 32) the following name.

PARREYSIA DALLIANA n. sp.

Shell small, rather thick, somewhat ovate. Length (of a medium specimen) 4.5; height (near end of ligament) 2.7; Diameter 2, mm. Rounded before, basal and dorsal lines nearly straight, post basal point rounded. Epidermis olivaceus, greenish in young shells, brownish in old specimens, and frequently *bleached*. Beaks very small, pointed, low, and incurved, with fine zigzag radial sculpture, which extends a little down the shell, and especially down the post slope; in dirty, or eroded specimens hardly to be noticed. The shell is a little inflated, the post ridge rounded. Inside the nacre is soft cream or white. The left valve has two cardinals, the anterior blade-like, and the posterior much smaller, and tubercular. The laterals are double (or even inclined to be trifid); in the right valve the cardinals are split or double, and the laterals single.

The anterior cicatrices are confluent, and also those of the posterior.

Habitat is stated to be "Birma."

The shell is stouter, shorter and more inflated than *P. foliaceus*, and the posterior portion is not as green. The species lacks the sculpturing of *P. dimotus*, and the direction of the cardinal teeth is different. The shell is a *Parreysia* according to Simpson's ideas of classification.

SHELLS COLLECTED AT MANZANILLO, WEST MEXICO, OCT., 1910, BY C. R. ORCUTT, IDENTIFIED BY WILLIAM H. DALL.

Gadinia peruviana Sby.
Siphonaria lecanium Cpr.
Bulla punctulata Ads.
Terebra variegata Gray.
Conus princeps Brod.
Conus arcuatus Sby.
Conus nux Brod.
Conus gladiator Brod.
Mitra funiculata Rve.
Vasum cestum Brod.
Lyria barnesii Gray.
Latirus ceratus Wood.
Purpura triserialis Blv.
Cantharus sanguinolentus Ducl.
Nassa corpulenta C. B. Ads.
Columbella major Sby.
Columbella festiva Sby.
Anachis coronata Sby.
Anachis sulcosa Sby.
Nitidella cribaria Lam.
Muricidea dubia Sby.
Craspedotriton scalariformis Brod.
Morum tuberculosum Sby.
Cypraea arabicula Lam.
Trivia pacifica Gray.
Strombus gracilior.
Cerithium maculosum Kien.
Cerithium uncinatum Gmelin.
Cerithium gemmatum Hinds.
Planaxis nigritella Forbes.
Planaxis nigritella var.
Littorina conspersa Phil.
Hipponix grayanus Mke.
Crepidula onyx Sby.
Calyptraea mammillaris Brod.
Trochita spirata Forbes.
Crucibulum imbricatum Brod.
Natica chemnitzianum Pfr.
Polinices uber Val.
Scurria mesoleuca Mke.
Turbo squamigerus Rve.
Uvanilla unguis Mawe.
Tegula globula Cpr.
Nerita bernhardi Recl.
Fissurella alba Cpr.
Fissurella microtrema Sby.
Fissurella rugosa Sby.
Chiton albolineatus Sby.
Arca grandis B. & S.
Perna chemnitziana Orb.
Lima pacifica Orb.
Phacoides lampsus Dall.
Cardium consors Sby.
Cardium magnificum Desh.
Venericardia crassicostata Sby.
Venericardia cuvieri Brod.
Anomalocardia subimbricata Sby.
Chione undatella Sby.
Paradione squalida Sby.
Tellina cumingi Hanley.
Tellinides viridotincta Cpr.
Semele proxima C. B. Ads.
Tagelus politus Cpr.
Thracia plicata Desh.

The above list has a value for geographical distribution, which justifies its publication.—W. H. Dall.

BOSTON MALACOLOGICAL CLUB.

The November meeting was devoted to reports by members on the work done during the summer, collecting or otherwise. At the December meeting Mr. C. J. Maynard gave an account of the distribution and variation of the species of *Cerion* in the vicinity of Nassau, New Providence, Bahama Islands, showing a large series of specimens and illustrating with the aid of a map how the various species and varieties were distributed from what he considered the original starting point. At the January meeting the Secretary gave an account of the various forms of the foot in mollusca, and its adaptation to the habits of the species. Mr. Olof O. Nylander, of Caribou, Maine, spoke at the meeting, February 10, on the distribution and variation of *Lymnæa emarginata* Say and the var. *mighelsi* W. G. Binney, in Aroostook Co., Me. A large map, on which was marked the various colonies, clearly showed its limited distribution throughout the Fish River lakes, and a beautiful series used in illustrating his paper published in 1901, with many additional specimens, showed all the gradations between *L. emarginata* and *mighelsi*. The speaker referred to the great abundance of the species in some years and their scarcity in others, and the probable causes; also of the changes which are now taking place in the region detrimental to their existence.

HENRY W. WINKLEY, *Secretary*.

NOTES.

A NEW FORM OF OREOHELIX.—*Oreohelix haydeni betheli*, n. subsp. Typically flat and broad, with a lower spire and wider umbilicus than true (Utah) *O. haydeni*; last embryonic whorl concave or even grooved next the suture. The spiral cords are strong throughout, often more numerous than in typical *haydeni*, 10 to 16; fine interstitial spirals visible. Alt. 10, diam. 21 mm. Abundant under rocks on north side of Grand River at Glenwood Springs, Colorado, collected by E. Bethel, J. Henderson, T. D. A. Cockerell and others. The true *betheli* occurs on the north side of the river, but on the opposite side is a related form (*alta* Pilsbry, n. v.) with the spire conoidal and the umbilicus narrower, and also other forms to be described later.—H. A. PILSBRY and T. D. A. COCKERELL.

Mr. C. J. Maynard is now in the Bahamas making further studies on the distribution of the various species of *Cerion*.

MBL/WHOI LIBRARY
WH 17WS V

297